Name: _____ ( ) Class: _____

PRELIMINARY EXAMINATION
GENERAL CERTIFICATE OF EDUCATION ORDINARY LEVEL

ADDITIONAL MATHEMATICS 4047/01

Paper 1 **Thursday 16 August 2018**

2 hours

Additional Materials: Answer Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class, and index number on all the work you hand in.
Write in dark blue or black pen on both sides of the paper.
You may use a pencil for any diagrams or graphs.
Do not use paper clips, highlighters, glue, or correction fluid.

Answer **all** the questions.
Write your answers on the separate Answer Paper provided
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, staple all your work together with this cover sheet.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is **80**.

FOR EXAMINER'S USE

Q1		Q6		Q11	
Q2		Q7			
Q3		Q8			
Q4		Q9			
Q5		Q10			

/ 80

This document consists of **5** printed pages.

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + ... + \binom{n}{r}a^{n-r}b^r + ... + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1).....(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$

$$\sin(A \pm B) = \sin A\cos B \pm \cos A\sin B$$

$$\cos(A \pm B) = \cos A\cos B \mp \sin A\sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A\tan B}$$

$$\sin 2A = 2\sin A\cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 Express $\dfrac{3x^3+2x^2+4x-1}{x^3+x^2}$ in partial fractions. [4]

2 A cylinder has a radius of $(1+2\sqrt{2})$ cm and its volume is $\pi(84+21\sqrt{2})\,\text{cm}^3$.
Find, **without using a calculator**, the exact length of the height of the cylinder in the form $(a+b\sqrt{2})$ cm, where *a* and *b* are integers. [5]

3 **(i)** Sketch the graph of $y=4-3\sin 2x$ for $0\le x\le \pi$. [3]

(ii) State the range of values of k for which $4-3\sin 2x=k$ has two roots for $0\le x\le \pi$. [2]

4 **Solutions to this question by accurate drawing will not be accepted.**

PQRS is a parallelogram in which the coordinates of the points *P* and *R* are $(-5, 8)$ and $(6, -2)$ respectively. Given that *PQ* is perpendicular to the line $y=-\dfrac{1}{2}x+3$ and *QR* is parallel to the *x* axis, find

(i) the coordinates of *Q* and of *S*, [5]

(ii) the area of *PQRS*. [2]

5 **(i)** Differentiate $\dfrac{\ln x}{x}$ with respect to *x*. [3]

(ii) Hence find $\int \dfrac{\ln x^2}{x^2}\,\text{d}x$. [4]


Downloaded from SingaKids.com

6 **(i)** Show that $\dfrac{2}{\tan\theta + \cot\theta} = \sin 2\theta$. [3]

(ii) Hence find the value of p, giving your answer in terms of π, for which

$\int_0^p \dfrac{4}{\tan 2x + \cot 2x}\,\mathrm{d}x = \dfrac{1}{4}$, where $0 < p < \dfrac{\pi}{4}$. [4]

7



In the diagram XBY is a structure consisting of a beam XB of length 35 cm attached at B to another beam BY of length 80 cm so that angle $XBY = 90^\circ$. Small rings at X and Y enable X to move along the vertical wire AP and Y to move along the vertical wire CQ. There is another ring at B that allows B to move along the horizontal line AC. Angle $ABX = \theta$ and θ can vary.

(i) Show that $AC = (35\cos\theta + 80\sin\theta)$ cm. [2]

(ii) Express AC in the form of $R\sin(\theta + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. [4]

(iii) Tom claims that the length of AC is 89cm. Without measuring, Mary said that this was not possible. Explain how Mary came to this conclusion. [1]

8 **(a)** Find the range of values of p for which $px^2 + 4x + p > 3$ for all real values of x. [5]

(b) Find the range of values of k for which the line $5y = k - x$ does not intersect the curve $5x^2 + 5xy + 4 = 0$. [5]

Downloaded from SingaKids.com


9 The diagram shows part of the graph of $y = 4 - |x + 1|$.

(i) Find the coordinates of the points *A*, *B*, *C* and *D*. [5]


(ii) Find the number of solutions of the equation $4 - |x + 1| = mx + 3$ when

(a) $m = 2$ **(b)** $m = -1$ [2]

(iii) State the range of values of *m* for which the equation $4 - |x + 1| = mx + 3$ has two solutions. [1]



10 The diagram shows a cone of radius *r* cm and height *h* cm. It is given that the volume of the cone is 10π cm^3.



(i) Show that the curved surface area, *A* cm^2, of the cone, is $A = \dfrac{\pi\sqrt{r^6 + 900}}{r}$. [3]

(ii) Given that *r* can vary, find the value of *r* for which *A* has a stationary value. [4]

(iii) Determine whether this value of *A* is a maximum or a minimum. [2]

11 The equation of a curve is $y = x(2 - x)^3$.

(i) Find the range of values of *x* for which *y* is an increasing function. [5]

(ii) Find the coordinates of the stationary points of the curve. [3]

(iii) Hence, sketch the graph of $y = x(2 - x)^3$. [3]

Downloaded from SingaKids.com

St Nicholas Girls School Additional Mathematics Preliminary Examination Paper I 2018


Answers

Paper 1

1. $3+\frac{5}{x}-\frac{1}{x^2}-\frac{6}{x+1}$

2. $(12-3\sqrt{2}\ )$ cm

3 (i)



(ii) $1<k<4$ or $4<k<7$

4 (i) $Q(-10,\ -2)$, $S(11,\ 8)$ (ii) 160 units^2

5 (i) $\frac{1-\ln x}{x^2}$ (ii) $2\left(-\frac{1}{x}-\frac{\ln x}{x}\right)+c$

6 (ii) $\frac{\pi}{12}$

7 (ii) $5\sqrt{305}\sin(\theta+23.6^\circ)$ cm or $87.3\sin(\theta+23.6^\circ)$ cm


(iii) The maximum value of AC=87.3cm <89 cm

8 (a) $p>4$ (b) $-8<k<8$

9 (i) $A(-5,0)$, B$(-1,4)$, $C(3,0)$, D$(0,3)$ (ii) (a) 1 (b) infinite (iii) 2

10 (ii) 2.77 (iii) minimum

11 (i) $x<\frac{1}{2}$ (ii) $(2,0)$ $\left(\frac{1}{2},\frac{27}{16}\right)$ (iii)



Downloaded from SingaKids.com

Name: _____ ( ) Class: _____

PRELIMINARY EXAMINATION
GENERAL CERTIFICATE OF EDUCATION ORDINARY LEVEL

ADDITIONAL MATHEMATICS 4047/01

Paper 1 **Marking Scheme** **Thursday 16 August 2018**

2 hours

Additional Materials: Answer Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class, and index number on all the work you hand in.
Write in dark blue or black pen on both sides of the paper.
You may use a pencil for any diagrams or graphs.
Do not use paper clips, highlighters, glue, or correction fluid.

Answer **all** the questions.
Write your answers on the separate Answer Paper provided
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, staple all your work together with this cover sheet.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is **80**.

FOR EXAMINER'S USE

Q1		Q6		Q11	
Q2		Q7			
Q3		Q8			
Q4		Q9			
Q5		Q10			

80

This document consists of **5** printed pages.

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1).....(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$

$$\sin(A \pm B) = \sin A\cos B \pm \cos A\sin B$$

$$\cos(A \pm B) = \cos A\cos B \mp \sin A\sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A\tan B}$$

$$\sin 2A = 2\sin A\cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 Express $\frac{3x^3+2x^2+4x-1}{x^3+x^2}$ in partial fractions. [4]

1 $x^3+x^2 \overline{)\,3x^3+2x^2+4x-1}$ $3x^3+3x^2$ $-x^2+4x-1$ $\frac{3x^3+2x^2+4x-1}{x^2+x^3}=3+\frac{-x^2+4x-1}{x^2(x+1)}$ $\frac{-x^2+4x-1}{x^2(x+1)}=\frac{A}{x}+\frac{B}{x^2}+\frac{c}{x+1}$ $-x^2+4x-1=Ax(x+1)+B(x+1)+cx^2$ Let $x=-1$ $\quad -1-4-1=c$ $c=-6$ Let $x=0$ $\quad B=-1$ $-x^2+4x-1=Ax(x+1)-1(x+1)-6x^2$ Let $x=1$ $\quad -1+4-1=2A-2-6$ $A=5$ $\frac{3x^3+2x^2+4x-1}{x^2+x^3}=3+\frac{5}{x}-\frac{1}{x^2}-\frac{6}{x+1}$ **[4]**	M1√ M1√ M1√ A1
If • $\frac{3x^3+2x^2+4x-1}{x^2+x^3}=\frac{A}{x}+\frac{B}{x^2}+\frac{c}{x+1}$ • $\frac{3x^3+2x^2+4x-1}{x^2+x^3}=3+\frac{Ax+B}{x^2}+\frac{c}{x+1}$ • $\frac{3x^3+2x^2+4x-1}{x^2+x^3}=\frac{Ax+B}{x^2}+\frac{c}{x+1}$	Max 3m 3m 2m

SGFREEPAPERS.COM


Downloaded from SingaKids.com

2 A cylinder has a radius of $(1+2\sqrt{2})$ cm and its volume is $\pi(84+21\sqrt{2})$ cm^3.
Find, **without using a calculator**, the exact length of the height of the cylinder in the form $(a+b\sqrt{2})$ cm, where a and b are integers. [5]

2.	$\pi(84+21\sqrt{2}) = \pi(1+2\sqrt{2})^2 \times h$ $h = \dfrac{84+21\sqrt{2}}{(1+2\sqrt{2})^2}$ $h = \dfrac{84+21\sqrt{2}}{1+4\sqrt{2}+8}$ $h = \dfrac{(84+21\sqrt{2})(4\sqrt{2}-9)}{(4\sqrt{2}+9)(4\sqrt{2}-9)}$ $h = \dfrac{756-336\sqrt{2}+189\sqrt{2}-168}{81-32}$ $h = \dfrac{588-147\sqrt{2}}{49}$ $h = (12-3\sqrt{2})$ cm **[5]**	B1 M1 M1√ M1√ A1	expansion Conjugate surd For either expansion No unit, overall -1m

3 **(i)** Sketch the graph of $y = 4-3\sin 2x$ for $0 \le x \le \pi$. [3]

(ii) State the range of values of k for which $4-3\sin 2x = k$ has two roots for $0 \le x \le \pi$. [2]

3 (a)  [3]	B1 any one pt B1 2nd pt The 2 pts must be different nature B1 perfect	• sine shape • -ve sine shape • 1 cycle • Amplitude • shift +4 up ignoring no labelling of axes	
3 (b) $1 < k < 4$ or $4 < k < 7$ [2] **Alternative** **[5]** $1 < k < 7, k \ne 4$	B1 + B1 B1 + B1		

Downloaded from SingaKids.com

4 Solutions to this question by accurate drawing will not be accepted.

PQRS is a parallelogram in which the coordinates of the points *P* and *R* are $(-5, 8)$ and $(6, -2)$ respectively. Given that *PQ* is perpendicular to the line $y = -\frac{1}{2}x + 3$ and *QR* is parallel to the *x* axis, find

(i) the coordinates of *Q* and of *S*, [5]

(ii) the area of *PQRS*. [2]

1(i)	Since *QR* parallel to the *x* axis, $y_Q = -2$.	B1
	Since *PQ* is perpendicular to the line $y = -\frac{1}{2}x + 3$, gradient of *PQ* = 2	B1 (⊥ gradient)
	$\frac{(-2)-(8)}{x_Q-(-5)} = 2$	M1
	$-10 = 2x_Q + 10$	
	$x_Q = -10$	
	$Q(-10, -2)$	A1
	Midpoint of *PR* = Midpoint of *QS* or by inspection	
	$\left(\frac{(-5)+(6)}{2}, \frac{(8)+(-2)}{2}\right) = \left(\frac{(-10)+x_s}{2}, \frac{(-2)+y_s}{2}\right)$	
	$1 = -10 + x_s$ $\quad 6 = -2 + y_s$	
	$x_s = 11$ $\quad y_s = 8$	
	$S(11, 8)$ [5]	B1
(ii)	Area of *PQRS*	
	$= \frac{1}{2}\begin{vmatrix} -5 & -10 & 6 & 11 & -5 \\ 8 & -2 & -2 & 8 & 8 \end{vmatrix}$	
	$= \frac{1}{2}\lvert(10 + 20 + 48 + 88) - (-80 - 12 - 22 - 40)\rvert$ or (5+11)(8+2)	√M1
	$= \frac{1}{2}\lvert 320 \rvert$ [2]	
	$= 160$ units^2 **[7]**	A1 no unit overall -1m

SGFREEPAPERS.COM

Downloaded from SingaKids.com

5 **(i)** Differentiate $\frac{\ln x}{x}$ with respect to x. [3]

(ii) Hence find $\int \frac{\ln x^2}{x^2}\,\mathrm{d}x$. [4]

(i)	$\frac{\mathrm{d}}{\mathrm{d}x}\left(\frac{\ln x}{x}\right) = \frac{x\left(\frac{1}{x}\right) - \ln x}{x^2}$ $= \frac{1-\ln x}{x^2}$ [3]	B1 +B1 A1	Either $v\frac{du}{dx}$ $oru\frac{dv}{dx}$ with the use of quotient rule /product rule perfect
(ii)	$\int \frac{1-\ln x}{x^2}\,\mathrm{d}x = \frac{\ln x}{x}$ $\int \frac{1}{x^2}\,\mathrm{d}x - \int \frac{\ln x}{x^2}\,\mathrm{d}x = \frac{\ln x}{x}$ $\int x^{-2}\,\mathrm{d}x - \frac{\ln x}{x} = \int \frac{\ln x}{x^2}\,\mathrm{d}x$ $\frac{x^{-1}}{-1} - \frac{\ln x}{x} = \int \frac{\ln x}{x^2}\,\mathrm{d}x$ $\int \frac{\ln x}{x^2}\,\mathrm{d}x = -\frac{1}{x} - \frac{\ln x}{x}$ $\int \frac{\ln x^2}{x^2}\,\mathrm{d}x = 2\int \frac{\ln x}{x^2}\,\mathrm{d}x$ $= 2\left(-\frac{1}{x} - \frac{\ln x}{x}\right) + c$ [4] **[7]**	M1 M1 B1 A1	Integration is the reverse process of differentiation Making $\int \frac{\ln x}{x^2}\,\mathrm{d}x$ the subject or split the expression Integration of x^{-2} With c

Downloaded from SingaKids.com

6 **(i)** Show that $\dfrac{2}{\tan\theta+\cot\theta}=\sin 2\theta$. [3]

(ii) Hence find the value of p, giving your answer in terms of π, for which


$\int_0^p \dfrac{4}{\tan 2x+\cot 2x}\,dx=\dfrac{1}{4}$, where $0<p<\dfrac{\pi}{4}$. [4]

(i)	$\dfrac{2}{\tan\theta+\cot\theta}=2\div\left(\dfrac{\sin\theta}{\cos\theta}+\dfrac{\cos\theta}{\sin\theta}\right)$ $=2\div\left(\dfrac{\sin^2\theta+\cos^2\theta}{\cos\theta\sin\theta}\right)$ $=2\div\left(\dfrac{1}{\cos\theta\sin\theta}\right)$ $=2\sin\theta\cos\theta$ $=\sin 2\theta$ [3]	B1 M1 M1	change to sin and cos combine terms for identity …to the end. (must show "1")
(ii)	$\int_0^p \dfrac{4}{\tan 2x+\cot 2x}\,dx$ $=2\int_0^p \sin 4x\,dx$ $=2\left[-\dfrac{\cos 4x}{4}\right]_0^p$ $=\left(-\dfrac{1}{2}\cos 4p\right)-\left(-\dfrac{1}{2}\cos 0\right)$ $=-\dfrac{1}{2}\cos 4p+\dfrac{1}{2}$ $\int_0^p \dfrac{4}{\tan 2x+\cot 2x}\,dx=\dfrac{1}{4}$ $-\dfrac{1}{2}\cos 4p+\dfrac{1}{2}=\dfrac{1}{4}$ $-\dfrac{1}{2}\cos 4p=-\dfrac{1}{4}$ $\cos 4p=\dfrac{1}{2}$ $4p=\dfrac{\pi}{3}$ $p=\dfrac{\pi}{12}$ [4] **[7]**	B1 M1 M1 A1	integrate their sinkx for substitution in their integral

SGFREEPAPERS.COM

Downloaded from SingaKids.com

7




In the diagram *XBY* is a structure consisting of a beam *XB* of length 35 cm attached at *B* to another beam *BY* of length 80 cm so that angle $XBY = 90^\circ$. Small rings at *X* and *Y* enable *X* to move along the vertical wire *AP* and Y to move along the vertical wire *CQ*. There is another ring at *B* that allows *B* to move along the horizontal line *AC*. Angle $ABX = \theta$ and θ can vary.

(i) Show that $AC = (35\cos\theta + 80\sin\theta)$ cm. [2]

(ii) Express *AC* in the form of $R\sin(\theta + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. [4]

(iii) Tom claims that the length of *AC* is 89cm. Without measuring, Mary said that this was not possible. Explain how Mary came to this conclusion. [1]

Downloaded from SingaKids.com

7 (i)	$AB = 35\cos\theta$ $\angle YBC = 90^\circ - \theta$ $\angle BYC = \theta$ $BC = 80\sin\theta$ $AC = (35\cos\theta + 80\sin\theta)$ cm  [2]	B1 either AB or BC B1 -1m overall for no unit
7 (ii)	$R\sin(\theta+\alpha) = R\sin\theta\cos\alpha + R\cos\theta\sin\alpha$ $AC = 35\cos\theta + 80\sin\theta$ $R\sin\alpha = 35$ $R\cos\alpha = 80$ $R^2\cos^2\alpha + R^2\sin^2\alpha = 80^2 + 35^2$ $R = \sqrt{80^2 + 35^2}$ $R^2 = 7625$ $R = 87.3$ *or* $5\sqrt{305}$ $\frac{R\sin\alpha}{R\cos\alpha} = \frac{35}{80}$ $\tan\alpha = \frac{35}{80}$ $\alpha = 23.6^\circ$ $AC = 35\cos\theta + 80\sin\theta$ $35\cos\theta + 80\sin\theta = 5\sqrt{305}\sin(\theta + 23.6^\circ)$ cm or $87.3\sin(\theta + 23.6^\circ)$ cm [4]	B1 M1 for R M1 for $\tan\alpha = \frac{35}{80}$ A1
7 (iii)	The maximum value of AC=87.3cm Therefore it is not possible for the length to be more than that. **Alternative** $5\sqrt{305}\sin(\theta + 23.6^\circ) = 89$ $\sin(\theta + 23.6^\circ) = \frac{89}{5\sqrt{305}}$ No Solution Therefore it is not possible for the length to be more than that. [1] **[7]**	DB1 DB1

SGFREEPAPERS.COM

Downloaded from SingaKids.com

8 **(a)** Find the range of values of p for which $px^2 + 4x + p > 3$ for all real values of x. [5]

(b) Find the range of values of k for which the line $5y = k - x$ does not intersect the curve $5x^2 + 5xy + 4 = 0$. [5]

(a)	$px^2 + 4x + p > 3$ for all real values of x $px^2 + 4x + p - 3 > 0$ for all real values of x, D<0 $4^2 - 4(p)(p-3) < 0$ $16 - 4p^2 + 12p < 0$ $4p^2 - 12p - 16 > 0$ $p^2 - 3p - 4 > 0$ $(p-4)(p+1) > 0$ $p < -1,\ p > 4$ NA As p >0 [5]		M1 M1 M1 DA1+DA1	D<0 with substitution For $b^2 - 4ac$ For factorisation Upon correct factorisation Ignore"and" and no p>0
(b)	$5y = k - x$ $5x^2 + 5xy + 4 = 0$ $5x^2 + 5x\left(\frac{k-x}{5}\right) + 4 = 0$ $5x^2 + kx - x^2 + 4 = 0$ $4x^2 + kx + 4 = 0$ $k^2 - 4(4)(4) < 0$	$5(k-5y)^2 + 5(k-5y)y + 4 = 0$ $5k^2 - 50ky + 125y^2 + 5ky - 25y^2 + 4 = 0$ $100y^2 - 45ky + 5k^2 + 4 = 0$ $(-45k)^2 - 400(5k^2 + 4) < 0$ $2025k^2 - 2000k^2 - 1600 < 0$ $k^2 - 64 < 0$ $(k-8)(k+8) < 0$	M1 M1 +M1√ M1	For substitution D<0 with substitution For $b^2 - 4ac$ factorisation
	$-8 < k < 8$ [5] **[10]**		DA1	Upon correct factorisation

Downloaded from SingaKids.com


9 The diagram shows part of the graph of $y = 4 - |x+1|$.

(i) Find the coordinates of the points *A*, *B*, *C* and *D*. [5]

(ii) Find the number of solutions of the equation $4 - |x+1| = mx + 3$ when

(a) $m = 2$ **(b)** $m = -1$ [2]

(iii) State the range of values of *m* for which the equation $4 - |x+1| = mx + 3$ has two solutions. [1]




SGFREEPAPERS.COM

(i)	B(−1,4), D(0,3) $4 - \|x+1\| = 0$ $\|x+1\| = 4$ $x+1 = \pm 4$ $x+1 = 4$ or $x+1 = -4$ $x = 3$ or $x = -5$ *A*(−5,0) *C*(3,0) [5]	A1+A1 B1 A1 +A1
(ii) (a) (b)	$4 - \|x+1\| = mx + 3$ When $m = 2$, the number of solutions is 1 When $m = -1$, the number of solutions is infinite [2]	 A1 A1
(iii)	When $-1 < m < 1$, the number of solutions is 2 [1] **[8]**	A1

Downloaded from SingaKids.com

10 The diagram shows a cone of radius r cm and height h cm. It is given that the volume of the cone is 10π cm^3.



(i) Show that the curved surface area, A cm^2, of the cone, is $A = \dfrac{\pi\sqrt{r^6 + 900}}{r}$. [3]

(ii) Given that r can vary, find the value of r for which A has a stationary value. [4]

(iii) Determine whether this value of A is a maximum or a minimum. [2]

Downloaded from SingaKids.com

10(i)	Volume $= \frac{1}{3}\pi r^2 h = 10\pi$ $h = \frac{30}{r^2}$ $l^2 = r^2 + h^2$ $= r^2 + \left(\frac{30}{r^2}\right)^2$ $l = \sqrt{r^2 + \frac{900}{r^4}}$ $A = \pi r l = \pi r\sqrt{r^2 + \frac{900}{r^4}}$ $A = \pi r\sqrt{\frac{(r^6+900)}{r^4}}$ $A = \frac{\pi r\sqrt{(r^6+900)}}{r^2}$ $A = \frac{\pi\sqrt{(r^6+900)}}{r}$ [3]	B1 M1 A1	If put cm^2 -1m over all
(ii)	$u = \pi\sqrt{r^6+900}$, $v = r$ $\frac{du}{dr} = \frac{1}{2}\times\pi\times(r^6+900)^{-\frac{1}{2}}\times 6r^5$ $\frac{dv}{dr} = 1$ $\frac{du}{dr} = 3\pi r^5(r^6+900)^{-\frac{1}{2}}$ $\frac{dA}{dr} = \frac{3\pi r^6(r^6+900)^{-\frac{1}{2}} - \pi(r^6+900)^{\frac{1}{2}}}{r^2}$ When $\frac{dA}{dr} = 0$ $\frac{\pi(r^6+900)^{-\frac{1}{2}}[3r^6-r^6-900]}{r^2} = 0$ $\frac{\pi[3r^6-r^6-900]}{r^2\,(r^6+900)^{\frac{1}{2}}} = 0$ $2r^6 - 900 = 0$ $r^6 = 450$ $r = 2.77$ [4]	B1 B1 M1 A1	Either $u\frac{dv}{dx}$ *or* $v\frac{du}{dx}$ With the use of quotient rule or product rule Perfect $\frac{dA}{dr} = 0$ with substitution With cm -1m overall

Downloaded from SingaKids.com

(iii)	<table><tr><td>r</td><td>r < 2.768</td><td>r = 2.768</td><td>r > 2.768</td></tr><tr><td>$\frac{dA}{dr}$</td><td>-</td><td>0</td><td>+</td></tr><tr><td>Sketch</td><td>\</td><td>–</td><td>/</td></tr></table> A is a minimum when r = 2.77 [2] **[9]**	M1 DA1	For subst with + r Upon correct $\frac{dA}{dr}$


Downloaded from SingaKids.com

11 The equation of a curve is $y = x(2-x)^3$.

(i) Find the range of values of x for which y is an increasing function. [5]

(ii) Find the coordinates of the stationary points of the curve. [3]

(iii) Hence, sketch the graph of $y = x(2-x)^3$. [3]

$y = x(2-x)^3$ $\frac{dy}{dx} = (2-x)^3(1) - 3x(2-x)^2$ $= (2-x)^2[2-x-3x]$ $= (2-x)^2(2-4x)$ when $\frac{dy}{dx} > 0$, $2-4x > 0$ $-4x > -2$ $x < \frac{1}{2}$ [5] Note: should not see this $2-x > 0$ $-x > -2$ $x < 2$ No [A1]	B1 +B1 A1 M1 A1	Either $v\frac{du}{dx}$ *or* $u\frac{dv}{dx}$ and the use of product rule Perfect for $\frac{dy}{dx} > 0$ with substitution
when $\frac{dy}{dx} = 0$, $(2-x)^2(2-4x) = 0$ $x = 2$, $x = \frac{1}{2}$ $y = 2(2-2)^3$ $y = \frac{1}{2}\left(2-\frac{1}{2}\right)^3$ $= 0$ $= \frac{27}{16}$ Ans $(2,0)$ $\left(\frac{1}{2}, \frac{27}{16}\right)$ [3]	M1 A1+A1	$\frac{dy}{dx} = 0$ with substitution If $(2-x)(2-4x) = 0$ don't penalise]
 [3] **[11]**	B1√ B1√ B1	their max pt $\left(\frac{1}{2}, \frac{27}{16}\right)$ (2,0) their pt of inflexion (0,0) -1m for less than perfect

Downloaded from SingaKids.com

Name: _____ ( ) Class: _____

PRELIMINARY EXAMINATION
GENERAL CERTIFICATE OF EDUCATION ORDINARY LEVEL

ADDITIONAL MATHEMATICS 4047/02

Paper 2

Friday 17 August 2018

2 hours 30 minutes

Additional Materials: Answer Paper
Graph Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class, and index number on all the work you hand in.
Write in dark blue or black pen on both sides of the paper.
You may use a pencil for any diagrams or graphs.
Do not use paper clips, highlighters, glue, or correction fluid.

Answer **all** the questions.
Write your answers on the separate Answer Paper provided
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, staple all your work together with this cover sheet.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is **100**.

FOR EXAMINER'S USE

Q1		Q5		Q9	
Q2		Q6		Q10	
Q3		Q7		Q11	
Q4		Q8		Q12	

100

This document consists of **5** printed pages.

圣尼各拉女校
CHIJ ST. NICHOLAS GIRLS' SCHOOL
Girls of Grace ● Women of Strength ● Leaders with Heart

[Turn over

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + ... + \binom{n}{r}a^{n-r}b^r + ... + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1).....(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$

$$\sin(A \pm B) = \sin A\cos B \pm \cos A\sin B$$

$$\cos(A \pm B) = \cos A\cos B \mp \sin A\sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A\tan B}$$

$$\sin 2A = 2\sin A\cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 **(i)** On the same axes sketch the curves $y^2 = 64x$ and $y = -x^2$. [2]

(ii) Find the equation of the line passing through the points of intersection of the two curves. [4]

2 The roots of the equation $x^2 + 2x + p = 0$, where p is a constant, are α and β.
The roots of the equation $x^2 + qx + 27 = 0$, where q is a constant, are α^3 and β^3.
Find the value of p and of q. [6]


3 **(a)** Given that $3^{2x-2} \times 5^{-2x} = 27^x \div 5^{x+1}$, evaluate the exact value of 15^x. [3]

(b) Given that $\log_x y = 64 \log_y x$, express y in terms of x. [4]

4 **(i)** Write down, and simplify, the first three terms in the expansion of $(1 - \frac{x^2}{2})^n$, in ascending powers of x, where n is a positive integer greater than 2. [2]

(ii) The first three terms in the expansion, in ascending powers of x, of $(2 + 3x^2)(1 - \frac{x^2}{2})^n$ are $2 - px^2 + 2x^4$, where p is an integer. Find the value of n and of p. [5]

5



In the figure, XYZ is a straight line that is tangent to the circle at X.
XQ bisects $\angle RXZ$ and cuts the circle at S. RS produced meets XZ at Y and $ZR = XR$.
Prove that

(a) $SR = SX$, [3]

(b) a circle can be drawn passing through Z, Y, S and Q. [4]

Downloaded from SingaKids.com

6 The expression $3x^3 + ax^2 + bx + 4$, where a and b are constants, has a factor of $x - 2$ and leaves a remainder of -9 when divided by $x + 1$.

(i) Find the value of a and of b. [4]

(ii) Using the values of a and b found in part **(i)**, solve the equation $3x^3 + ax^2 + bx + 4 = 0$, expressing non-integer roots in the form $\frac{c \pm \sqrt{d}}{3}$, where c and d are integers. [4]

7 **(a)** Prove that $\sec\theta + 1 = \frac{\tan\theta \sin\theta}{1 - \cos\theta}$. [4]

(b) Hence or otherwise, solve $\frac{\tan\theta \sin\theta}{1 - \cos\theta} = \frac{3}{4}\sec^2\theta$ for $0 \le \theta \le 2\pi$. [4]

8 The temperature, A °C, of an object decreases with time, t hours. It is known that A and t can be modelled by the equation $A = A_0 e^{-kt}$, where A_0 and k are constants.
Measured values of A and t are given in the table below.

t (hours)	2	4	6	8
A (°C)	49.1	40.2	32.9	26.9

(i) Plot ln A against t for the given data and draw a straight line graph. [2]

(ii) Use your graph to estimate the value of A_0 and of k. [4]

(iii) Assuming that the model is still appropriate, estimate the number of hours for the temperature of the object to be halved. [2]

9 The curve $y = f(x)$ passes through the point $(0, 3)$ and is such that $f'(x) = \left(e^x + \frac{1}{e^x}\right)^2$.

(i) Find the equation of the curve. [4]

(ii) Find the value of x for which $f''(x) = 3$. [4]

Downloaded from SingaKids.com

10 A circle has the equation $x^2 + y^2 + 4x + 6y - 12 = 0$.

(i) Find the coordinates of the centre of the circle and the radius of the circle. [3]

The highest point of the circle is *A*.


(ii) State the equation of the tangent to the circle at *A*. [1]

(iii) Determine whether the point $(0, -7)$ lies within the circle. [2]

The equation of a chord of the circle is $y = 7x - 14$.

(iv) Find the length of the chord. [5]

11



The diagram shows part of the curve of $y = x^2 - 7x + 12$ passing through the point *B* and meeting the *x*-axis at the point *A*.

(i) Find the gradient of the curve at *A*. [4]

The normal to the curve at *A* intersects the curve at *B*.

(ii) Find the coordinates of *B*. [4]

The line *BC* is perpendicular to the *x*-axis.

(iii) Find the area of the shaded region. [4]

12 A particle *P* moves in a straight line, so that, *t* seconds after passing through a fixed point *O*, its velocity, *v* m s^{-1}, is given by $v = \cos t - \sin 2t$, where $0 \le t \le \frac{\pi}{2}$. Find

(i) in terms of π, the values of *t*, when *P* is at instantaneous rest, [5]

(ii) the distance travelled by *P* from $t = 0$ to $t = \frac{\pi}{2}$, [6]


(iii) an expression for the acceleration of *P* in terms of *t*. [1]

Downloaded from SingaKids.com

St Nicholas Girls School Additional Mathematics Preliminary Examination Paper II 2018

Answers

1 (i)



(ii) $y = -4x$

2 $p = 3, q = -10$

3 (a) $\frac{5}{9}$ (b) $y = x^8$, $y = x^{-8}$

4 (i) $1 - n\left(\frac{x^2}{2}\right) + \frac{n(n-1)}{8}x^4 + \cdots$ (ii) $n = 8, \quad p = 5$

6 (i) $a = -8, b = 2$ (ii) $x = 2$, $x = \frac{1 \pm \sqrt{7}}{3}$

7 (b) $\frac{\pi}{3}, \frac{5\pi}{3}$

8 (ii) $A_0 = 59.7$, $k = 0.1$ (iii) 6.93

9 (i) $y = \frac{1}{2}e^{2x} + 2x - \frac{1}{2}e^{-2x} + 3$ (ii) $\frac{1}{2}\ln 2$

10 (i) Centre = $(-2, -3)$, Radius $= 5$ units (ii) $y = 2$

(iii) The distance of the point from the centre of the cicle $= \sqrt{20} < \sqrt{25}$ radius of the circle, so the point lies within the circle.

(iv) $5\sqrt{2}$ units

11 (i) -1 (ii) $B(5, 2)$ (iii) 1sq unit.

12 (i) $\frac{\pi}{2}$, $\frac{\pi}{6}$ (ii) $\frac{1}{2}$m (iii) $-\sin t - 2\cos 2t$

Downloaded from SingaKids.com

Name: _____ ( ) Class: _____

PRELIMINARY EXAMINATION
GENERAL CERTIFICATE OF EDUCATION ORDINARY LEVEL

ADDITIONAL MATHEMATICS 4047/02

Paper 2

Friday 17 August 2018

2 hours 30 minutes

Marking Scheme

Additional Materials: Answer Paper
Graph Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class, and index number on all the work you hand in.
Write in dark blue or black pen on both sides of the paper.
You may use a pencil for any diagrams or graphs.
Do not use paper clips, highlighters, glue, or correction fluid.

Answer **all** the questions.
Write your answers on the separate Answer Paper provided
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, staple all your work together with this cover sheet.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is **100**.

FOR EXAMINER'S USE

Q1		Q5		Q9	
Q2		Q6		Q10	
Q3		Q7		Q11	
Q4		Q8		Q12	

100

This document consists of **5** printed pages.

Downloaded from SingaKids.com

Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + ... + \binom{n}{r}a^{n-r}b^r + ... + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1).....(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$

$$\sin(A \pm B) = \sin A\cos B \pm \cos A\sin B$$

$$\cos(A \pm B) = \cos A\cos B \mp \sin A\sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A\tan B}$$

$$\sin 2A = 2\sin A\cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$


$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 **(i)** On the same axes sketch the curves $y^2 = 64x$ and $y = -x^2$. [2]

(ii) Find the equation of the line passing through the points of intersection of the two curves. [4]

(i)



B1 +B1

-1 mark if no label

[2]

(ii)	$y^2 = 64x$ ------- (1) $y = -x^2$ ------- (2) Sub (2) into (1), $(-x^2)^2 = 64x$ $x^4 = 64x$ $x^4 - 64x = 0$ $x(x^3 - 64) = 0$ $x = 0$ or $x^3 - 64 = 0$ $y = 0$ $x^3 = 64$ $x = 4$ $y = -16$ $m = \frac{-16-0}{4-0}$ $= -4$ $y = -4x$ [4] **[6]**	M1 B1+B1 DA1	Solving Simultaneous Equations Either 1 pairs of x values or y values. [or 1m for each pair of x and y values] Must have (–4,16)

Downloaded from SingaKids.com

2 The roots of the equation $x^2 + 2x + p = 0$, where p is a constant, are α and β.

The roots of the equation $x^2 + qx + 27 = 0$, where q is a constant, are α^3 and β^3.

Find the value of p and of q. [6]

2	$x^2 + 2x + p = 0$ $\quad$ $x^2 + qx + 27 = 0$ $\alpha + \beta = -2$ $\quad$ $\alpha^3 + \beta^3 = -q$	B1	For both sum of roots or first pair of sum & product of roots.
	$\alpha\beta = p$ $\quad$ $\alpha^3\beta^3 = 27$	B1	For both product of roots or 2nd pair of product and sum of roots
	$\alpha\beta = 3$		
	$p = 3$	A1	
	$(\alpha+\beta)(\alpha^2 - \alpha\beta + \beta^2) = -q$ or $(\alpha+\beta)^3 - 3\alpha^2\beta + 3\beta^2\alpha = -q$	B1	For $\alpha^3 + \beta^3$
	$(\alpha+\beta)[(\alpha+\beta)^2 - 2\alpha\beta - \alpha\beta] = -q$ or $(\alpha+\beta)^3 - 3\alpha\beta(\alpha+\beta) = -q$		
	$(-2)[4 - 9] = -q$ or $(-2)^3 - 3p(-2) = -q$	M1√	
	$q = -10$ **[6]**	A1	

SGFREEPAPERS.COM

Downloaded from SingaKids.com

3 **(a)** Given that $3^{2x-2} \times 5^{-2x} = 27^{x} \div 5^{x+1}$, evaluate the exact value of 15^{x}. [3]

(b) Given that $\log_x y = 64 \log_y x$, express y in terms of x. [4]

(a)	$3^{2x-2} \times 5^{-2x} = 27^{x} \div 5^{x+1}$ Method (i) $3^{2x-2} \times 5^{-2x} = 3^{3x} \times 5^{-1-x}$ $\frac{3^{2x-2}}{3^{3x}} = \frac{5^{-1-x}}{5^{-2x}}$ $3^{2x-2-3x} = 5^{-1-x+2x}$ $3^{-x-2} = 5^{x-1}$ $3^{-x} \times 3^{-2} = 5^{x} \times 5^{-1}$ $3^{x} \times 5^{x} = 5^{-1} \div 3^{-2}$	M1 M1√	applying index Law correctly on either LHS or RHS grouping and making power of x on one side
	Method (ii) $3^{2x} \times 3^{-2} \times 5^{-2x} = 3^{3x} \times 5^{-x} \times 5^{-1}$ $3^{x} \times 5^{x} = 5^{-1} \div 3^{-2}$	M1 M1√	Applying index law grouping and making power of x on one side
	$15^{x} = \frac{5}{9}$ [3]	A1	
(b)	$\log_x y = 64 \log_y x$ $\log_x y = \frac{64 log_x x}{log_x y}$ $(log_x y)^2 = 64$ $log_x y = \pm 8$ $y = x^{8}$, $y = x^{-8}$ [4] **[7]**	B1 M1√ A1+A1	change of base

Downloaded from SingaKids.com

4 **(i)** Write down, and simplify, the first three terms in the expansion of $(1-\frac{x^2}{2})^n$, in ascending powers of x, where n is a positive integer greater than 2. [2]


(ii) The first three terms in the expansion, in ascending powers of x, of $(2+3x^2)(1-\frac{x^2}{2})^n$ are $2 - px^2 + 2x^4$, where p is an integer. Find the value of n and of p. [5]

(i)	$\left(1-\frac{x^2}{2}\right)^n = 1 - n\left(\frac{x^2}{2}\right) + {}^nC_2\left(\frac{x^4}{4}\right) + \cdots \ldots\ldots\ldots$ $\left(1-\frac{x^2}{2}\right)^n = 1 - n\left(\frac{x^2}{2}\right) + \frac{n(n-1)}{8}x^4 + \cdots \ldots\ldots\ldots$	M1 B1	Or any two terms 1m, perfect 2m [2]
(ii)	$(2+3x^2)(1-\frac{x^2}{2})^n = (2+3x^2)(1-\frac{nx^2}{2}+\frac{n(n-1)}{8}x^4+\cdots.)$ $= 2 - nx^2 + \frac{n(n-1)}{4}x^4 + 3x^2 - \frac{3n}{2}x^4 + \cdots \ldots\ldots..$ $= 2 - (n-3)x^2 + \left(\frac{n^2-7n}{4}\right)x^4 + \cdots ....$ $= 2 - px^2 + 2x^4 + \cdots ....$ $\frac{n^2-7n}{4} = 2$ $n^2 - 7n - 8 = 0$ $(n-8)(n+1) = 0$ $n = 8, n = -1$(NA) $-n + 3 = -p$ $-8 + 3 = -p$ $p = 5$	M1√ M1√ DA1 M1√ A1	factorisation Upon correct factorisation [5] **[7]**

SGFREEPAPERS.COM

Downloaded from SingaKids.com

5



In the figure, *XYZ* is a straight line that is tangent to the circle at *X*.

XQ bisects $\angle RXZ$ and cuts the circle at *S*. *RS* produced meets *XZ* at Y and *ZR* = *XR*.

Prove that

(a) *SR* = *SX*, [3]

(b) a circle can be drawn passing through *Z*, *Y*, *S* and *Q*. [4]

(a)	$\angle ZXQ = \angle SRX$ (Alternate Segment Theorem) $\angle ZXQ = \angle QXR$ (XQ is the angle bisector of $\angle RXZ$) $\angle QXR = \angle SRX$ By base angles of isosceles triangles, SR=SX [3]	B1 B1 B1
(b)	Let $\angle QXR$ be *x* $\angle RSX = 180^\circ - 2x$ (Isosceles Triangle) $\angle YSQ = 180^\circ - 2x$ (Vertically Opposite Angles) $\angle RZX = \angle ZXR = 2x$ (Base angles of Isosceles Triangle) $\angle RZX + \angle YSQ = 180^\circ - 2x + 2x = 180^\circ$ Since opposite angles are supplementary in cyclic quadrilaterals, a circle that passes through Z, Y, S and Q can be drawn **Alternative** [4] Similar but use of tangent secant theorem. **[7]**	B1 B1 B1 B1

Downloaded from SingaKids.com

6 The expression $3x^3 + ax^2 + bx + 4$, where a and b are constants, has a factor of $x - 2$ and leaves a remainder of -9 when divided by $x + 1$.

(i) Find the value of a and of b. [4]

(ii) Using the values of a and b found in part **(i)**, solve the equation $3x^3 + ax^2 + bx + 4 = 0$, expressing non-integer roots in the form $\frac{c \pm \sqrt{d}}{3}$, where c and d are integers. [4]

(i)	$f(x) = 3x^3 + ax^2 + bx + 4$ x-2 is a factor $f(2) = 0$ $3(8) + 4a + 2b + 4 = 0$ $4a + 2b + 28 = 0$ $2a + b + 14 = 0$---------(1) $f(-1) = -9$ $-3 + a - b + 4 = -9$ $a - b = -10$---------(2) (1)+(2) $3a = -24$ $a = -8$ Sub into (2) $-8 - b = -10$ $b = 2$ [4]	M1 M1 A1 A1
(ii)	$x - 2 \overline{)\, 3x^3 - 8x^2 + 2x + 4}$ $3x^3 - 6x^2$ $-2x^2 + 2x$ $-2x^2 + 4x$ $-2x + 4$ $-2x + 4$ $3x^3 - 8x^2 + 2x + 4 = 0$ $(x - 2)(3x^2 - 2x - 2) = 0$ $x - 2 = 0$ $3x^2 - 2x - 2 = 0$ $x = \frac{2 \pm \sqrt{(-2)^2 - 4 \times 3 \times -2}}{2 \times 3}$ $x = \frac{2 \pm \sqrt{28}}{6}$ $x = \frac{2(1 \pm \sqrt{7})}{6}$ $x = 2$ $x = \frac{1 \pm \sqrt{7}}{3}$ [4] **[8]**	B1 M1√ A1 +A1

SGFREEPAPERS.COM

Downloaded from SingaKids.com

7 **(a)** Prove that $\sec\theta + 1 = \dfrac{\tan\theta\sin\theta}{1-\cos\theta}$. [4]

(b) Hence or otherwise, solve $\dfrac{\tan\theta\sin\theta}{1-\cos\theta} = \dfrac{3}{4}\sec^2\theta$ for $0 \le \theta \le 2\pi$. [4]

(a)	$RHS = \dfrac{\tan\theta\sin\theta}{1-\cos\theta}$ $= \dfrac{\frac{\sin\theta}{\cos\theta}\sin\theta}{1-\cos\theta}$ $= \dfrac{\frac{\sin^2\theta}{\cos\theta}}{1-\cos\theta}$ $= \dfrac{1-\cos^2\theta}{(1-\cos\theta)\cos\theta}$ $= \dfrac{(1-\cos\theta)(1+\cos\theta)}{(1-\cos\theta)\cos\theta}$ $= \dfrac{1+\cos\theta}{\cos\theta}$ $= \dfrac{1}{\cos\theta} + 1$ $= \sec\theta + 1$ [4]	B1 B1 B1 B1	change *tan* change sin² to cos² identity $a^2 - b^2$ split and bring to answer
(b)	$\dfrac{\tan\theta\sin\theta}{1-\cos\theta} = \dfrac{3}{4}\sec^2\theta$ $1 + \sec\theta = \dfrac{3}{4}\sec^2\theta$ $3\sec^2\theta - 4\sec\theta - 4 = 0$ $(\sec\theta - 2)(3\sec\theta + 2) = 0$ $\sec\theta = 2$ or $\sec\theta = -\dfrac{2}{3}$ $\cos\theta = \dfrac{1}{2}$ or $\theta = \dfrac{\pi}{3}, \dfrac{5\pi}{3}$ $\cos\theta = -\dfrac{3}{2}$ (No Solution) = 1.05, 5.24 [4] **[8]**	B1 M1 DA1+ DA1	substitution factorization 1st DA1 for change to cos & no soln Upon correct factorisation

Downloaded from SingaKids.com

8 The temperature, A °C, of an object decreases with time, t hours. It is known that A and t can be modelled by the equation $A = A_0 e^{-kt}$, where A_0 and k are constants.

Measured values of A and t are given in the table below.

t (hours)	2	4	6	8
A (°C)	49.1	40.2	32.9	26.9

(i) Plot ln A against t for the given data and draw a straight line graph. [2]

(ii) Use your graph to estimate the value of A_0 and of k. [4]

(iii) Assuming that the model is still appropriate, estimate the number of hours for the temperature of the object to be halved. [2]

8 **(i)** B1 for correct points, values & correct axes.


B1 best fit line . [2]

t	2	4	6	8
ln A	3.89	3.69	3.49	3.29

SGFREEPAPERS.COM

(ii)	$A = A_0 e^{-kt}$ $\ln A = -kt + \ln A_0$ $-k$ = gradient $-k = \dfrac{3.39 - 3.99}{7 - 1}$ $k = 0.1 \pm 0.02$ $\ln A_0 = 4.09$ $A_0 = e^{4.09}$ $A_0 = 59.7$ (3s.f.) ±4 [4]	M1 A1 M1 A1	gradient vertical intercept
(iii)	$\frac{1}{2}A_0 = 29.865$ Or $\frac{1}{2}A_0 = A_0 e^{-kt}$ $\ln 29.865 = 3.396$ OR $\frac{1}{2} = e^{-0.1t}$ From the graph, t = 6.9 $t = 6.93$ (3s.f.) [2] **[8]**	√M1 A1 ±0.5	

Downloaded from SingaKids.com



[Turn over

Downloaded from SingaKids.com

9 The curve $y = f(x)$ passes through the point $(0,3)$ and is such that $f'(x) = \left(e^x + \frac{1}{e^x}\right)^2$.

(i) Find the equation of the curve. [4]

(ii) Find the value of x for which $f''(x) = 3$. [4]

(i)	$y = \int \left(e^x + \frac{1}{e^x}\right)^2 \mathrm{d}x$ $= \int e^{2x} + 2 + e^{-2x} \mathrm{d}x$ $= \frac{e^{2x}}{2} + 2x + \frac{e^{-2x}}{-2} + c$ at $(0,3)$, $3 = \frac{1}{2}e^0 + 2(0) - \frac{1}{2}e^0 + c$ $c = 3$ $y = \frac{1}{2}e^{2x} + 2x - \frac{1}{2}e^{-2x} + 3$ [4]	M1 B1 M1 A1	knowing $y = \int f(x)\mathrm{d}x$ ignore no + c for ubstitution
(ii)	$f'(x) = e^{2x} + 2 + e^{-2x}$ $\quad f'(x) = \left(e^x + e^{-x}\right)^2$ $f''(x) = 2e^{2x} - 2e^{-2x}$ $\quad f''(x) = 2\left(e^x + e^{-x}\right)\left(e^x - e^{-x}\right)$ when $f''(x) = 3$, $2e^{2x} - 2e^{-2x} = 3$ Let $e^{2x} = a$, $\quad 2a - \frac{2}{a} = 3$ $2a^2 - 2 = 3a$ $2a^2 - 3a - 2 = 0$ $(2a+1)(a-2) = 0$ $a = -\frac{1}{2}$ $\quad a = 2$ $e^{2x} = -\frac{1}{2}$ $\quad e^{2x} = 2$ no solution $\quad 2x = \ln 2$ $x = \frac{1}{2}\ln 2 = \ln\sqrt{2} = 0.347$ [4] **[8]**	B1 M1 +DA1 +DA1	 factorisation Upon correct factorisation

SGFREEPAPERS.COM

Downloaded from SingaKids.com

10 A circle has the equation $x^2 + y^2 + 4x + 6y - 12 = 0$.

(i) Find the coordinates of the centre of the circle and the radius of the circle. [3]

The highest point of the circle is *A*.

(ii) State the equation of the tangent to the circle at *A*. [1]

(iii) Determine whether the point $(0, -7)$ lies within the circle. [2]

The equation of a chord of the circle is $y = 7x - 14$.

(iv) Find the length of the chord. [5]


Downloaded from SingaKids.com

(i)	$x^2 + y^2 + 4x + 6y - 12 = 0$ $x^2 + y^2 + 2gx + 2fy + c = 0$ $2g = 4 \qquad 2f = 6$ $g = 2 \qquad f = 3$ Centre = $(-2, -3)$ Radius $= \sqrt{g^2 + f^2 - C}$ $= \sqrt{(-2)^2 + (-3)^2 - (-12)}$	$(x)^2 + 2(x)(2) + (2)^2 + (y)^2 + 2(y)(3) + (3)^2$ $= 12 + (2)^2 + (3)^2$ $(x+2)^2 + (y+3)^2 = 25$	A1 M1
	Radius $= 5$ units [3]		A1 ignore no unit
(ii)	$y = 2$ (y= their y coord of centre +radius) [1]		B1 √
(iii)	The distance of the point from the centre of the cicle $= \sqrt{(0-(-2))^2 + (-7-(-3))^2}$ $= \sqrt{20} \quad < \sqrt{25}$ Since it is lesser than the radius of the circle, it lies within the circle. [2]		M1√ their centre DA1
(iv)	$y = 7x - 14$ ------- (1) $x^2 + y^2 + 4x + 6y - 12 = 0$ ------- (2) Sub (1) into (2), $x^2 + (7x-14)^2 + 4x + 6(7x-14) - 12 = 0$ $x^2 + 49x^2 - 196x + 196 + 4x + 42x - 84 - 12 = 0$ $50x^2 - 150x + 100 = 0$ $x^2 - 3x + 2 = 0$ $(x-1)(x-2) = 0$ $x = 1$ or $x = 2$ Sub into (1), $y = -7$ or $y = 0$ The length of the chord $= \sqrt{(1-2)^2 + (-7-0)^2}$ $= \sqrt{50}$ $= 5\sqrt{2}$ units [5] **[11]**		M1 Solving simultaneous equations M1 Factorizing B1 Either 1 pair correct or both x solutions are correct √M1 A1 accept 7.07

SGFREEPAPERS.COM

Downloaded from SingaKids.com

11



The diagram shows part of the curve of $y = x^2 - 7x + 12$ passing through the point B and meeting the x-axis at the point A.

(i) Find the gradient of the curve at A. [4]

The normal to the curve at A intersects the curve at B.

(ii) Find the coordinates of B. [4]

The line BC is perpendicular to the x-axis.

(iii) Find the area of the shaded region. [4]

Downloaded from SingaKids.com

(i)	$y = x^2 - 7x + 12$ $= (x-3)(x-4)$ $\frac{dy}{dx} = 2x - 7$ *when* $x = 3$, $\frac{dy}{dx} = 2(3) - 7$ $= -1$ [4]	M1 B1 M1 A1	using smaller (positive) x value
(ii)	$\perp m = 1$ sub $m = 1$ and $(3,0)$ into $y = mx + c$ $0 = 1(3) + c$ $c = -3$ equation of normal: $y = x - 3$ $x^2 - 7x + 12 = x - 3$ or $(x-3)(x-4) = x - 3$ $x^2 - 8x + 15 = 0$ $x - 4 = 1$ $(x-3)(x-5) = 0$ $x = 5$ $x = 3$ $x = 5$ $y = 2$ $B(5,2)$ [4]	M1 M1 M1 A1	sub $\perp$m and their(3,0) curve and normal factorisation
(iii)	Area $= \left\lvert \int_3^4 x^2 - 7x + 12\,dx \right\rvert + \int_4^5 x^2 - 7x + 12\,dx$ $= \left\lvert \left[\frac{x^3}{3} - \frac{7x^2}{2} + 12x \right]_3^4 \right\rvert + \left[\frac{x^3}{3} - \frac{7x^2}{2} + 12x \right]_4^5$ $= \left\lvert \left(\frac{64}{3} - \frac{7(16)}{2} + 12(4) \right) - \left(\frac{27}{3} - \frac{7(9)}{2} + 12(3) \right) \right\rvert$ $+ \left(\frac{125}{3} - \frac{7(25)}{2} + 12(5) \right) - \left(\frac{64}{3} - \frac{7(16)}{2} + 12(4) \right)$ $= \left\lvert 13\frac{1}{3} - 13\frac{1}{2} \right\rvert + 14\frac{1}{6} - 13\frac{1}{3}$ $= \left\lvert -\frac{1}{6} \right\rvert + \frac{5}{6}$ $= 1$ sq unit [4] **[12]**	M1 B1 M1 A1	Area $= \left\lvert \int y\,dx \right\rvert + \int y\,dx$ √their limits from (i) and (ii) for integration substitution

SGFREEPAPERS.COM

Downloaded from SingaKids.com

12 A particle P moves in a straight line, so that, t seconds after passing through a fixed point O, its velocity, v m s^{-1}, is given by $v = \cos t - \sin 2t$, where $0 \le t \le \frac{\pi}{2}$. Find

(i) in terms of π, the values of t, when P is at instantaneous rest, [5]

(ii) the distance travelled by P from $t = 0$ to $t = \frac{\pi}{2}$, [6]

(iii) an expression for the acceleration of P in terms of t. [1]

Downloaded from SingaKids.com

(i)	$v = \cos t - \sin 2t$ when $v = 0$, $\cos t - \sin 2t = 0$ $\cos t - 2\sin t \cos t = 0$ $\cos t(1 - 2\sin t) = 0$ $\cos t = 0$ $\sin t = \frac{1}{2}$ $t = \frac{\pi}{2}$ $t = \frac{\pi}{6}$ [5]	B1 B1 M1 A1+A1	For v=0 for double angle factorisation
(ii)	$s = \int \cos t - \sin 2t \, dt$ $= \sin t + \frac{1}{2}\cos 2t + c$ when $t = 0, s = 0$ $0 = \sin 0 + \frac{1}{2}\cos 0 + c$ $c = -\frac{1}{2}$ $s = \sin t + \frac{1}{2}\cos 2t - \frac{1}{2}$ when $t = \frac{\pi}{6}$, $s = \sin\frac{\pi}{6} + \frac{1}{2}\cos\frac{\pi}{3} - \frac{1}{2}$ $= \frac{1}{2} + \frac{1}{2}\left(\frac{1}{2}\right) - \frac{1}{2}$ $= \frac{1}{4}$ when $t = \frac{\pi}{2}$, $s = \sin\frac{\pi}{2} + \frac{1}{2}\cos\pi - \frac{1}{2}$ $= 1 + \frac{1}{2}(-1) - \frac{1}{2}$ $= 0$ Distance travelled = $2\left(\frac{1}{4}\right)$ $= \frac{1}{2}$ m [6]	B1 B1+B1 M1 M1 DA1	For $s = \int v \, dt$ Integration ignore no +c Sub either $t = \frac{\pi}{6}$ or $t = \frac{\pi}{2}$ For both s for $t = \frac{\pi}{6}$ and $t = \frac{\pi}{2}$ found
(iii)	$a = \frac{dv}{dt} = (-\sin t - 2\cos 2t) m/s^2$ [1] **[12]**	B1	

Downloaded from SingaKids.com

Downloaded from SingaKids.com