COMMONWEALTH SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2018

ADDITIONAL MATHEMATICS
PAPER 1

Name: _____ ( ) Class: _____

SECONDARY FOUR EXPRESS **Wednesday 12 September 2018**
SECONDARY FIVE NORMAL ACADEMIC **11 00 – 13 00**
4047/1 **2 h**

READ THESE INSTRUCTIONS FIRST

Write your name, index number and class on all the work you hand in.
Write in dark blue or black pen on both sides of the paper.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Write your answers on the separate writing paper provided.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

Name of setter: Mrs Margaret Loh

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots\ldots + \binom{n}{r}a^{n-r}b^r + \ldots\ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\ldots\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$

Downloaded from SingaKids.com

1. The slope at any point (x, y) of a curve is given by $\frac{dy}{dx} = \frac{k}{(2x+3)^2} - 1$ where k is a constant. If the tangent to the curve at $(-1, 0)$ is perpendicular to the line $3y = x + 1$, find

   (i) the value of k, [3]

   (ii) the equation of the curve. [3]

2. (i) On the same axes, sketch the curves $y = -8x^{-\frac{1}{2}}$ and $y^2 = \frac{1}{4}x$. [2]

   (ii) Find the equation of the line passing through the origin and the point of intersection of the two curves. [3]

3. The equation $y = \frac{x+c}{x+d}$, where c and d are constants, can be represented by a straight line when $xy - x$ is plotted against y. The line passes through the points $(0, 4)$ and $(0.2, 0)$.

   (i) Find the value of c and of d, [4]

   (ii) If $(2.5, a)$ is a point on the straight line, find the value of a. [1]

4. The roots of a quadratic equation are α and β, where $\alpha^3 + \beta^3 = 0$, $\alpha\beta = \frac{27}{64}$, $\alpha + \beta > 0$.

   (i) Find this quadratic equation with integral coefficient. [4]

   The roots of another quadratic equation $x^2 + px + q = 0$ are $\alpha - \beta$ and $\beta - \alpha$.


   (ii) Find the value of p and of q. [3]

5. (i) Prove the identity $\sin^2 2x(\cot^2 x - \tan^2 x) = 4\cos 2x$. [4]

   (ii) Hence find, for $0 \le x \le 2\pi$, the values of x for which $\sin^2 2x = \frac{e}{\cot^2 x - \tan^2 x}$. [3]

Downloaded from SingaKids.com

6.



(a) The diagram shows a cylinder of height h cm and base radius r cm inscribed in a cone of height 28 cm and base radius 10 cm. Show that

(i) the height, h cm, of the cylinder is given by

$$h = 28 - \frac{14}{5}r .$$ [1]

(ii) the volume, V cm^3, of the cylinder is given by

$$V = 14\pi r^2(2 - \frac{r}{5}).$$ [1]

(b) (i) Given that r can vary, find the maximum volume of the cylinder. [5]

(ii) Show that, in this case, the cylinder occupies $\frac{4}{9}$ of the volume of the cone. [2]

Downloaded from SingaKids.com

7. (a) A circle with centre P lies in the first quadrant of the Cartesian plane. It is tangential to the x-axis and the y-axis, and passes through the points $A(4, 18)$ and $B(18, 16)$.

Find

(i) the equation of the perpendicular bisector of the line segment AB, [3]

(ii) the coordinates of the centre P, [2]

(iii) the equation of the circle, [1]

The tangent at A touches the x-axis at R. The line joining A and P is produced to touch the x-axis at S.

(b) Find the area of triangle ARS. [4]

8. Use the result $(\sqrt{x}+\sqrt{y})^2 \equiv x+y+2\sqrt{(xy)}$, or otherwise, find the square root of $12+\sqrt{140}$ in the form $\sqrt{a}+\sqrt{b}$, where a and b are constants to be determined. [5]

9. Given that $P(x) = 2x^4 - 5x^3 + 5x^2 - x - 10$,

(i) find the quotient when $P(x)$ is divided by $(2x-1)(x^2+3)$, [2]

(ii) hence express $\dfrac{P(x)}{(2x-1)(x^2+3)}$ in partial fractions. [5]

10. The velocity, v ms^{-1}, of a particle travelling in a straight line at time t seconds after leaving a fixed point O, is given by

$$v = 2t^2 + (1-3k)t + 8k - 1,$$

where k is a constant. The velocity is a minimum at $t = 5$.

(i) Show that $k = 7$. [2]


(ii) Show that the particle will never return to O with time. [2]

(iii) Find the duration when its velocity is less than 13 ms^{-1}. [2]

(iv) Find the distance travelled by the particle during the third second. [2]

Downloaded from SingaKids.com

11.




The diagram shows part of curve $y = \frac{3}{1-2x}$ intersecting with a straight line $y = 2x + 3$ at the point A. Find

(i) the coordinates of A. [2]

(ii) the area of the shaded region bounded by the line and the curve. [4]

Downloaded from SingaKids.com

12.



In the diagram, two circles touch each other at A. TA is tangent to both circles at A and FE is a tangent to the smaller circle at C. Chords AE and AF intersect the smaller circle at B and D respectively. Prove that

(i) line BD is parallel to line FE, [2]

(ii) $\angle FAC = \angle CAE$, [3]

END OF PAPER


Downloaded from SingaKids.com

2018 CWSS Prelim AM P1 Answer Key

1.	(i) -2 (ii) $y=\frac{1}{(2x+3)}-x-2$	10.	(iii) 4s (iv) $17\frac{2}{3}$ m or 17.7 m
2.	(ii) $y=-\frac{1}{8}x$	11.	(i) $A=(-1,1)$ (ii) 0.352 units^2
3.	(i) $c=4$; $d=20$ (ii) -46	12.	(i)To prove: BD // FE
			Proof: Let $\angle TAF$ be θ.
4.	(i) $64x^2-72x+27=0$		$\angle ABD=\angle TAF=\theta$ (alt seg thm)
	(ii) $p=0;\ q=\frac{27}{64}$		$\angle AEF=\angle TAF=\theta$ (alt seg thm)
			$\therefore\ \angle ABD=\angle AEF=\theta$
5.	(ii) 0.412, 2.73, 3.55, 5.87		Using property of corresponding angles, BD // EF (shown)
6.	b(i) $\frac{11200\pi}{27}cm^3$ or $1300\,cm^3$		(ii)To prove: $\angle FAC=\angle CAE$
			Proof: Let $\angle BCE=\alpha$
7	a(i) $y=7x-60$ (ii) (10, 10)		$\angle CBD=\angle BCE=\alpha$ (alt $\angle$s, $BD//EF$)
	(iii) $(x-10)^2+(y-10)^2=100$		$\angle FAC=\angle CBD=\alpha$ ($\angle$s in same segment)
	b. 337.5 units^2		Also, $\angle CAE=\angle BCE=\alpha$ (alt seg thm)
			$\therefore\ \angle FAC=\angle CAE=\alpha$ (shown)
8.	$\sqrt{7}+\sqrt{5}$		
9.	(i) $x-2$ (ii) $x-2-\frac{3}{(2x-1)}+\frac{7}{(x^2+3)}$		

END

Downloaded from SingaKids.com


Qn No	Solutions	Marks
1	$3y = x+1$	
	$y = \frac{1}{3}x + \frac{1}{3}$	
	$\therefore$ grad of tangent $= -3$	M1
(i)	$-3 = \frac{k}{(2x+3)^2} - 1$	M1
	$k = -2$	A1
(ii)	$\frac{dy}{dx} = \frac{-2}{(2x+1)^2} - 1$	
	$y = \int [-2(2x+1)^{-2} - 1]dx$	M1
	$= \frac{-2(2x+1)^{-1}}{(-1)(2)} - x + c$	
	$= \frac{1}{(2x+3)} - x + c$	M1
	When $y = 0,\ x = -1$ $\quad 0 = \frac{1}{-2+3} + 1 + c$	
	$c = -2$	
	$\therefore\ y = \frac{1}{(2x+3)} - x - 2$	A1
2(i)		Graphs are [B1] & [B1]
(ii)	$(-8x^{-\frac{1}{2}})^2 = \frac{1}{4}x$	M1
	$64x^{-1} = \frac{1}{4}x$	
	$256 = x^2$	
	$x = 16$ or -16 (NA)	M1
	When $x = 16,\ y = \frac{-8}{\sqrt{16}} = -2$	

Downloaded from SingaKids.com

	Grad of line $= \frac{-2}{16} = -\frac{1}{8}$	
	$\therefore$ Eqn of line is $y = -\frac{1}{8}x$	A1
3(i)	$y(x+d) = x+c$	
	$xy - x = -yd + c$	M1
	$\therefore \quad c = 4$	B1
	Grad $= -\frac{4}{0.2} = -20$	M1
	$\therefore \quad -d = -20$	
	$d = 20$	A1
(ii)	$\therefore \quad xy - x = -20y + 4$	
	$a = -20(2.5) + 4 \quad = -46$	B1
4	$\alpha^3 + \beta^3 = 0$	
	$(\alpha+\beta)[(\alpha+\beta)^2 - 3\alpha\beta] = 0$	
	$(\alpha+\beta)[(\alpha+\beta)^2 - 3\left(\frac{27}{64}\right)] = 0$	M1
	Since $\alpha \neq -\beta, \quad (\alpha+\beta)^2 = \frac{81}{64}$	
	$\alpha + \beta = \frac{9}{8}$ or $-\frac{9}{8}$ (NA)	A1
(i)	Quad eqn is $x^2 - \frac{9}{8}x + \frac{27}{64} = 0$	M1
	$64x^2 - 72x + 27 = 0$	B1
(ii)	Sum of roots $= \alpha - \beta + \beta - \alpha \quad = 0$	
	Prod of roots $= (\alpha - \beta)(\beta - \alpha)$	
	$= \alpha\beta - \alpha^2 - \beta^2 + \alpha\beta$	
	$= 2\alpha\beta - (\alpha^2 + \beta^2)$	
	$= 2\alpha\beta - [(\alpha+\beta)^2 - 2\alpha\beta]$	M1
	$= 4\alpha\beta - (\alpha+\beta)^2$	
	$= 4\left(\frac{27}{64}\right) - \left(\frac{9}{8}\right)^2$	
	$= \frac{108}{64} - \frac{81}{64} \quad = \frac{27}{64}$	
	$\therefore \quad p = 0 \quad \& \quad q = \frac{27}{64}$	B1, B1

SGFREEPAPERS.COM

Downloaded from SingaKids.com

5(i)	To prove: $\sin^2 2x(\cot^2 x - \tan^2 x) = 4\cos 2x$	
	Proof: LHS = $\sin^2 2x(\cot^2 x - \tan^2 x)$	
	$= \sin^2 2x\left(\frac{\cos^2 x}{\sin^2 x} - \frac{\sin^2 x}{\cos^2 x}\right)$	M1
	$= \sin^2 2x\left(\frac{\cos^4 x - \sin^4 x}{\sin^2 x \cos^2 x}\right)$	M1
	$= 4\sin^2 x\cos^2 x\left(\frac{(\cos^2 x - \sin^2 x)(\cos^2 x + \sin^2 x)}{\sin^2 x \cos^2 x}\right)$	M1
	$= 4(\cos^2 x - \sin^2 x)$	M1
	$= 4\cos 2x$	
	= RHS (proved)	
(ii)	$\sin^2 2x(\cot^2 x - \tan^2 x) = e$	
	$4\cos 2x = e$	M1
	$\cos 2x = \frac{e}{4}$	
	$2x \approx 0.8236,\ 5.4596,\ 7.1068,\ 11.743$	
	$x \approx 0.412,\ 2.73,\ 3.55,\ 5.87$	A1, A1
6a(i)	 Using Similar triangles, $\frac{28-h}{28} = \frac{r}{10}$ $28 - h = \frac{28}{10}r$ $h = 28 - \frac{14}{5}r$ (shown)	M1
(ii)	Vol of cylinder = $\pi r^2 h$	
	$V = \pi r^2\left(28 - \frac{14}{5}r\right)$	M1
	$V = 14\pi r^2\left(2 - \frac{1}{5}r\right)$ (shown)	
b(i)	$\frac{dV}{dr} = 56\pi r - \frac{14}{5}\pi(3r^2)$	
	$= 14\pi r(4 - \frac{3}{5}r)$	M1
	At stat pt, $\frac{dV}{dr} = 0$	
	$14\pi r(4 - \frac{3}{5}r) = 0$	M1

Downloaded from SingaKids.com


	$r = 0$ (NA), $\quad 4 - \frac{3}{5}r = 0 \quad \Rightarrow \quad r = 6\frac{2}{3}$	A1
	$\frac{d^2V}{dr^2} = 56\pi - \frac{84}{5}\pi r$	
	$= 56\pi - \frac{84}{5}\pi\left(6\frac{2}{3}\right)$	
	$= -175.93 (2dp) \quad < 0$	M1
	Since $\frac{d^2V}{dr^2} < 0$, $\therefore$ r = $6\frac{2}{3}$ will make V a maximum.	
	Max volume = $14\pi\left(\frac{20}{3}\right)\left(\frac{20}{3}\right)\left(2 - \frac{1}{5}\left[\frac{20}{3}\right]\right)$	
	$= \frac{11200}{27}\pi$ cm^3 or 1300 cm^3 $(3sf)$	B1
(ii)	<u>To show</u>: Vol of cylinder = $\frac{4}{9}$ (Vol of cone)	
	<u>Proof</u>: Vol of cone = $\frac{1}{3}\pi(10)^2(28) = \frac{2800}{3}\pi$ cm^3	M1
	$\frac{\text{Vol of cylinder}}{\text{Vol of cone}} = \frac{11200\pi}{27} \times \frac{3}{2800\pi} = \frac{4}{9}$	M1
	$\therefore$ Vol of cylinder = $\frac{4}{9}$ (Vol of cone) (shown)	
7a(i)	Mid-pt of AB = $\left(\frac{4+18}{2}, \frac{18+16}{2}\right)$ = (11, 17)	M1
	Grad of AB = $\frac{18-16}{4-18} = -\frac{1}{7}$	
	Grad of perpendicular bisector = 7	M1
	Eqn of perpendicular bisector is $y - 17 = 7(x - 11)$	
	$y = 7x - 60$	A1
(ii)	Let the centre P be (m, m).	
	$m = 7m - 60$	M1
	$m = 10$	
	$\therefore \quad P = (10, 10)$	A1
(iii)	Eqn of circle is $(x-10)^2 + (y-10)^2 = 100$	B1
	Or $x^2 + y^2 - 20x - 20y + 100 = 0$	
(b)	Grad of AP = $\frac{18-10}{4-10}$	
	$= -\frac{4}{3}$	

SGFREEPAPERS.COM

Downloaded from SingaKids.com

	$\therefore$ Grad of tangent at $A = \frac{3}{4}$	
	Eqn of tangent at A is $y - 18 = \frac{3}{4}(x-4)$	
	$y = \frac{3}{4}x + 15$	
	$\therefore R = (-20, 0)$	B1
	Eqn of AP is $y - 10 = -\frac{4}{3}(x-10)$	
	$y = -\frac{4}{3}x + 23\frac{1}{3}$	
	$\therefore S = \left(17\frac{1}{2}, 0\right)$	B1
	$\therefore$ Area of $\Delta ARS = \frac{1}{2}\left(20 + 17\frac{1}{2}\right)(18)$	M1
	$= 337.5$ units^2	A1
8	$x + y = 12$ ----------(1)	B1
	$4xy = 140$ ----------(2)	B1
	From eqn (1): $y = 12 - x$ substi into eqn (2)	
	$4x(12-x) = 140$	M1
	$x^2 - 12x + 35 = 0$	
	$(x-7)(x-5) = 0$	
	$\therefore \quad x = 7$ or $x = 5$	
	When $x = 7, \quad y = 5$	A1
	When $x = 5, \quad y = 7$	
	$\therefore \quad \sqrt{12 + \sqrt{140}} = \left(\sqrt{7} + \sqrt{5}\right)$	A1
9(i)	$(2x-1)(x^2+3) = 2x^3 - x^2 + 6x - 3$	
	$2x^3 - x^2 + 6x - 3 \overline{)\, 2x^4 - 5x^3 + 5x^2 - x - 10}$ $x - 2$ $-(2x^4 - x^3 + 6x^2 - 3x)$ $-4x^3 - x^2 + 2x - 10$ $-(-4x^3 + 2x^2 - 12x + 6)$ $-3x^2 + 14x - 16$	M1
	$\therefore \quad$ Quotient $= x - 2$	A1
(ii)	$\frac{P(x)}{(2x-1)(x^2+3)} = x - 2 + \frac{(-3x^2 + 14x - 16)}{(2x-1)(x^2+3)}$	

Downloaded from SingaKids.com

	$\frac{(-3x^2+14x-16)}{(2x-1)(x^2+3)} = \frac{A}{(2x-1)} + \frac{(Bx+C)}{(x^2+3)}$ where A, B and C are constants	
	$-3x^2+14x-16 = A(x^2+3)+(Bx+C)(2x-1)$	M1
	When $x = \frac{1}{2}$, $-3\left(\frac{1}{4}\right)+14\left(\frac{1}{2}\right)-16 = A\left(3\frac{1}{4}\right)$	
	$A = -3$	B1
	When $x = 0$, $-16 = 3A - C$	
	$-16 = -9 - C$	
	$C = 7$	B1
	Comparing coeff of x^2 : $-3 = A + 2B$	
	$-3 = -3 + 2B$	
	$B = 0$	B1
	$\therefore \quad \frac{P(x)}{(2x-1)(x^2+3)} = x - 2 - \frac{3}{(2x-1)} + \frac{7}{(x^2+3)}$	A1
10(i)	$\frac{dv}{dt} = 4t + (1-3k)$	
	When vel is a minimum, $\frac{dv}{dt} = 0$	
	$4(5) + (1-3k) = 0$	M1
	$3k = 21$	
	$k = 7$ (shown)	A1
(ii)	When $k = 7$, $v = 2t^2 - 20t + 55$	
	Discriminant = $(-20)^2 - 4(2)(55)$	
	$= 400 - 440$	
	$= -40$	
	< 0	M1
	$\Rightarrow$ there is no real values of t such that vel = 0, also vel > 0 hence particle will never return to O with time.	A1
(iii)	$2t^2 - 20t + 55 < 13$	M1
	$2t^2 - 20t + 42 < 0$	
	$t^2 - 10t + 21 < 0$	
	$(t-7)(t-3) < 0$	
	 $\therefore \quad 3 < t < 7$ Duration = $7 - 3 = 4$ s	A1

SGFREEPAPERS.COM

Downloaded from SingaKids.com

(iv)	$s = \int_2^3 (2t^2 - 20t + 55)dt$	M1
	$= \left[\frac{2t^3}{3} - 10t^2 + 55t \right]_2^3$	
	$= [18 - 90 + 165] - \left[\frac{16}{3} - 40 + 110 \right]$	
	$= 17\frac{2}{3}$ m or 17.7 m(3sf)	A1
11(i)	$\frac{3}{1-2x} = 2x + 3$	M1
	$3 = (2x + 3)(1 - 2x)$	
	$3 = 2x - 4x^2 + 3 - 6x$	
	$4x^2 + 4x = 0$	
	$4x(x + 1) = 0$	
	$x = 0$ or $x = -1$	
	For pt A: When $x = -1$, $y = -2 + 3 = 1$	
	$\therefore \quad A = (-1, 1)$	A1
(ii)	Area of shaded region = $\frac{1}{2}(1 + 3) - \int_{-1}^{0} \frac{3}{1-2x} dx$	M1, M1
	$= 2 - \left[\frac{3\ln(1-2x)}{-2} \right]_{-1}^{0}$	M1
	$= 2 - \left[0 + \frac{3}{2}\ln 3 \right]$	
	$= 2 - 1.6479 \quad \approx \quad 0.352$ units^2	A1
12(i)	To prove: BD // FE	
	Proof: Let $\angle TAF$ be θ.	
	$\angle ABD = \angle TAF = \theta$ (alt seg thm)	M1
	$\angle AEF = \angle TAF = \theta$ (alt seg thm)	
	$\therefore \ \angle ABD = \angle AEF = \theta$	
	Using property of corresponding angles, BD // EF (shown)	A1
(ii)	To prove: $\angle FAC = \angle CAE$	
	Proof: Let $\angle BCE = \alpha$	
	$\angle CBD = \angle BCE = \alpha$ (alt $\angle$s, $BD//EF$)	B1
	$\angle FAC = \angle CBD = \alpha$ ($\angle$s in same segment)	B1
	Also, $\angle CAE = \angle BCE = \alpha$ (alt seg thm)	B1
	$\therefore \ \angle FAC = \angle CAE = \alpha$ (shown)	
	END	

Downloaded from SingaKids.com

1 **(i)** A particle moves along the curve $y = \ln\left(x^2 + 1\right)$ in such a way that the *y*-coordinate of the particle is decreasing at a constant rate of 0.2 units per second. Find the rate at which the *x*-coordinate of the particle is changing at the instant when $x = -0.5$. [3]

(ii) Find the *x*-coordinates of the point on the curve where the gradient is **stationary**. [3]

2 **(i)** Solve the equation $\log_3\left(2x+1\right) - \log_3\left(2x-3\right) = 1 + \log_3 \frac{2}{5}$. [4]

(ii) Solve the equation $\ln y + 1 = 2\log_y \mathrm{e}$, giving your answer(s) in terms of e. [5]

3 Given that $y = \mathrm{e}^x \sin x$,

(i) show that $2\frac{\mathrm{d}y}{\mathrm{d}x} - \frac{\mathrm{d}^2 y}{\mathrm{d}x^2} = 2y$. [4]

(ii) Hence, or otherwise, find the value of $\int_0^{\frac{\pi}{3}} \mathrm{e}^x \sin x \, \mathrm{d}x$. [4]

4 Given that the first three terms, in ascending powers of *y*, of the expansion of $\left(a + y\right)^n$, where *a* and *n* are positive real constants, are $64 + 192y + 240y^2$.

(i) By considering the ratio of the coefficients of the first two terms, show that $a = \frac{1}{3}n$. [3]

(ii) Find the value of *a* and of *n*. [4]

5 **(a)** Using the substitution $u = 2^x$, solve the equation $4^{x+1} = 2^x + 3$. [4]

(b) The quantity, *N*, of a radioactive substance, at time *t* years, is given by $N = N_0 \mathrm{e}^{-kt}$, where N_0 and *k* are positive constants.

(i) Sketch the graph of *N* against *t*, labelling any axes intercepts. [2]

(ii) State the significance of N_0. [1]

(iii) The quantity halves every 5 years. Calculate the value of *k*. [3]

Downloaded from SingaKids.com

6 Solutions to this question by accurate drawing will not be accepted.

The coordinates of the points P and Q are $(-5,2)$ and $(7,6)$ respectively. Find


(i) the equation of the line parallel to PQ and passing through the point $(-2,3)$, [3]

(ii) the equation of the perpendicular bisector of PQ. [3]

A point R is such that the shortest distance of R from the line passing through P and Q is $\sqrt{10}$ units.

(iii) Find the area of triangle PQR. [3]

7 The diagram shows a sketch of the curve $y = \mathrm{f}(x)$. The x-coordinates of the maximum and minimum points are $-\alpha$ and α, where $k > 0$.



It is given that $\mathrm{f}'(x) = ax^2 + bx + c$, where a, b and c are real constants. For each of the following, state, with reasons, whether they are positive, negative or zero.


(i) $b^2 - 4ac$, [2]

(ii) $\frac{b}{a}$, [2]

(iii) $\frac{c}{a}$. [2]

Downloaded from SingaKids.com

8 The diagram shows the cross-section of a house with a rooftop *BAC*. The length of *AB* and *AC* are 10 m and 24 m respectively. The angle between *AB* and the horizontal through *A* is θ degrees and $\angle BAC = 90°$.



The base of the house is of length *L* m.

(i) Show that $L = 10\cos\theta + 24\sin\theta$. [2]

(ii) Express *L* in the form $R\sin(\theta + \alpha)$, where $R > 0$ and α is an acute angle. [4]

(iii) Find the longest possible base of the house and the corresponding value of θ. [3]

9 **(a)** The equation of a curve is $y = \dfrac{2x}{1+x}$.

(i) Find the equation of the tangent to the curve at point $P(1,1)$. [4]

(ii) The tangent cuts the axes at *Q* and *R* respectively. Find the area of triangle *OPQ*. [2]

(b) A curve has equation $y = \mathrm{f}(x)$, where $\mathrm{f}(x) = \dfrac{1}{3}x^3 - 2x^2 + 13x + 5$.

Determine, with explanation, whether f is an increasing or decreasing function. [4]


Downloaded from SingaKids.com

10 (a) (i) Solve the equation $\left|x^2-3x+2\right|+x=1$. [3]

(ii) What can be deduced about the number of points of intersections of the graphs of $y=\left|x^2-3x+2\right|$ and $y=-x+1$? [1]

(iii) Hence, on a single diagram, sketch the graphs of $y=\left|x^2-3x+2\right|$ and $y=-x+1$, indicating the coordinates of any axial intercepts and turning point. [4]

(b) The diagram shows part of the graph of $y=\left|k-x\right|$, where k is a constant.



A line $y=mx+c$ is drawn to determine the number of solutions to the equation $\left|k-x\right|=mx+c$.

(i) If $m=1$, state the range of values of c, in terms of k, such that the equation has one solution. [1]

(ii) If $c=0$, state the range of values of m such that the equation has no solutions. [2]

11 (a) State the principal range of $\sin^{-1}x$, leaving your answers in terms of π. [1]

(b) (i) Prove that $\dfrac{1+\tan x}{1-\tan x}=\sec 2x+\tan 2x$. [5]

(ii) Hence find the reflex angle x such that $3\sec 2x+3\tan 2x=1$. [3]

(c) A buoy floats and its height above the seabed, h m, is given by $h=a\cos bt+c$, where t is time measured in hours from 0000 hours and a, b and c are constants. The least height of the buoy above seabed is 180 metres and is recorded at 0000 hours. The greatest height of the buoy above seabed is 196 metres and is first recorded at 0600 hours.

(i) Find the values of a, b and c. [3]

(ii) Using values found in **(i)**, sketch the graph of $h=a\cos bt+c$ for $0\le t\le 24$. [2]

(ii) The buoy floats above the top of a huge rock first at 0500 hours. State the number of hours in each day that the buoy is above the rock. [1]

END OF PAPER

Downloaded from SingaKids.com

Question 1

(i)	0.25 units/s
(ii)	$x = \pm 1$

Question 2

(i)	$x = \dfrac{23}{2}$
(ii)	$y = \mathrm{e}^{-2}$ or $y = \mathrm{e}$


Question 3

(ii)	$1.02 (3\text{s.f.})$

Question 4

(ii)	$n = 6, a = 2$

Question 5

(a)	$x = 0$
(a)(i)	
(ii)	It represents the initial amount of radioactive substance.
(iii)	0.139

Question 6

(i)	$y = \dfrac{1}{3}x + 3\dfrac{2}{3}$
(ii)	$y = -3x + 7$
(iii)	20 units^2

Question 7

(i)	$b^2 - 4ac > 0$.
(ii)	$\dfrac{b}{a} > 0$
(iii)	$\dfrac{c}{a} < 0$

Question 8

(ii)	$L = 26\sin(\theta + 22.6°)$
(iii)	Longest possible base is 26 m.

SGTREEPAPERS.COM


Downloaded from SingaKids.com

	$\theta = 67.4°\,(1\text{ d.p.})$

Question 9

(a)(i)	$y = \frac{1}{2}x + \frac{1}{2}$
(ii)	$\frac{1}{4}$ units^2

Question 10

(a)(i)	$x = 1$
(ii)	The line $y = -x + 1$ is tangential to $y = \lvert x^2 - 3x + 2 \rvert$.
(ii)	
(b)(i)	$c > -k$
(ii)	$-1 \le m < 0$

Question 11

(a)	$-\frac{\pi}{2} \le \sin^{-1} x \le \frac{\pi}{2}$
(b)(ii)	$x = 333.3°\,(1\text{ d.p.})$
(c)(i)	$a = -8, b = \frac{\pi}{6}, c = 188$
(iii)	4 hours

Downloaded from SingaKids.com

1 **(i)** A particle moves along the curve $y = \ln\left(x^2 + 1\right)$ in such a way that the y-coordinate of the particle is decreasing at a constant rate of 0.2 units per second. Find the rate at which the x-coordinate of the particle is changing at the instant when $x = -0.5$. [3]

(ii) Find the x-coordinates of the point on the curve where the gradient is **stationary**. [3]

(i)	$\frac{dy}{dx} = \frac{2x}{x^2+1}$ $\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$ $-0.2 = \frac{2(-0.5)}{(-0.5)^2+1} \times \frac{dx}{dt}$ $\Rightarrow \frac{dx}{dt} = 0.25$ units/s	**B1** **M1** **A1**
(ii)	$\frac{d^2y}{dx^2} = \frac{(x^2+1)(2)-2x(2x)}{(x^2+1)^2}$ $= \frac{2-2x^2}{(x^2+1)^2}$ $\frac{d^2y}{dx^2} = 0$ $2-2x^2 = 0$ $x = \pm 1$	**√M1** **M1** **A1**

SGFREEPAPERS.COM

Downloaded from SingaKids.com

2 **(i)** Solve the equation $\log_3(2x+1) - \log_3(2x-3) = 1 + \log_3 \frac{2}{5}$. [4]

(ii) Solve the equation $\ln y + 1 = 2\log_y \mathrm{e}$, giving your answer(s) in terms of e. [5]

(i)	$\log_3(2x+1) - \log_3(2x-3) = 1 + \log_3 \frac{2}{5}$	
	$\log_3 \frac{2x+1}{2x-3} = \log_3\left(3 \times \frac{2}{5}\right)$	**B1**, **B1**
	$\frac{2x+1}{2x-3} = \frac{6}{5}$	**M1** – remove log
	$10x + 5 = 12x - 18$	
	$2x = 23$	
	$x = \frac{23}{2}$	**A1**
(ii)	$\ln y + 1 = 2\log_y \mathrm{e}$	
	$\ln y + 1 = \frac{2}{\ln y}$	**B1** – change base
	$(\ln y)^2 + \ln y - 2 = 0$	**B1**
	$(\ln y + 2)(\ln y - 1) = 0$	**M1** – attempt to solve
	$\ln y = -2$ or 1	
	$y = \mathrm{e}^{-2}$ or $y = \mathrm{e}$	**A2**

Downloaded from SingaKids.com

3 Given that $y = e^x \sin x$,

(i) show that $2\frac{dy}{dx} - \frac{d^2y}{dx^2} = 2y$. [4]

(ii) Hence, or otherwise, find the value of $\int_0^{\frac{\pi}{3}} e^x \sin x \, dx$. [4]

(i)	$y = e^x \sin x$ $\frac{dy}{dx} = e^x \sin x + e^x \cos x$ $\frac{d^2y}{dx^2} = e^x \sin x + e^x \cos x - e^x \sin x + e^x \cos x$ $= 2e^x \cos x$ $-\frac{d^2y}{dx^2} + 2\frac{dy}{dx} = -2e^x \cos x + 2\left(e^x \sin x + e^x \cos x\right)$ $= 2e^x \sin x$ $= 2y$ $2\frac{dy}{dx} - \frac{d^2y}{dx^2} = 2y$	**M1** – product rule **B1** **M1** – product rule **M1** **a.g.**
(ii)	$-\frac{d^2y}{dx^2} + 2\frac{dy}{dx} = 2y$ $\therefore -\frac{dy}{dx} + 2y = 2\int e^x \sin x \, dx$ $\Rightarrow -e^x \sin x - e^x \cos x + 2e^x \sin x = 2\int e^x \sin x \, dx$ $\therefore \int e^x \sin x \, dx = \frac{1}{2}\left(e^x \sin x - e^x \cos x\right) + c$ $\int_0^{\frac{\pi}{3}} e^x \sin x \, dx = \left[\frac{1}{2}\left(e^x \sin x - e^x \cos x\right)\right]_0^{\frac{\pi}{3}}$ $= 1.02 (3\text{s.f.})$	**M1** – integration **B1** – making integral the subject **M1** – substitution of limits **A1**

SGFREEPAPERS.COM

Downloaded from SingaKids.com

4 Given that the first three terms, in ascending powers of y, of the expansion of $(a+y)^n$, where a and n are positive real constants, are $64+192y+240y^2$.

(i) By considering the ratio of the coefficients of the first two terms, show that $a=\frac{1}{3}n$. [3]

(ii) Find the value of a and of n. [4]

(i)	$(a+y)^n = a^n + na^{n-1}y + \frac{n(n-1)}{2}a^{n-2}y^2 + \ldots$ By comparing coefficents, $a^n = 64 \quad ---(1)$ $na^{n-1} = 192 \quad ---(2)$ $\frac{n(n-1)}{2}a^{n-2} = 240 \quad ---(3)$ $\frac{(1)}{(2)}: \frac{a}{n} = \frac{64}{192} = \frac{1}{3} \Rightarrow a = \frac{1}{3}n ---(4)$	**B1** – award for first two terms **M1**, **A1**
(ii)	$\frac{(2)}{(3)}: \frac{2a}{n-1} = \frac{192}{240} = \frac{4}{5} \Rightarrow a = \frac{2}{5}(n-1) ---(5)$ $(4)=(5):$ $\frac{1}{3}n = \frac{2}{5}(n-1)$ $5n = 6n - 6$ $n = 6$ $\Rightarrow a = 2$	**√M1** **√M1** – simultaneous eqn **A1** **A1**

Downloaded from SingaKids.com


5 **(a)** Using the substitution $u = 2^x$, solve the equation $4^{x+1} = 2^x + 3$. [4]

(b) The quantity, N, of a radioactive substance, at time t years, is given by $N = N_0 e^{-kt}$, where N_0 and k are positive constants.

(i) Sketch the graph of N against t, labelling any axes intercepts. [2]

(ii) State the significance of N_0. [1]

(iii) The quantity halves every 5 years. Calculate the value of k. [3]

(a)	$4u^2 = u + 3$ $4u^2 - u - 3 = 0$ $(4u + 3)(u - 1) = 0$ $u = 1$ or $-\frac{3}{4}$ $x = 0$ or $2^x = -\frac{3}{4}$ (no solutions)	**B1** **M1** **A1, A1**
(a)(i)		**B1** – shape **B1** – $t > 0$ and label N_0
(ii)	It represents the initial amount of radioactive substance.	**B1**
(iii)	$\frac{1}{2} N_0 = N_0 e^{-k(5)}$ $\frac{1}{2} = e^{-5k}$ $-5k = \ln \frac{1}{2} = -\ln 2$ $t = \frac{\ln 2}{5} \approx 0.139$	**M1** **M1** **A1**

SGFREEPAPERS.COM

Downloaded from SingaKids.com

6 Solutions to this question by accurate drawing will not be accepted.

The coordinates of the points P and Q are $(-5,2)$ and $(7,6)$ respectively. Find

(i) the equation of the line parallel to PQ and passing through the point $(-2,3)$. [3]

(ii) the equation of the perpendicular bisector of PQ. [3]


A point R is such that the shortest distance of R from the line passing through P and Q is $\sqrt{10}$ units.

(iii) Find the area of triangle OQR. [3]

(i)	$m_{PQ} = \dfrac{6-2}{7-(-5)} = \dfrac{1}{3}$	**B1**
	$y-3 = \dfrac{1}{3}\left[x-(-2)\right]$	**M1**
	$y = \dfrac{1}{3}x + 3\dfrac{2}{3}$	**A1**
(ii)	Midpoint of $PQ = \left(\dfrac{-5+7}{2}, \dfrac{2+6}{2}\right) = (1,4)$	**B1**
	Gradient of perpendicular bisector $= -3$ $y-4 = -3(x-1)$	**√M1**
	$y = -3x+7$	**A1**
(iii)	$PQ = \sqrt{(7-(-5))^2 + (6-2)^2} = 4\sqrt{10}$ units	**M1**
	Area $= \dfrac{1}{2}\left(4\sqrt{10}\right)\sqrt{10}$	**√M1**
	$= 20$ units^2	**A1**

Downloaded from SingaKids.com

7 The diagram shows a sketch of the curve $y = \mathrm{f}(x)$. The x-coordinates of the minimum and maximum points are α and $-\alpha$, where $\alpha > 0$.



It is given that $\mathrm{f}'(x) = ax^2 + bx + c$, where a, b and c are real constants. For each of the following, state, with reasons, whether they are positive, negative or zero.

(i) $b^2 - 4ac$, [2]

(ii) $\frac{b}{a}$, [2]


(iii) $\frac{c}{a}$. [2]

SGFREEPAPERS.COM

(i)	Since there are two stationary points, $\mathrm{f}'(x) = 0$ has two real roots, therefore $b^2 - 4ac > 0$.	**M1** **A1**
(ii)	Since $\lvert\alpha\rvert > \lvert\beta\rvert$ and $\alpha < 0$, $\alpha + \beta < 0$, $\therefore \frac{b}{a} = -(\alpha + \beta) > 0$	**M1** **A1**
(iii)	Since $\alpha < 0$ and $\beta > 0$, $\alpha\beta < 0$, $\therefore \frac{c}{a} = \alpha\beta < 0$	**M1** **A1**

Downloaded from SingaKids.com

8 The diagram shows the cross-section of a house with a rooftop *BAC*. The length of *AB* and *AC* are 10 m and 24 m respectively. The angle between *AB* and the horizontal through *A* is θ degrees and $\angle BAC = 90°$.



The base of the house is of length *L* m.

(i) Show that $L = 10\cos\theta + 24\sin\theta$. [2]

(ii) Express *L* in the form $R\sin(\theta + \alpha)$, where $R > 0$ and α is an acute angle. [4]

(iii) Find the longest possible base of the house and the corresponding value of θ. [3]

(i)	Let the point vertically above *B* and *C* be *M* and *N* respectively. $\angle ACN = 90°$ $AM = 10\cos\theta$ and $AN = 24\sin\theta$ $L = MN = 10\cos\theta + 24\sin\theta$	**B1**, **B1**
(ii)	$R = \sqrt{10^2 + 24^2}$ $= 26$ $\alpha = \tan^{-1}\left(\frac{10}{24}\right)$ $= 22.620°\,(3\text{ d.p.})$ $L = 26\sin(\theta + 22.6°)$	**M1** **A1** **M1** **A1**
(iii)	Longest possible base is 26 m. $\theta + 22.620° = 90°$ $\theta = 67.4°\,(1\text{ d.p.})$	**B1** **√M1** **A1**

Downloaded from SingaKids.com

9 **(a)** The equation of a curve is $y = \dfrac{2x}{1+x}$.

(i) Find the equation of the tangent to the curve at point $P(1,1)$. [4]

(ii) The tangent cuts the axes at Q and R respectively. Find the area of triangle PQR. [2]

(b) A curve has equation $y = \mathrm{f}(x)$, where $\mathrm{f}(x) = \dfrac{1}{3}x^3 - 2x^2 + 13x + 5$.

Determine, with explanation, whether f is an increasing or decreasing function. [4]

(a) **(i)**	$\dfrac{dy}{dx} = \dfrac{(1+x)(2)-(2x)(1)}{(1+x)^2}$ $= \dfrac{2}{(1+x)^2}$	**M1**
	$\left.\dfrac{dy}{dx}\right\|_{x=1} = \dfrac{1}{2}$	**B1**
	Equation of Tangent: $y-1 = \dfrac{1}{2}(x-1) \Rightarrow y = \dfrac{1}{2}x + \dfrac{1}{2}$	**M1** – substitution of point **A1**
(ii)	$Q(-1,0)$ and $R\left(0,\dfrac{1}{2}\right)$	**√B1**
	Area of Triangle $= \dfrac{1}{2}(1)\left(\dfrac{1}{2}\right) = \dfrac{1}{4}$ units^2	**√B1**
(b)	$\mathrm{f}'(x) = x^2 - 4x + 13$	**B1**
	$= (x-2)^2 - 2^2 + 13$	**M1** - complete the square
	$= (x-2)^2 + 9$	
	$(x-2)^2 \geq 0 \Rightarrow (x-2)^2 + 9 > 0$	**M1**
	$\therefore \mathrm{f}'(x) > 0$, f is an increasing function.	**A1**

SGFREEPAPERS.COM


Downloaded from SingaKids.com

10 (a) (i) Solve the equation $\left|x^2-3x+2\right|+x=1$. [3]

(ii) What can be deduced about the number of points of intersections of the graphs of $y=\left|x^2-3x+2\right|$ and $y=-x+1$? [1]

(iii) Hence, on a single diagram, sketch the graphs of $y=\left|x^2-3x+2\right|$ and $y=-x+1$, indicating any axial intercepts. [4]

(b) The diagram shows part of the graph of $y=\left|k-x\right|$, where k is a constant.




A line $y=mx+c$ is drawn to determine the number of solutions to the equation $\left|k-x\right|=mx+c$.

(i) If $m=1$, state the range of values of c, in terms of k, such that the equation has one solution. [1]

(ii) If $c=0$, state the range of values of m such that the equation has no solutions. [2]

(a) (i)	$x^2-3x+2=-x+1$ or $x^2-3x+2=-(-x+1)$ $x^2-2x+1=0$ $x^2-4x+3=0$ $(x-1)^2=0$ $(x-3)(x-1)=0$ $x=1$ $x=1$ or $x=3$(rejected)	**M1** **A1, A1**
(ii)	The line $y=-x+1$ is tangential to $y=\left\|x^2-3x+2\right\|$.	**B1**

Downloaded from SingaKids.com

(ii)		**B1** - modulus graph **B1** – axial intercepts & turning point **B1** – line with intercepts **B1** – line tangent to modulus
(b)(i)	$c > -k$	**B1**
(ii)	$-1 \le m < 0$	**A1**, **A1**

SGFREEPAPERS.COM

Downloaded from SingaKids.com

11 (a) State the principal range of $\sin^{-1} x$, leaving your answers in terms of π. [1]

(b) (i) Prove that $\dfrac{1+\tan x}{1-\tan x} = \sec 2x + \tan 2x$. [5]

(ii) Hence find the reflex angle x such that $\sec 2x + \tan 2x = \dfrac{1}{3}$. [3]

(c) A buoy floats and its height above the seabed, h m, is given by $h = a\cos bt + c$, where t is time measured in hours from 0000 hours and a, b and c are constants. The least height of the buoy above seabed is 180 metres and is recorded at 0000 hours. The greatest height of the buoy above seabed is 196 metres and is first recorded at 0600 hours.


(i) Find the values of a, b and c. [3]

(ii) Using values found in **(i)**, sketch the graph of $h = a\cos bt + c$ for $0 \le t \le 24$. [2]

(ii) The buoy floats above the top of a huge rock first at 0500 hours. State the number of hours in each day that the buoy is above the rock. [1]

(a)	$-\dfrac{\pi}{2} \le \sin^{-1} x \le \dfrac{\pi}{2}$	**B1**
(b) (i)	$\dfrac{1+\tan x}{1-\tan x} = \dfrac{1+\dfrac{\sin x}{\cos x}}{1-\dfrac{\sin x}{\cos x}}$	
	$= \dfrac{\cos x + \sin x}{\cos x - \sin x}$	**M1**
	$= \dfrac{(\cos x + \sin x)^2}{\cos^2 x - \sin^2 x}$	**M1**
	$= \dfrac{1 + 2\sin x \cos x}{\cos 2x}$	**M1** – double angle
	$= \dfrac{1 + \sin 2x}{\cos 2x}$	**M1** – double angle
	$= \sec 2x + \tan 2x$	**A1**
(ii)	$\dfrac{1+\tan x}{1-\tan x} = \dfrac{1}{3}$	**M1**
	$3 + 3\tan x = 1 - \tan x$	
	$4\tan x = -2$	
	$\tan x = -\dfrac{1}{2}$	
	$\alpha = 26.565°\,(3\text{ d.p.})$	**B1**
	$x = 333.3°\,(1\text{ d.p.})$	**A1**

Downloaded from SingaKids.com

(c) **(i)**	$\frac{196-180}{2}=8 \Rightarrow a=-8$ $c=\frac{196+180}{2}=188$ $b=\frac{2\pi}{12}=\frac{\pi}{6}$	**B1** **B1** **B1**
(ii)		**B1** – shape **B1** - points
(iii)	4 hours	**B1**

SGFREEPAPERS.COM

Downloaded from SingaKids.com