ZHONGHUA SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2019
SECONDARY 4E, 5N

Candidate's Name	Class	Register Number

ADDITIONAL MATHEMATICS

4047/01

PAPER 1

17 Sep 2019
2 hours

Candidates answer on the question paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number in the spaces at the top of this page..
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer all questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

For Examiner's Use
80

This question paper consists of 16 printed pages (including this cover page)

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \cdots + \binom{n}{r}a^{n-r}b^r + \cdots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!\,r!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$
$$\sec^2 A = 1 + \tan^2 A.$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$
$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$
$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$
$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$
$$\sin 2A = 2 \sin A \cos A$$
$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$
$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for ΔABC


$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}ab \sin C$$

Downloaded from SingaKids.com

1 The diagram shows a straight line passing through $P(0,b)$ and $Q(a,-2b)$




Given that the gradient of PQ is $-\frac{3}{7}$ and that the distance $PQ = 3\sqrt{58}$, find the value of a and of b. [5]

Downloaded from SingaKids.com

2 The equation of a curve is $f(x) = |3cos2x| - 2$

(i) State the minimum and maximum values of $f(x)$. [2]

(ii) Sketch the graph of $f(x) = |3cos2x| - 2$ for $0° \leq x \leq 180°$. [3]



Downloaded from SingaKids.com


3 The expression $ax^3 - 4x^2 + bx + 6$ is exactly divisible by $x^2 - x - 2$.

(i) Find the value of a and of b. [4]

(ii) Using these values of a and of b, solve the equation $ax^3 - 4x^2 + bx + 6 = 0$. [2]

Downloaded from SingaKids.com

4



In the diagram, TE and TC are tangents to the circle at A and B respectively.
BD is parallel to TE.

(i) Prove that $AB = AD$. [3]

(ii) Prove that triangle TAB is similar to triangle ABD. [3]

Downloaded from SingaKids.com

5 The function f is defined by $f(x) = \frac{5-x^2}{x^2+3}$, $x > 0$

(i) Explain, with working, whether f is an increasing or decreasing function. [4]

(ii) A point P moves along the curve $y = f(x)$ in such a way that the y-coordinate of P is increasing at a rate of 0.2 units per second. Find the rate of change of the x-coordinate of P when $x = 4$. [2]

Downloaded from SingaKids.com

6 At m minutes after an oven is switched on, its temperature, T°C, is given by $T = 200 - 175e^{-km}$. The oven reached a temperature of 150°C after 16 minutes 15 seconds.

(i) State the initial temperature of the oven. [1]

(ii) Estimate the temperature after 10 minutes. [3]

(iii) "If the oven is switched on for a very long time, it will never exceed a certain temperature." Do you agree with the statement? Justify your answer with clear explanation. [2]

Downloaded from SingaKids.com

7 (i) Express $-2x^2 + 3x + 2$ in the form $a(x + b)^2 + c$ where a, b and c are constants. [2]

(ii) Use your answer from part (i) to explain why the curves with equations $y = -2x^2 + 3x + 2$ and $(x + 5)^2 + (y - 9)^2 = 30$ will not intersect. [4]

Downloaded from SingaKids.com

8 (i) Form a quadratic equation for which the sum of roots is 4 and the sum of the squares of the roots is 20. [3]

(ii) Given that the equation $(2m+3)x^2-(8m+8)x+8m+3=0$, find the value of m, for which

(a) one root is the negative of the other, [2]

(b) one root is the reciprocal of the other. [2]

Downloaded from SingaKids.com


9 Without using a calculator,

(i) show that $sin105° = \frac{1+\sqrt{3}}{2\sqrt{2}}$, [2]

(ii) hence, express $1 + cot^2 105°$ in the form $a + b\sqrt{3}$, where a and b are integers. [5]

Downloaded from SingaKids.com

10



The figure shows a piece of iron bar of length 8 metres, bent to form a pentagonal window frame $PQRST$.

$PT = QR = y$ metres, $RS = ST = 2x$ metres and angle PTS =angle $QRS = 150°$.

(i) Show that $y = 4 - 3x$. [3]

(ii) Express the area, A, enclosed by the frame, in terms of x. [3]

Downloaded from SingaKids.com

10 (iii) Given that x can vary, find the exact value of x for which the area enclosed by the frame is a maximum. [2]

Downloaded from SingaKids.com

11 (i) Prove the identity $\frac{tanA-cotA}{tanA+cotA}+1 = 2sin^2A$ [3]

(ii) Hence, solve the equation $\frac{tanA-cotA}{tanA+cotA}+1 = 5sinAcosA$ for $0° < A < 360°$ [5]

Downloaded from SingaKids.com

12 A particle starts from rest, travels in a straight line so that t is the time in seconds after passing a fixed point O. Its velocity, $v\,m/s$, is given by $v = 6t - 2t^2$. The particle comes to instantaneous rest at A.

(i) Find the acceleration of the particle at A. [3]

(ii) Find the maximum velocity of the particle. [2]

Downloaded from SingaKids.com

12 (iii) Find the total distance travelled by the particle during the first 5 seconds. [5]

End of Paper

Downloaded from SingaKids.com

ZHONGHUA SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2019
SECONDARY 4E, 5N

Candidate's Name	Class	Register Number
Solutions		

ADDITIONAL MATHEMATICS

4047/01

PAPER 1

17 Sep 2019
2 hours

Candidates answer on the question paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number in the spaces at the top of this page..
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.


The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is **80**.

For Examiner's Use
80

This question paper consists of 16 printed pages (including this cover page)

Downloaded from SingaKids.com

1 The diagram shows a straight line passing through $P(0, b)$ and $Q(a, -2b)$



Given that the gradient of PQ is $-\frac{3}{7}$ and that the distance $PQ = 3\sqrt{58}$, find the value of a and of b. [5]

$$Grad\ PQ = \frac{b-(-2b)}{0-a} = -\frac{3}{7}$$ M1

$$\frac{3b}{-a} = -\frac{3}{7}$$

$$a = 7b$$ M1

$$PQ = 3\sqrt{58}$$

$$\sqrt{(0-a)^2 + (b+2b)^2} = 3\sqrt{58}$$ M1

$$a^2 + 9b^2 = 9(58)$$

$$(7b)^2 + 9b^2 = 9(58)$$

$$58b^2 = 522$$

$$b^2 = 9$$

$$b = 3\ (b > 0\ as\ shown\ in\ diagram)$$ A1

$$\therefore a = 21$$ A1

Downloaded from SingaKids.com


2 The equation of a curve is $f(x) = |3cos2x| - 2$

(i) State the minimum and maximum values of $f(x)$. [2]

$Min\ of\ f(x) = 1$ B1

$Max\ of\ f(x) = -2$ B1

(ii) Sketch the graph of $f(x) = |3cos2x| - 2$ for $0° \leq x \leq 180°$. [3]



One complete cycle B1

Starts and ends at f(x)=1 B1

Correct curvature B1

Downloaded from SingaKids.com

3 The expression $ax^3 - 4x^2 + bx + 6$ is exactly divisible by $x^2 - x - 2$.

(i) Find the value of a and of b. [4]

Let $f(x) = ax^3 - 4x^2 + bx + 6$

$x^2 - x - 2 = (x - 2)(x + 1)$

$f(2) = 8a - 16 + 2b + 6 = 0$ M1

$f(-1) = -a - 4 - b + 6 = 0$ M1

$4a + b = 5$

$a + b = 2$

$3a = 3$

$a = 1$ A1

$b = 1$ A1

(ii) Using these values of a and of b, solve the equation $ax^3 - 4x^2 + bx + 6 = 0$. [2]


$x^3 - 4x^2 + x + 6 = 0$

By inspection, $(x - 3)$ is the linear factor of $f(x)$

$(x^2 - x - 2)(x - 3) = 0$ B1

$(x - 2)(x + 1)(x - 3) = 0$


$x = -1, 2, 3$ A1



$y = x^3 - 4x^2 + x + 6 = (x - 2)(x + 1)(x - 3)$

Downloaded from SingaKids.com

4



In the diagram, TE and TC are tangents to the circle at A and B respectively.
BD is parallel to TE.

(i) Prove that $AB = AD$. [3]

$$Let \measuredangle ABD = \theta$$

$$\measuredangle TAB = \theta \ (alternate\ angles, BD \parallel TE)$$ B1

$$\measuredangle ADB = \measuredangle TAB = \theta \ (tangent - chord\ theorem)$$ B1

Since $\measuredangle ADB = \measuredangle ABD = \theta$,

$$therefore, \Delta ABD\ is\ an\ isosceles\ triangle\ with\ AB = AD$$ B1

(ii) Prove that triangle TAB is similar to triangle ABD. [3]

In triangles TAB and ABD,

$$\measuredangle TAB = \measuredangle ABD = \theta \ (alternate\ angles, BD \parallel TE)$$ B1

$$\measuredangle TBA = \measuredangle ADB = \theta \ (tangent - chord\ theorem)$$ B1

$$\measuredangle ATB = \measuredangle BAD \ (angle\ sum\ of\ triangles)$$

Since all 3 pairs of corresponding angles are equal, triangle TB is similar to triangle ABD. B1

Downloaded from SingaKids.com

5 The function f is defined by $f(x) = \frac{5-x^2}{x^2+3}$, $x > 0$


(i) Explain, with working, whether f is an increasing or decreasing function. [4]

$$f'(x) = \frac{(x^2+3)(-2x)-(5-x^2)(2x)}{(x^2+3)^2}$$

$$= \frac{-2x^3 - 6x - 10x + 2x^3}{(x^2+3)^2}$$

$$= \frac{-16x}{(x^2+3)^2}$$

Since $x > 0, -16x < 0$ *and* $(x^2+3)^2 > 0$, $f'(x) < 0$ and thus $f(x)$ is a decreasing function for $x > 0$.



$$f(x) = \frac{5-x^2}{x^2+3}$$

(ii) A point P moves along the curve $y = f(x)$ in such a way that the y-coordinate of P is increasing at a rate of 0.2 units per second. Find the rate of change of the x-coordinate of P when $x = 4$. [2]

$$\frac{dy}{dt} = +0.2$$

$$\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$$

$$0.2 = \frac{-16(4)}{(4^2+3)^2} \times \frac{dx}{dt} \quad \text{M1}$$

$$\frac{dx}{dt} = -1.1281 = -1.13 \textit{units per sec} \quad \text{A1}$$

Downloaded from SingaKids.com

6 At m minutes after an oven is switched on, its temperature, T°C, is given by $T = 200 - 175e^{-km}$. The oven reached a temperature of 150°C after 16 minutes 15 seconds.

(i) State the initial temperature of the oven. [1]

When $m = 0, T = 200 - 175e^{-0}$

$= 25°C$ B1

(ii) Estimate the temperature after 10 minutes. [3]

$$150 = 200 - 175e^{-k(16.25)}$$

$$175e^{-k(16.25)} = 50$$

$$e^{-k(16.25)} = \frac{2}{7}$$ M1

$$-16.25k = \ln\frac{2}{7}$$

$$k = \frac{\ln\frac{2}{7}}{-16.25} \text{ or } 0.077093$$ A1

When $m = 10mins$, $T = 200 - 175e^{-k(10)}$

$= 119.04°C = 119°C\ (3\ sig\ fig)$ A1


(iii) "If the oven is switched on for a very long time, it will never exceed a certain temperature." Do you agree with the statement? Justify your answer with clear explanation. [2]

$$T = 200 - 175e^{-km}$$

$$T = 200 - \frac{175}{e^{km}}$$

Yes, I agree with the statement.

Since $k > 0, m > 0$, and when m becomes large, $\frac{175}{e^{km}}$ becomes a very small positive value. Therefore, T will never exceed 200°C. B2



$T = 200 - \frac{175}{e^{km}}$

Downloaded from SingaKids.com

7 (i) Express $-2x^2+3x+2$ in the form $a(x+b)^2+c$ where a, b and c are constants. [2]

$$-2x^2+3x+2 = -2\left[x^2-\frac{3}{2}x-1\right]$$

$$= -2\left[\left(x-\tfrac{3}{4}\right)^2-\tfrac{9}{16}-1\right] \qquad \left(x-\tfrac{3}{4}\right)^2 \text{ seen} \quad \text{B1}$$

$$= -2\left[\left(x-\frac{3}{4}\right)^2-\frac{25}{16}\right]$$

$$= -2\left(x-\frac{3}{4}\right)^2+\frac{25}{8} \qquad \text{B1}$$

(ii) Use your answer from part (i) to explain why the curves with equations $y=-2x^2+3x+2$ and $(x+5)^2+(y-9)^2=30$ will not intersect. [4]


$y=-2x^2+3x+2=-2\left(x-\frac{3}{4}\right)^2+\frac{25}{8}$ has a maximum value at $y=\frac{25}{8}$ *or* 3.125 B1

$(x+5)^2+(y-9)^2=30$ is a circle with center $(-5,9)$ and radius $\sqrt{30}$. M1

for either centre or radius

The minimum value of $y=9-\sqrt{30}=3.522$ B1

Since maximum of $y=-2x^2+3x+2$ < minimum of $(x+5)^2+(y-9)^2=30$, the 2 curves will not intersect. B1



Downloaded from SingaKids.com

8 (i) Form a quadratic equation for which the sum of roots is 4 and the sum of the squares of the roots is 20. [3]

$$\alpha + \beta = 4$$
$$\alpha^2 + \beta^2 = 20 \quad \text{M1}$$
$$\Rightarrow (\alpha + \beta)^2 - 2\alpha\beta = 20$$
$$(4)^2 - 2\alpha\beta = 20$$
$$\alpha\beta = -2 \quad \text{M1}$$

$$\therefore \textit{the quadratic equation is } x^2 - 4x - 2 = 0 \quad \text{A1}$$

(ii) Given that the equation $(2m+3)x^2 - (8m+8)x + 8m + 3 = 0$, find the value of m, for which

(a) one root is the negative of the other, [2]

Let the roots be α and $-\alpha$.

Sum of roots: $\alpha + (-\alpha) = \frac{8m+8}{2m+3}$ attempt to form eqn M1

$$\frac{8m+8}{2m+3} = 0$$
$$8m + 8 = 0$$
$$m = -1 \quad \text{A1}$$

(b) one root is the reciprocal of the other. [2]

Let the roots be α and $\frac{1}{\alpha}$.

Product of roots: $\alpha \times \frac{1}{\alpha} = \frac{8m+3}{2m+3}$ attempt to form eqn M1

$$8m + 3 = 2m + 3$$
$$6m = 0$$
$$m = 0 \quad \text{A1}$$

Downloaded from SingaKids.com

9 Without using a calculator,

(i) show that $sin105° = \frac{1+\sqrt{3}}{2\sqrt{2}}$, [2]

$$\sin 105° = \sin(45° + 60°) \quad \text{M1}$$

$$= \sin 45° \cos 60° + \cos 45° \sin 60°$$

$$= \left(\frac{1}{\sqrt{2}}\right)\left(\frac{1}{2}\right) + \left(\frac{1}{\sqrt{2}}\right)\left(\frac{\sqrt{3}}{2}\right) \quad \text{A1}$$

$$= \frac{1+\sqrt{3}}{2\sqrt{2}} \quad \text{AG}$$

(ii) hence, express $1 + cot^2 105°$ in the form $a + b\sqrt{3}$, where a and b are integers. [5]

$$1 + cot^2 105° = cosec^2 105° \quad \text{M1}$$

$$= \frac{1}{sin^2 105°}$$

$$= \left(\frac{2\sqrt{2}}{1+\sqrt{3}}\right)^2 \quad \text{M1}$$


$$= \frac{8}{4+2\sqrt{3}} \quad \text{For correctly squaring } 1+\sqrt{3} \quad \text{M1}$$

$$= \frac{4}{2+\sqrt{3}} \times \frac{2-\sqrt{3}}{2-\sqrt{3}} \quad \text{For rationalising} \quad \text{M1}$$

$$= 8 - 4\sqrt{3} \quad \text{A1}$$

Downloaded from SingaKids.com

10



The figure shows a piece of iron bar of length 8 metres, bent to form a pentagonal window frame $PQRST$.

$PT = QR = y$ metres, $RS = ST = 2x$ metres and angle PTS =angle $QRS = 150°$.

(i) Show that $y = 4 - 3x$. [3]

PQRT is a rectangle

$\Rightarrow \measuredangle STR = \measuredangle SRT = 150° - 90° = 60°$ M1

$\therefore \measuredangle TSR = 180° - 2(60°) = 60°$

Since TSR is an equilateral triangle,

$TR = 2x$

$\Rightarrow PQ = 2x$ M1

$3(2x) + 2y = 8$ B1

$2y = 8 - 6x$

$y = 4 - 3x$ AG

(ii) Express the area, A, enclosed by the frame, in terms of x. [3]

$A = 2x(y) + \frac{1}{2}(2x)(2x)\sin 60°$ For ½(absin c) o.e. B1

$= 2x(4 - 3x) + 2x^2\left(\frac{\sqrt{3}}{2}\right)$ For substituting y B1

$= 8x - 6x^2 + \sqrt{3}x^2$ A1


Downloaded from SingaKids.com

10 (iii) Given that x can vary, find the exact value of x for which the area enclosed by the frame is a maximum. [2]

$$A = 8x - 6x^2 + \sqrt{3}x^2$$
$$\frac{dA}{dx} = 8 - 12x + 2\sqrt{3}x \quad \text{M1}$$

For max area, $\frac{dA}{dx} = 0$

$$8 - 12x + 2\sqrt{3}x = 0$$
$$12x - 2\sqrt{3}x = 8$$
$$x = \frac{8}{12 - 2\sqrt{3}}$$
$$= \frac{4}{6 - \sqrt{3}} \quad \text{A1}$$



$A = 8x - 6x^2 + \sqrt{3}x^2$

Downloaded from SingaKids.com

11 (i) Prove the identity $\frac{tanA-cotA}{tanA+cotA}+1=2sin^2A$ [3]

$$LHS=\frac{tanA-cotA}{tanA+cotA}+1$$

$$=\frac{\left(\frac{\sin A}{\cos A}-\frac{\cos A}{\sin A}\right)}{\left(\frac{\sin A}{\cos A}+\frac{\cos A}{\sin A}\right)}+1$$ For converting tan A or cot A B1

$$=\frac{sin^2A-cos^2A}{sin^2A+cos^2A}+1$$ M1

$$=(sin^2A-cos^2A)+(sin^2A+cos^2A)$$ For $sin^2A+cos^2A=1$ and leading to final answer. B1

$$=2sin^2A$$

$$=RHS$$

(ii) Hence, solve the equation $\frac{tanA-cotA}{tanA+cotA}+1=5sinAcosA$ for $0°<A<360°$ [5]

From (i), subs $\frac{tanA-cotA}{tanA+cotA}+1=2sin^2A$

$$2sin^2A=5\sin A\cos A$$ B1

$$\sin A(2\sin A-5\cos A)=0$$ M1

$\sin A=0$ *or* $2\sin A=5\cos A$

$A=180°(rejected)$ *or* $\tan A=\frac{5}{2}$


$basic\ angle, \alpha=tan^{-1}\frac{5}{2}$

$=68.2°$ M1

$A=68.2°,\ 248.2°$

Ans: $A=68.2°,\ 248.2°$ For 180°(rejected) A1

For 68.2°, 248.2° A1



Downloaded from SingaKids.com

12 A particle starts from rest, travels in a straight line so that t is the time in seconds after passing a fixed point O. Its velocity, $v\ m/s$, is given by $v = 6t - 2t^2$. The particle comes to instantaneous rest at A.

(i) Find the acceleration of the particle at A. [3]

$$\text{at } A,\ v = 0$$
$$\Rightarrow 2t(3 - t) = 0 \qquad \text{For } v = 0 \qquad \text{M1}$$
$$t = 0 \quad or \quad t = 3$$


$$a = \frac{dv}{dt}$$
$$a = 6 - 4t \qquad \text{M1}$$

$$\text{at } A,\ \ acceleration = 6 - 4(3) = -6\ m/s \qquad \text{A1}$$

(ii) Find the maximum velocity of the particle. [2]

$$For \max velocity,\ \frac{dv}{dt} = 0$$
$$6 - 4t = 0 \qquad \text{M1}$$
$$t = \frac{3}{2}$$

$$\therefore \max velocity = 6\left(\frac{3}{2}\right) - 2\left(\frac{3}{2}\right)^2 = 4\frac{1}{2}\ m/s \qquad \text{A1}$$



$v = 6t - 2t^2$.

Downloaded from SingaKids.com

12 (iii) Find the total distance travelled by the particle during the first 5 seconds. [5]


$$s = \int v\,dt$$
$$= \int 6t - 2t^2\,dt$$
$$= 3t^2 - \frac{2}{3}t^3 + c \qquad \text{M1}$$

$$\text{when } t = 0, s = 0 \quad \Rightarrow \quad c = 0$$
$$\therefore \; s = 3t^2 - \frac{2}{3}t^3 \qquad \text{B1}$$

$$\text{when } t = 3, \; s = 3(3)^2 - \frac{2}{3}(3)^3 = 9 \qquad \text{A1}$$

$$\text{when } t = 5, \; s = 3(5)^2 - \frac{2}{3}(5)^3 = -8\frac{1}{3} \qquad \text{A1}$$

$$\text{Total distance } = 9 + 9 + 8\frac{1}{3} = 26\frac{1}{3}\ m \qquad \text{A1}$$



End of Paper

Downloaded from SingaKids.com

ZHONGHUA SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2019
SECONDARY 4E/5N

Candidate's Name	Class	Register Number

ADDITIONAL MATHEMATICS 4047/02

PAPER 2

20 September 2019
2 hours 30 minutes

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer all the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the presentation, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 100.

For Examiner's Use
100

This question paper consists of 21 printed pages (including this cover page)

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \cdots + \binom{n}{r}a^{n-r}b^r + \cdots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!\,r!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$
$$\sec^2 A = 1 + \tan^2 A.$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$
$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$
$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$
$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$
$$\sin 2A = 2 \sin A \cos A$$
$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$
$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2}ab \sin C$$

Downloaded from SingaKids.com

Answer all the questions

1 The equation of a curve is $y=-x^2+(k-2)x+2k$, where k is a constant.

(i) Find the set of values of k for which the line $y=2kx+2k+1$ lies entirely above the curve. [3]

(ii) State the values of k for which the line $y=2kx+2k+1$ is a tangent to the curve. [1]

Downloaded from SingaKids.com

1 (iii) Explain why there is only one value of k for which y cannot be positive and state this value. [4]



Downloaded from SingaKids.com

2 (i) Find the first 3 terms in the expansion of $\left(2-\frac{x}{4}\right)^n$ in ascending powers of x, where n is a positive integer greater than 2. Give the terms in their simplest forms. [2]

(ii) In the expansion of $(4+x)^2\left(2-\frac{x}{4}\right)^n$, there is no term in x^2.

Find the value of n. [5]

Downloaded from SingaKids.com

3 (i) Express $\frac{4x^3+21x^2-4x-6}{(x^2+2)(2x-1)}$ as $a+\frac{bx^2+cx+d}{(x^2+2)(2x-1)}$

where a, b, c and d are integers. [2]

(ii) Differentiate $\ln(x^2+2)$ with respect to x. [2]

Downloaded from SingaKids.com

3 Using the results from (i) and (ii) and expressing $\frac{bx^2+cx+d}{(x^2+2)(2x-1)}$ as partial fractions, find

(iii) $\int \frac{4x^3+21x^2-4x-6}{(x^2+2)(2x-1)}\,dx$. [8]

Downloaded from SingaKids.com

4 (a) Variables x and y are related by the equation $px^2 + qy = x$, where p and q are constants. Y is plotted against X to obtain a straight line graph.

(i) If $X = x$, state Y in terms of x and/or y. [1]

(ii) Explain clearly how p and q can be obtained from the straight line graph drawn. [2]

(b) The number of snails in a colony is being studied. The number of snails in a colony after t weeks is given by $P = P_o e^{kt}$, where P_o and k are constants.
The table below gives some values of P and t.

t (weeks)	4	8	12	16	20
P	30	45	66	99	147

(i) On the grid on page 9, draw a suitable straight line graph to illustrate this data. [3]


Use your graph to estimate

(ii) the number of snails in the colony when the study began, [2]

(iii) the value of k. [2]

[Turn over

Downloaded from SingaKids.com



Downloaded from SingaKids.com

5 (a) Given that $\frac{\log_x y}{\log_y x} + \log_x y - 6 = 0$, express y in terms of x. [4]

(b) Solve the equation $9^{x+1} - 3^{x+1} = 2$. [3]

Downloaded from SingaKids.com

6 (i) On the same diagram, sketch the curve $y = x^{\frac{3}{2}}$ and $y = 8x^{-\frac{3}{2}}$. [2]

(ii) Find the coordinates of the point of intersection of the two curves. [3]

[Turn over

Downloaded from SingaKids.com

7 A circle, C_1, has the equation $x^2+y^2-4x+6y=12$.

(i) Find the coordinates of the centre and the radius of C_1. [3]

The equation of the tangent to C_1 at the point P is $4y-3x+43=0$.

Hence, using (i), find

(ii) the coordinates of the point P on C_1. [5]


Downloaded from SingaKids.com

7 A second circle, C_2, passes through the points (19, 8) and (−9, 12) and has a radius of 20 units.

(iii) Find the two possible centres of circle, C_2. [6]

Downloaded from SingaKids.com

8



The diagram shows two triangles, ABC and DCB. It is given that $BC = 12$ cm, $DC = 3$ cm, angle CAB = angle $DCB = 90°$ and angle $ABC = \theta°$, where θ varies.

A perpendicular is dropped from D to meet AB at E.

(i) Show that the perimeter of triangle DEB, P can be expressed as $\sqrt{153} + 9\sin\theta + 15\cos\theta$. [4]

Downloaded from SingaKids.com

8 (ii) Express P in the form $\sqrt{153}+R\cos(\theta-\alpha)$, where $R>0$ and α is an acute angle. [3]

(iii) Find the value of θ for which $P=27$. [3]

Downloaded from SingaKids.com

9 (a) Show that $\frac{1+\sin 4A-\cos 4A}{1+\sin 4A+\cos 4A}=\tan 2A$ [3]

(b) The temperature, T, in degrees Celsius, of a metal plate undergoing a chemical process after x seconds, is given by $T=-\frac{25}{9}x^2+50x+\frac{55}{3}$.

Determine when the metal plate starts cooling. [3]

Downloaded from SingaKids.com

10 A cubic curve has the stationary points at $(1, -7)$ and $\left(-\frac{7}{3}, \alpha\right)$, where α is a constant.

(i) Explain why $\frac{dy}{dx} = k(x-1)(3x+7)$, where k is a constant. [3]

Downloaded from SingaKids.com


10 (ii) Given that the curve passes through the point $(-1, 17)$, determine its equation. [6]

Downloaded from SingaKids.com

10 (iii) Determine the nature of the two stationary points. [3]

Downloaded from SingaKids.com

11



The diagram shows part of the curve $y = 5 - e^{1-2x}$ passing through the point P where $x = k$, where k is a constant. The tangent to the curve at P meets the line $y = 5$ at $Q(1, 5)$ and x-axis at $R\left(-\frac{3}{2}, 0\right)$. A vertical line is drawn at Q.

(i) Find the value of k. [4]

Downloaded from SingaKids.com

11 (ii) Find the total area of the shaded region. [5]

End of paper

Downloaded from SingaKids.com

ZHONGHUA SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2019
SECONDARY 4E/5N

Candidate's Name	Class	Register Number
MARKING SCHEME		

ADDITIONAL MATHEMATICS 4047/02

PAPER 2

20 September 2019
2 hours 30 minutes

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the presentation, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is **100**.

For Examiner's Use
100

This question paper consists of **20** printed pages (including this cover page)

Downloaded from SingaKids.com

Answer all the questions

1 The equation of a curve is $y=-x^2+(k-2)x+2k$, where k is a constant.

(i) Find the set of values of k for which the line $y=2kx+2k+1$ lies entirely above the curve. [3]

For $2kx+2k+1 > -x^2+(k-2)x+2k$

$x^2+(k+2)x+1 < 0$ M1


Discriminant < 0

$x^2+(k+2)x+1 < 0$

$(k+2)^2-4(1)(1) < 0$ M1

$(k+2-2)(k+2+2) < 0$

$k(k+4) < 0$



$-4 < k < 0$ A1

(ii) State the values of k for which the line $y=2kx+2k+1$ is a tangent to the curve. [1]

$k = -4$ or $k = 0$ A1

[Turn over

Downloaded from SingaKids.com

1 (iii) Explain why there is only one value of k for which y cannot be positive and state this value. [4]

Discriminant $= (k-2)^2 - 4(-1)(2k)$

$= k^2 - 4k + 4 + 8k$

$= k^2 + 4k + 4$

$= (k+2)^2$ **B1**

When $k = -2$, Discriminant $= 0$
$y = 0$ is a tangent to the curve.
Since coefficient of x^2 is negative, the maximum value of $y = 0$.
$\therefore y$ cannot be positive. **B1**

For any other values of k, $(k+2)^2 > 0$
Discriminant > 0
The curve intersects the x- axis at two points,
Maximum value of $y > 0$
y can be positive. **B1**

Ans: $k = -2$

Downloaded from SingaKids.com

2 (i) Find the first 3 terms in the expansion of $\left(2-\frac{x}{4}\right)^n$ in ascending powers of x,

where n is a positive integer greater than 2. Give the terms in their simplest forms. [2]

$$\left(2-\frac{x}{4}\right)^n = 2^n + \binom{n}{1}(2)^{n-1}\left(-\frac{x}{4}\right) + \binom{n}{2}(2)^{n-2}\left(-\frac{x}{4}\right)^2 + \ldots$$

$$= 2^n - \frac{n\times 2^n x}{8} + \frac{n(n-1)}{2}\times 2^{n-2}\times\frac{x^2}{16} + \ldots$$

$$= 2^n - \frac{n\times 2^n x}{8} + \frac{n(n-1)}{128}\times 2^n \times x^2 + \ldots$$

B1 for any 2 terms correct

B2 for all 3 terms correct

(ii) In the expansion of $(4+x)^2\left(2-\frac{x}{4}\right)^n$, there is no term in x^2.

Find the value of n. [5]

$$(16+8x+x^2)\left(2^n - \frac{n\times 2^n x}{8} + \frac{n(n-1)}{128}\times 2^n \times x^2 + \ldots\right)$$

expansion of $(4+x)^2$ B1

Term in x^2,

$$= 16\times\frac{n(n-1)}{128}\times 2^n \times x^2 + 8x\left(-\frac{n\times 2^n x}{8}\right) + x^2 \times 2^n \quad \text{M1-adding of 3 product}$$

coefficient of $x^2 = \frac{n(n-1)}{8}\times 2^n - n\times 2^n + 2^n = 0$

$$2^n\left(\frac{n(n-1)}{8} - n + 1\right) = 0 \quad \text{M1 (Factorisation)}$$

$2^n \neq 0,\ n(n-1) - 8n + 8 = 0$

$n^2 - 9n + 8 = 0$ B1

$(n-1)(n-8) = 0$

$n = 1$ (rejected) or $n = 8$ A1

Downloaded from SingaKids.com

3 (i) Express $\frac{4x^3+21x^2-4x-6}{(x^2+2)(2x-1)}$ as $a+\frac{bx^2+cx+d}{(x^2+2)(2x-1)}$

where a, b, c and d are integers. [2]

$$2x^3-x^2+4x-2 \;\overline{)\; 4x^3+21x^2-4x-6}$$

$$\begin{array}{r} 2 \\ \hline 4x^3+21x^2-4x-6 \\ -(4x^3-2x^2+8x-4) \\ \hline 23x^2-12x-2 \end{array}$$

$$\frac{4x^3+21x^2-4x-6}{(x^2+2)(2x-1)} = 2+\frac{23x^2-12x-2}{(x^2+2)(2x-1)} \quad \text{M1, A1}$$

(ii) Differentiate $\ln(x^2+2)$ with respect to x. [2]

$$\frac{d}{dx}\ln(x^2+2)$$

$$= \frac{2x}{x^2+2} \quad \frac{1}{x^2+2} \text{ seen B1, A1}$$

Downloaded from SingaKids.com

3 Using the results from (i) and (ii) and expressing $\frac{bx^2+cx+d}{(x^2+2)(2x-1)}$ as partial fractions,

find

(iii) $\int \frac{4x^3+21x^2-4x-6}{(x^2+2)(2x-1)}\,dx$. [8]

$$\int \frac{4x^3+21x^2-4x-6}{(x^2+2)(2x-1)}\,dx = \int 2+\frac{23x^2-12x-2}{(x^2+2)(2x-1)}\,dx$$

Let $\frac{23x^2-12x-2}{(x^2+2)(2x-1)} = \frac{A}{2x-1}+\frac{Bx+C}{(x^2+2)}$ M1

$23x^2-12x-2 = A(x^2+2)+(2x-1)(Bx+C)$ M1

Let $x=\frac{1}{2}$,

$\frac{23}{4}-\frac{12}{2}-2 = A\left(\frac{1}{4}+2\right)$

$-\frac{9}{4} = A\left(\frac{9}{4}\right)$

$A=-1$ A1

Comparing the coefficient of x^2 and constant,

$23 = A+2B$

$B=12$ A1

$-2 = 2A-C$

$C=0$ A1

$$\int \frac{4x^3+21x^2-4x-6}{(x^2+2)(2x-1)}\,dx = 2x+\int -\frac{1}{2x-1}+\frac{12x}{(x^2+2)}\,dx$$

$$= 2x-\frac{\ln(2x-1)}{2}+6\int \frac{2x}{(x^2+2)}\,dx$$

$$= 2x-\frac{\ln(2x-1)}{2}+6\ln(x^2+2)+c$$ A1

B1 B1

Downloaded from SingaKids.com

4 (a) Variables x and y are related by the equation $px^2 + qy = x$, where p and q are constants. Y is plotted against X to obtain a straight line graph.

(i) If $X = x$, state Y in terms of x and/or y. [1]

(ii) Explain clearly how p and q can be obtained from the straight line graph drawn. [2]

(i) $qy = -px^2 + x$

$$\frac{y}{x} = -\frac{p}{q}x + \frac{1}{q}$$

$$Y = \frac{y}{x} \qquad \text{A1}$$

(ii) Y-intercept $= \frac{1}{q}$ B1

gradient of graph $= -\frac{p}{q}$ B1

(b) The number of snails in a colony is being studied. The number of snails in a colony after t weeks is given by $P = P_o e^{kt}$, where P_o and k are constants

The table below gives some values of P and t.

t (weeks)	4	8	12	16	20
P	30	45	66	99	147

(i) On the grid on page 9, draw a suitable straight line graph to illustrate this data. [3]

$\ln P = \ln P_o + kt$ M1

Plot $\ln P$ against t. Straight line graph drawn B2

t	4	8	12	16	20
$\ln P$	3.40	3.81	4.19	4.60	4.99

Use your graph to estimate

(ii) the number of snails in the colony when the study began, [2]

From graph, $\ln P_o = 3$ M1

$P_o = e^3 = 20$ Number of snails = 20 A1


(iii) the value of k. [2]

Gradient $= \frac{4.60 - 3.40}{16 - 4}$ M1

k = gradient = 0. 1 A1

[Turn over

Downloaded from SingaKids.com



[Turn over

Downloaded from SingaKids.com

5 (a) Given that $\frac{\log_x y}{\log_y x} + \log_x y - 6 = 0$, express y in terms of x. [4]

$\frac{\log_x y}{\log_y x} + \log_x y - 6 = 0$

$\log_x y \div \frac{\log_x x}{\log_x y} + \log_x y - 6 = 0$ M1 change of base

$(\log_x y)^2 + \log_x y - 6 = 0$

$(\log_x y + 3)(\log_x y - 2) = 0$ Any method to solve quadratic eqn M1

$\log_x y = -3$ or $\log_x y = 2$

$y = x^{-3}$ or $y = x^2$ A1, A1

(b) Solve the equation $9^{x+1} - 3^{x+1} = 2$. [3]

Let $u = 3^x$

$9u^2 - 3u - 2 = 0$ either $3^{2x} \times 9 = 9u^2$ seen or $3^{x+1} = 3u$ seen B1

$(3u+1)(3u-2) = 0$

$u = -\frac{1}{3}$ or $u = \frac{2}{3}$ M1


$3^x = -\frac{1}{3}$ (rejected) or $3^x = \frac{2}{3}$

$x \lg 3 = \lg \frac{2}{3}$

$x = \frac{\lg \frac{2}{3}}{\lg 3} = -0.369$ (to 3 sf) A1

Downloaded from SingaKids.com

6 (i) On the same diagram, sketch the curve $y = x^{\frac{3}{2}}$ and $y = 8x^{-\frac{3}{2}}$. [2]



(ii) Find the coordinates of the point of intersection of the two curves. [3]

For points of intersection,

$$x^{\frac{3}{2}} = 8x^{-\frac{3}{2}}$$

$$x^{\frac{3}{2}} = \frac{8}{x^{\frac{3}{2}}}$$

$x^3 = 8$ M1

$x = 2$ A1

$y = 2^{\frac{3}{2}}$

Point of intersection = $(2, 2.83)$ A1

Downloaded from SingaKids.com

7 A circle, C_1, has the equation $x^2+y^2-4x+6y=12$.

(i) Find the coordinates of the centre and the radius of C_1. [3]

centre = (2, −3)

$r=\sqrt{2^2+(-3)^2-(-12)}$ M1

$=\sqrt{25}$

$= 5$ A1

The equation of the tangent to C_1 at the point P is $4y-3x+43=0$.

Hence, using (i), find

(ii) the coordinates of the point P on C_1. [5]

Gradient of normal at $P=-\frac{4}{3}$ M1

$C=(2,-3)$

Equation of the normal at P is $y+3=-\frac{4}{3}(x-2)$

$y=-\frac{4}{3}x-\frac{1}{3}$ ——(1) B1

For P, substitute (1) into $4y-3x+43=0$

$4\left(-\frac{4}{3}x-\frac{1}{3}\right)-3x+43=0$ M1

$-\frac{25}{3}x=-\frac{125}{3}$

$x=5$ A1

$y=-\frac{4}{3}\times 5-\frac{1}{3}=-7$

$P=(5,-7)$ A1

Downloaded from SingaKids.com

7 A second circle, C_2, passes through the points (19, 8) and (−9, 12) and has a radius of 20 units.

(iii) Find the two possible centres of circle, C_2. [6]

For perpendicular bisector of $(19, 8)$ *and* $(-9, 12)$

gradient of perpendicular bisector $= -1 \div \left(\frac{12-8}{-28}\right)$ M1

$= 7$

mid point $= (5, 10)$

Equation of perpendicular bisector is

$y - 10 = 7(x - 5)$

$y = 7x - 25$ B1

Let centre $= (x, 7x - 25)$

radius $= 20$

$\sqrt{(x+9)^2 + (7x-37)^2} = 20$ form an eqn in one unknown M1

$(x+9)^2 + (7x-37)^2 = 400$

$x^2 + 18x + 81 + 49x^2 - 2(7x)(37) + 1369 = 400$

$50x^2 - 500x + 1050 = 0$

$x^2 - 10x + 21 = 0$


$(x-3)(x-7) = 0$ any method seen to solve quadratic eqn M1

$x = 3$ or $x = 7$

Centre $= (3, -4)$ or centre $= (7, 24)$ A1, A1

Downloaded from SingaKids.com

8



The diagram shows two triangles, ABC and DCB. It is given that $BC = 12$ cm, $DC = 3$ cm, angle CAB = angle $DCB = 90°$ and angle $ABC = \theta°$, where θ varies.

A perpendicular is dropped from D to meet AB at E.

$\sin\theta = \frac{AC}{12}$ $\quad \cos\theta = \frac{AB}{12}$

$AC = 12\sin\theta$ or $AB = 12\cos\theta$ B1 for correct AC or AB

Let the perpendicular from C to DE intersects at F.

angle $FCB = \theta$ (alternate angles)

angle $CDF = 180^\circ - (90^\circ - \theta) - 90^\circ$

$= \theta$

$\cos\theta = \frac{DF}{3}$

$DF = 3\cos\theta$

$\sin\theta = \frac{CF}{3}$ [4]

$CF = 3\sin\theta$ B1 for correct DF or CF

Perimeter of triangle DFB, $P = DE + EB + DB$

$= 3\cos\theta + 12\sin\theta + 12\cos\theta - 3\sin\theta + \sqrt{3^2 + 12^2}$

(B1) or (B1) (B1)

$= \sqrt{153} + 9\sin\theta + 15\cos\theta$

Downloaded from SingaKids.com

8 (ii) Express P in the form $\sqrt{153}+R\cos(\theta-\alpha)$, where $R>0$ and α is an acute angle. [3]

$$P = \sqrt{153}+9\sin\theta+15\cos\theta$$
$$= \sqrt{153}+15\cos\theta+9\sin\theta$$
$$= \sqrt{153}+R\cos(\theta-\alpha)$$
$$R=\sqrt{15^2+9^2}=\sqrt{306} \quad \text{A1}$$
$$\alpha = \tan^{-1}\frac{9}{15} \quad \text{M1}$$
$$= 30.964^\circ$$
$$P = \sqrt{153}+\sqrt{306}\cos(\theta-31.0^\circ) \quad \text{A1}$$

(iii) Find the value of θ for which $P=27$. [3]

$$\sqrt{153}+\sqrt{306}\cos(\theta-30.964^\circ)=27$$
$$\sqrt{306}\cos(\theta-30.964^\circ) = \frac{27-\sqrt{153}}{\sqrt{306}} \quad \text{M1}$$
$$\theta-30.964^\circ = 33.240^\circ \quad \text{M1}$$
$$\theta = 64.2^\circ \quad \text{A1}$$

Downloaded from SingaKids.com

9 (a) Show that $\dfrac{1+\sin 4A-\cos 4A}{1+\sin 4A+\cos 4A}=\tan 2A$ [3]

$$\text{LHS} = \frac{1+\sin 4A-\cos 4A}{1+\sin 4A+\cos 4A}$$

$$= \frac{1+2\sin 2A\cos 2A-(1-2\sin^2 2A)}{1+2\sin 2A\cos 2A+(2\cos^2 2A-1)}$$ use of double angle identity B1

$$= \frac{2\sin 2A\cos 2A+2\sin^2 2A}{2\sin 2A\cos 2A+2\cos^2 2A}$$ B1

$$= \frac{2\sin 2A(\cos 2A+\sin 2A)}{2\cos 2A(\sin 2A+\cos 2A)}$$ M1 Factorisation

$$= \frac{\sin 2A}{\cos 2A}$$

$$= \tan 2A$$

(b) The temperature, T, in degrees Celsius, of a metal plate undergoing a chemical process after x seconds, is given by $T=-\dfrac{25}{9}x^2+50x+\dfrac{55}{3}$.

Determine when the metal plate starts cooling. [3]

$\dfrac{dT}{dx}=-\dfrac{50}{9}x+50$ M1 correct differentiation

For cooling to start,

$\dfrac{dT}{dx}<0$ B1

$-\dfrac{50}{9}x+50<0$

$x>9$ A1

[Turn over

Downloaded from SingaKids.com

10 A cubic curve has the stationary points at $(1, -7)$ and $\left(-\frac{7}{3}, \alpha\right)$, where α is a constant.

(i) Explain why $\frac{dy}{dx} = k(x-1)(3x+7)$, where k is a constant. [3]

Since stationary points occur at $x = 1$ or $x = -\frac{7}{3}$

$\frac{dy}{dx} = 0$ when $x = 1$ or $x = -\frac{7}{3}$ B1

$x - 1 = 0$ or $3x + 7 = 0$
$(x - 1)(3x + 7) = 0$ } either award B1

Since the curve is cubic, $\frac{dy}{dx}$ is quadratic,
Thus a cubic curve has at most 2 stationary points. B1

$\frac{dy}{dx} = k(x-1)(3x+7)$, k is a constant

Downloaded from SingaKids.com

10 (ii) Given that the curve passes through the point $(-1, 17)$, determine its equation. [6]

$y = \int k(3x^2 + 4x - 7)\,dx$ M1

$= k(x^3 + 2x^2 - 7x) + c$ B1 correct integration

substitute $x = 1$, $y = -7$

$-7 = -4k + c$ (1) B1

substitute $x = -1$, $y = 17$

$17 = 8k + c$ (2) B1

(2) − (1), $24 = 12k$

$k = 2$, $c = 1$ either ans for k or c A1

Equation is $y = 2x^3 + 4x^2 - 14x + 1$ A1

Downloaded from SingaKids.com

10 (iii) Determine the nature of the two stationary points. [3]

$$\frac{d^2y}{dx^2} = \frac{d}{dx}2(x-1)(3x+7)$$
$$= 2(x-1)(3)+2(3x+7)$$
$$= 12x+8 \qquad \text{M1}$$

when $x = 1$,


$\frac{d^2y}{dx^2} = 20 > 0$, $(1, -7)$ is a minimum point. B1

when $x = -\frac{7}{3}$

$\frac{d^2y}{dx^2} = -20 < 0$, $\left(-\frac{7}{3}, \alpha\right)$ is a maximum point B1

Downloaded from SingaKids.com

11



The diagram shows part of the curve $y = 5 - e^{1-2x}$ passing through the point P where $x = k$, where k is a constant. The tangent to the curve at P meets the line $y = 5$ at $Q(1, 5)$ and x-axis at $R\left(-\frac{3}{2}, 0\right)$. A vertical line is drawn at Q.

(i) Find the value of k. [4]

gradient of tangent at $P = \frac{5}{2.5}$

$= 2$ B1

$\frac{dy}{dx} = -(-2)e^{1-2x}$ B1

when $x = k$, $\frac{dy}{dx} = 2$,

$2 = 2e^{1-2k}$ M1

$1 = e^{1-2k}$

$1 - 2k = 0$

$k = \frac{1}{2}$ A1

[Turn over

Downloaded from SingaKids.com

11 (ii) Find the total area of the shaded region. [5]

Total area $= \frac{1}{2}\times 5\times\left(1-\left(-\frac{3}{2}\right)\right) - \int_0^1 5-e^{1-2x}\,dx$ M1

$= \frac{1}{2}\times 5\times\frac{5}{2} - \left[5x-\frac{e^{1-2x}}{-2}\right]_0^1$ A1, A1

$= \frac{25}{4} - \left(5+\frac{1}{2}e^{-1} - \left(\frac{1}{2}e\right)\right)$ M1 correct evaluation of definite integral

$= \frac{5}{4} - \frac{1}{2e} + \left(\frac{1}{2}e\right)$

$= 2.43$ (to 3 sf) A1

End of paper

Downloaded from SingaKids.com