1 Solve the equation $x+1+\sqrt{5-x}=0$. [3]

2 (a) (i) Given the equation $13\left(3^{2x}\right)=5^{3x}$, show that $\left(\frac{125}{9}\right)^{x}=13$. [1]

(ii) Hence find the value of x. [2]

www.testpapersfree.com

Downloaded from SingaKids.com

2 (b) Solve the equation $\log_x(2x-1)=\log_{x^2}(4-5x)$. [4]

3 **(i)** State the third term in the expansion of $(2+x)^n$ in terms of a and n. [2]

(ii) Write down the first three terms in the expansion of $(1-3x)^5$. [2]

(iii) In the expansion of $(1-3x)^5(2+ax)^6$, the coefficient of x is -384. Find the value of a. [4]

www.testpapersfree.com

Downloaded from SingaKids.com

4 Differentiate the following with respect to x.

(i) $3\tan^2\left(\frac{x}{2}\right)$ [2]

(ii) $e^{-x}x^2$ [3]

www.testpapersfree.com

Downloaded from SingaKids.com

5 Given that $y = \dfrac{e^{2x}}{\sqrt{1-4x}}$, show that $\dfrac{dy}{dx} = \dfrac{4e^{2x}(1-2x)}{(1-4x)\sqrt{1-4x}}$. [3]

www.testpapersfree.com

Downloaded from SingaKids.com

6 **(i)** The equation of a curve is given by $y = x + \sin^2 x$ for $0 \le x \le \pi$.

Write down the expression for $\frac{dy}{dx}$. [1]

(ii) Find stationary point of the curve. [3]

(iii) Determine the nature of the stationary point. [2]

7 Find the range of values of x such that the graph $y = \ln\left(\frac{x-2}{x-3}\right)^2$ is decreasing. [4]

www.testpapersfree.com

Downloaded from SingaKids.com

8 **(a)** The expression $f(x) = ax^3 + (a-3b)x^2 + 3bx + c$ is exactly divisible by $x^2 + 3x$. When $f(x)$ is divided by $x+2$, the remainder is 10. Find the values of a, b and c. Hence factorise $f(x)$ completely. [6]

(b) Given that $3x^3 - 2x^2 + x - 4 = A(x-1) + B(x-1)(x+1) + Cx(x^2-1) + D$ for all values of x, find the value of A, B, C and D. [4]

www.testpapersfree.com

Downloaded from SingaKids.com

9 The function f is defined, for $x \geq 0°$, by $f(x) = 3\cos\left(\frac{2x}{3}\right) - 1$.

(i) State the maximum and minimum values of $f(x)$. [2]

(ii) State the amplitude of $f(x)$. [1]

(iii) State the period of $f(x)$. [1]

(iv) Sketch the graph of $y = 3\cos\left(\frac{2x}{3}\right) - 1$ for $0° \leq x \leq 270°$. [2]

10 **(i)** Express the equations and in the form of

$$y = a(x+b)^2 + c.$$ [4]

(ii) Hence explain whether the curves will intersect or not. [3]

www.testpapersfree.com

Downloaded from SingaKids.com


11 **(a)** Find the range of values of k for which the line $y = kx + 3$ lies entirely below the curve $xy + 20 = 5y$ for all real values of x. [5]

(b) The quadratic equation $mx^2 + 5nx + 9m = 0$ has equal roots. Given that m and n are positive, find $\frac{m}{n}$. Determine also the range of values of $\frac{m}{n}$ for which the quadratic equation has real roots. [5]

www.testpapersfree.com

Downloaded from SingaKids.com

12



The picture above shows the path of a dolphin jumping out of the water to see clearly and to watch the surface of the oceans. The path is modelled by a quadratic function, $y = ax^2 + bx + c$ as shown in the diagram. The dimensions of the path is given in the table below.

How high can the dolphin jump out of the water?	450 cm
How far can the dolphin jump out of the water?	600 cm

(i) Find the coordinates of the highest point in the path. [1]

(ii) Find the equation, in the form $y = ax^2 + bx + c$, of the path of the dolphin, where a, b and c are real numbers. [4]

www.testpapersfree.com

Downloaded from SingaKids.com

13(i)	Factorise $x^4 - 1$ completely.
(ii)	Given that $\frac{x^4}{x^4-1} = a + \frac{b}{x^4-1}$, find the values of a and of b.
(iii)	Express $\frac{x^4}{x^4-1}$ in partial fractions.

www.testpapersfree.com

Downloaded from SingaKids.com

14 John deposits \$ P into a bank which pays a compound interest r % annually. The table shows the total amount \$ A in John's bank account over a period of 5 years.

n	1	2	3	4	5
A	8960	10035	11239	12588	14099

It is known that n and A are related by the equation $A = P\left(1+\frac{r}{100}\right)^n$.

(i) Express this equation in a form suitable for drawing a straight line graph. [2]

(ii) Draw a straight line graph for the given data. [2]

(iii) Use your graph to estimate the value of

(a) P, correct to the nearest thousand, and [2]

(b) r, correct to 2 significant figures. [2]

www.testpapersfree.com

Downloaded from SingaKids.com

www.testpapersfree.com

Downloaded from SingaKids.com

NAME: _____ ( ) CLASS: 4 ( )

ANGLICAN HIGH SCHOOL
SECONDARY FOUR
PRELIMINARY EXAMINATIONS 2021

S4

ADDITIONAL MATHEMATICS **4049 / 02**
Paper 2 **31st August 2021 Tuesday**
Candidates answer on the Question Paper. **2 hours 15 minutes**
No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your name, index number and class in the space at the top of this page.
Write in dark blue or black pen.
You may use a HB pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters and glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 100.

For Examiners' Use

Question	Marks	Question	Marks	Table of Penalties	
1		7			
2		8		Clarity/Logic	
3		9		Accuracy/Precision	
4		10		Units	
5		11		Total:	
6		12			
Parent's Name & Signature: Date:				90	

This document consists of **21** printed pages.

www.testpapersfree.com

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2\sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\text{Area of } \Delta = \frac{1}{2}ab\sin C$$

www.testpapersfree.com

Downloaded from SingaKids.com

1 Angles A and B are both obtuse and $\tan(A+B)=\frac{3}{4}$.

Without the use of calculators,

(i) state the range of angle $(A+B)$, [1]

(ii) find the value of $\sin(A+B)$, [2]

(iii) given that $\tan B=-\frac{12}{5}$, find the value of $\cos A$. [3]

www.testpapersfree.com

Downloaded from SingaKids.com

2 A slice of bread is toasted in an oven to a temperature of 154 °C. It is subsequently left to cool in such a way that its temperature, T °C, after removal from the oven, is

given by $T = 26 + Ae^{-kt}$, where A and k are constants, and t is the time after removal from oven in minutes.

(i) Find the value of A. [1]

(ii) What does the value "26" represent? [1]

When $t = 2$, the temperature of the bread is 116 °C.

(iii) Find the value of k. [2]

The toasted bread is recommended to be consumed at a temperature of 45 °C.

(iv) Determine, with clear working, the waiting time, in minutes, after removal from the oven before consuming the toasted bread. [2]

www.testpapersfree.com

Downloaded from SingaKids.com

3 Given that the function $f(x) = 1 + 2\sin^2 3x + 4\cos^2 3x$,

(i) Express $f(x)$ in the form $a + b\cos 6x$. [2]

(ii) Hence, find the exact value of $f(x) = 3.5$ for $0 \le x \le \frac{\pi}{2}$. [3]

www.testpapersfree.com

Downloaded from SingaKids.com

4 **(i)** Find the derivative of the functions $y = \ln x$ and $y = \ln 2x$ respectively. [1]

(ii) Explain why, in **(i)**, having the same derivative does not mean that the two functions are the same. [1]

www.testpapersfree.com

Downloaded from SingaKids.com

5 (i) Using $R\cos(\theta-\alpha)$, where $R>0$ and $0°\le\alpha\le90°$, solve the equation

$$5\cos\theta+8\sin\theta=6 \text{ for } 0°\le\theta\le180°.$$ [4]


(ii) State the largest value of $20-(5\cos x+8\sin x)^2$ and

find the corresponding value of x, such that $0°\le x\le180°$ [2]

www.testpapersfree.com

Downloaded from SingaKids.com

6 **Solutions to this question by accurate drawing will NOT be accepted.**



The figure shows a right-angled triangle ABC, where points A, B and C are $(k,0)$, $(2,8)$ and $(-2,-4)$ respectively. BC cuts the y-axis at N. M is a point on AC.

(i) Given that , find the value of k. [3]

www.testpapersfree.com

Downloaded from SingaKids.com

(ii) Find the coordinates of N and show that N is the midpoint of BC. [3]

www.testpapersfree.com

Downloaded from SingaKids.com

(iii) Given that the area of quadrilateral $ABNM$ is 25 units^2, find the coordinates of M. [6]

www.testpapersfree.com

Downloaded from SingaKids.com

7 **(i)** Show that $\frac{d}{dx}\ln(e^x \cos x) = 1 - \tan x$. [3]

(ii) Using part (i), find $\int \tan x \, dx$. [2]

www.testpapersfree.com

Downloaded from SingaKids.com

(iii) Hence show that $\int_0^{\frac{\pi}{3}} \tan x \, dx = \ln 2$. [2]

8 Point P lies on the curve $y = \frac{kx^2}{\sqrt{2-x}}$, where k is a constant. The x-coordinate of P is -2.

www.testpapersfree.com


Downloaded from SingaKids.com

(i) Given that the gradient of the normal at P is $\frac{1}{7}$, show that $k = 4$. [4]

www.testpapersfree.com

Downloaded from SingaKids.com

(i) The point P moves along the curve. Given that x is increasing at a rate of 0.9 units per second, find the rate of change of $(OP)^2$ at the point $x = 0.5$. [4]




www.testpapersfree.com

Downloaded from SingaKids.com

9 The diagram shows the glass test tube of radius r cm and height $(h+r)$ cm with a hemisphere of radius r cm as the base. The surface area of the test tube is 12π.

(i) Show that $h = \dfrac{6-r^2}{r}$. [2]



(ii) Show that the volume, V cm^3, of the test tube is $V = 6\pi r - \dfrac{1}{3}\pi r^3$. [2]

www.testpapersfree.com

Downloaded from SingaKids.com

(iii) Calculate the stationary value of V and determine if the volume is a maximum or minimum. [5]

www.testpapersfree.com

Downloaded from SingaKids.com

10 The equation of a circle, C_1, is .

(i) Find the coordinates of the centre of C_1 and its radius. [3]

(ii) Find the equation of the circle, C_2, which is a reflection of C_1 in the line $y = 1$. [2]

(iii) Given that the line intersects the circle, C_1, at two points, R and S, find the equation of the perpendicular bisector of the line RS. [4]

www.testpapersfree.com

Downloaded from SingaKids.com

11 A particle moves in a straight line, so that, t seconds after leaving a fixed point O its velocity, v is given by $v = \frac{36}{(t-3)^2} - 9$. Find

(i) the initial acceleration of the particle. [2]

(ii) the values of t when the particle is at instantaneous rest. [2]

www.testpapersfree.com


Downloaded from SingaKids.com

(iii) the total distance travelled by the particle in the first 2 seconds. [6]

www.testpapersfree.com

Downloaded from SingaKids.com

12



The diagram shows part of the curve $y = \sqrt{3x+4}$. The line AB is tangent to the curve at A. The points B and C lie on the x-axis, and C is a point on the curve. The gradient of the tangent of the curve at A is $\frac{3}{2}$. Find

(i) the coordinates of A and equation of the line AB. [5]

www.testpapersfree.com

Downloaded from SingaKids.com

(ii) the coordinates of B and of C. [2]

(iii) the area of the shaded region ABC. [3]

www.testpapersfree.com

Downloaded from SingaKids.com

www.testpapersfree.com

Downloaded from SingaKids.com

1 Solve the equation $x+1+\sqrt{5-x}=0$. [3]

$x+1+\sqrt{5-x}=0$

$x+1=-\sqrt{5-x}$

$(x+1)^2=5-x$

$x^2+2x+1-5+x=0$

$x^2+3x-4=0$

$(x+4)(x-1)=0$

$x=-4$ or $x=1$ (rejected)

2 (a) (i) Given the equation $13\left(3^{2x}\right)=5^{3x}$, show that $\left(\frac{125}{9}\right)^x=13$. [1]

$13\left(3^{2x}\right)=5^{3x}$

$13=\frac{\left(5^3\right)^x}{\left(3^2\right)^x}$

$\left(\frac{125}{9}\right)^x=13$

(ii) Hence find the value of x. [2]

$\left(\frac{125}{9}\right)^x=13$

$\lg\left(\frac{125}{9}\right)^x=\lg 13$

$x=\lg 13\div\lg\left(\frac{125}{9}\right)$

$x\approx 0.975$ (3 s.f.)

www.testpapersfree.com

Downloaded from SingaKids.com

2 (b) Solve the equation $\log_x(2x-1)=\log_{x^2}(4-5x)$. [4]

$\log_x(2x-1)=\log_{x^2}(4-5x)$

$\log_x(2x-1)=\dfrac{\log_x(4-5x)}{\log_x x^2}$

$\log_x(2x-1)=\dfrac{\log_x(4-5x)}{2}$

$2\log_x(2x-1)-\log_x(4-5x)=0$

$\log_x\dfrac{(2x-1)^2}{4-5x}=0$

$\dfrac{(2x-1)^2}{4-5x}=1$

$4x^2-4x+1=4-5x$

$4x^2+x-3=0$

$(4x-3)(x+1)=0$

$x=\dfrac{3}{4}$ or $x=-1$ (N.A.)

3 **(i)** State the third term in the expansion of $(2+x)^n$ in terms of a and n. [2]

(ii) Write down the first three terms in the expansion of $(1-3x)^5$. [2]

(iii) In the expansion of $(1-3x)^5(2+ax)^6$, the coefficient of x is -384. Find the value of a. [4]

3(i)

$T_3=\binom{n}{2}2^{n-2}(x)^2$

$=\dfrac{n(n-1)}{2}\dfrac{2^n}{2^2}x^2$

$=n(n-1)2^{n-3}x^2$ *or* $\dfrac{n(n-1)2^{n-2}x^2}{2}$

(ii)

$(1-3x)^5$

$=1+\binom{5}{1}(-3x)+\binom{5}{2}(-3x)^2+\ldots$

$=1-15x+90x^2+\ldots$

www.testpapersfree.com

Downloaded from SingaKids.com

(iii)	$(1-3x)^5(2+ax)^6$ $=\left(1-15x+90x^2+\ldots\right)\left[2^6+\binom{6}{1}2^5(ax)+\ldots\right]$ $=\left(1-15x+90x^2+\ldots\right)\left(64+192ax+\ldots\right)$ Coefficient of $x=(1)(192a)+(-15)(64)=-384$ $192a-960=-384$ $192a=576$ $a=3$

4 Differentiate the following with respect to x.

(i) $3\tan^2\left(\frac{x}{2}\right)$ [2]

(ii) $e^{-x}x^2$ [3]

4(i)	$\frac{d}{dx}\left[3\tan^2\left(\frac{x}{2}\right)\right]=3\left(2\tan\frac{x}{2}\right)\left(\sec^2\frac{x}{2}\right)\left(\frac{1}{2}\right)$ $=3\tan\frac{x}{2}\sec^2\frac{x}{2}$
(ii)	$\frac{d}{dx}\left[e^{-x}x^2\right]=e^{-x}(2x)+\left(-e^{-x}\right)x^2$ $=e^{-x}\left(2x-x^2\right)$ OR $xe^{-x}(2-x)$

www.testpapersfree.com
Downloaded from SingaKids.com

5 Given that $y = \dfrac{e^{2x}}{\sqrt{1-4x}}$, show that $\dfrac{dy}{dx} = \dfrac{4e^{2x}(1-2x)}{(1-4x)\sqrt{1-4x}}$. [3]

5	$\dfrac{dy}{dx} = \dfrac{e^{2x}}{\sqrt{1-4x}}$ $= \dfrac{\sqrt{1-4x}\dfrac{d}{dx}e^{2x} - e^{2x}\dfrac{d}{dx}\sqrt{1-4x}}{\left(\sqrt{1-4x}\right)^2}$ $= \dfrac{2\sqrt{1-4x}\,e^{2x} - e^{2x}\left(\dfrac{1}{2}\right)(1-4x)^{-\frac{1}{2}}(-4)}{1-4x}$ $= \dfrac{e^{2x}\left(2\sqrt{1-4x} + 2(1-4x)^{-\frac{1}{2}}\right)}{1-4x}$ $= \dfrac{2e^{2x}(1-4x+1)}{(1-4x)\sqrt{1-4x}}$ $= \dfrac{4e^{2x}(1-2x)}{(1-4x)\sqrt{1-4x}}$ (shown)

6 **(i)** The equation of a curve is given by $y = x + \sin^2 x$ for $0 \le x \le \pi$.

Write down the expression for $\dfrac{dy}{dx}$. [1]

www.testpapersfree.com

Downloaded from SingaKids.com

(ii) Find stationary point of the curve. [3]

(iii) Determine the nature of the stationary point. [2]

6

(i) $y = x + \sin^2 x$

$\frac{dy}{dx} = 1 + 2\sin x \cos x \ (or = 1 + \sin 2x)$

(ii) At stationary point, $\frac{dy}{dx} = 0$

$1 + \sin 2x = 0$

$\sin 2x = -1$

$2x = \frac{3\pi}{2} or\ 2\pi + \frac{3\pi}{2} rej, since \quad 0 \le 2x \le 2\pi$

$x = \frac{3\pi}{4} \ or \ 2.36$

$y = \frac{3\pi}{4} \qquad + \sin^2\left(\frac{3\pi}{4}\right)$

$= \frac{3\pi}{4} + \frac{1}{2} \ or \ \frac{3\pi + 2}{4} \ or \ 2.86$

Therefore the stationary point is $(2.36,\ 2.86)$

$or \left(\frac{3\pi}{4}, \frac{3\pi + 2}{4}\right) or \left(\frac{3\pi}{4}, 0.5 + \frac{3\pi}{4}\right)$

(iii)

x	$\frac{3\pi}{4}-$	$\frac{3\pi}{4}$	$\frac{3\pi}{4}+$
$\frac{dy}{dx}$	+ve	0	+ve

Therefore the stationary point is a point of inflexion

7 Find the range of values of x such that the graph $y = \ln\left(\frac{x-2}{x-3}\right)^2$ is decreasing. [4]

7

www.testpapersfree.com

Downloaded from SingaKids.com

$$y = \ln\left(\frac{x-2}{x-3}\right)^2$$


$$= 2\left(\ln(x-2) - \ln(x-3)\right)$$

$$\frac{dy}{dx} = 2\left(\frac{1}{x-2} - \frac{1}{x-3}\right)$$

$$= 2\left(\frac{x-3-x+2}{(x-2)(x-3)}\right)$$

$$= \frac{-2}{(x-2)(x-3)}$$

$$(x-2)(x-3) > 0$$



$x < 2$ or $x > 3$

8 **(a)** The expression $f(x) = ax^3 + (a-3b)x^2 + 3bx + c$ is exactly divisible by $x^2 + 3x$. When $f(x)$ is divided by $x+2$, the remainder is 10. Find the values of a, b and c. Hence factorise $f(x)$ completely. [6]

(b) Given that $3x^3 - 2x^2 + x - 4 = A(x-1) + B(x-1)(x+1) + Cx(x^2-1) + D$ for all values of x, find the value of A, B, C and D. [4]

8(a)

$f(0) = c = 0 \quad \Rightarrow c = 0$

$f(-3) = 0$

$-27a + 9(a-3b) - 9b = 0$

$-18a - 36b = 0$

$a = -2b$

www.testpapersfree.com

Downloaded from SingaKids.com

	$f(-2)=10$ $-8a+4(a-3b)-6b=10$ $-4a-18b=10$ $8b-18b=10$ $b=-1$ $a=2$ $f(x)=2x^3+5x^2-3x$ $=x(2x^2+5x-3)$ $=x(x+3)(2x-1)$
(b)	$3x^3-2x^2+x-4=A(x-1)+B(x-1)(x+1)+Cx(x^2-1)+D$ When $x=1$, $D=-2$ When $x=-1$, $A=4$ When $x=0$, $B=-2$ When $x=2$, $C=3$

www.testpapersfree.com

Downloaded from SingaKids.com


9 The function f is defined, for $x \geq 0°$, by $f(x) = 3\cos\left(\frac{2x}{3}\right) - 1$.

(i) State the maximum and minimum values of $f(x)$. [2]

(ii) State the amplitude of $f(x)$. [1]

(iii) State the period of $f(x)$. [1]

(iv) Sketch the graph of $y = 3\cos\left(\frac{2x}{3}\right) - 1$ for $0° \leq x \leq 270°$. [2]

9(i)	Max value of $f(x) = 2$ Min value of $f(x) = -4$
(ii)	Amplitude = 3
(iii)	Period = $360° \div \frac{2}{3} = 540°$
(iv)	

10 **(i)** Express the equations $y = 2x^2 - 3x + 9$ and $y = -2x^2 - 3x + \frac{27}{4}$ in the form of

$$y = a(x+b)^2 + c.$$ [4]

(ii) Hence explain whether the curves will intersect or not. [3]


www.testpapersfree.com

Downloaded from SingaKids.com

10(i)	$y = 2x^2 - 3x + 9$ $y = 2\left(x^2 - \frac{3}{2}x\right) + 9$ $y = 2\left[\left(x - \frac{3}{4}\right)^2 - \left(\frac{3}{4}\right)^2\right] + 9$ $y = 2\left(x - \frac{3}{4}\right)^2 - \frac{9}{8} + 9$ $y = 2\left(x - \frac{3}{4}\right)^2 + \frac{63}{8}$ $y = -2x^2 - 3x + \frac{27}{4}$ $y = -2\left(x^2 + \frac{3}{2}x\right) + \frac{27}{4}$ $y = -2\left[\left(x + \frac{3}{4}\right)^2 - \left(\frac{3}{4}\right)^2\right] + \frac{27}{4}$ $y = 2\left(x + \frac{3}{4}\right)^2 + \frac{9}{8} + \frac{27}{4}$ $y = -2\left(x + \frac{3}{4}\right)^2 + \frac{63}{8}$
(ii)	The minimum point of $y = 2x^2 - 3x + 9$ is $\left(\frac{3}{4}, \frac{63}{8}\right)$. The maximum point of $y = -2x^2 - 3x + \frac{27}{4}$ is $\left(-\frac{3}{4}, \frac{63}{8}\right)$. Both of the y coordinates are the same but their x coordinates are different.

www.testpapersfree.com

Downloaded from SingaKids.com



From the graphs, the two graphs will not intersect.

11 **(a)** Find the range of values of k for which the line $y = kx + 3$ lies entirely below the curve $xy + 20 = 5y$ for all real values of x. [5]

(b) The quadratic equation $mx^2 + 5nx + 9m = 0$ has equal roots. Given that m and n are positive, find $\frac{m}{n}$. Determine also the range of values of $\frac{m}{n}$ for which the quadratic equation has real roots. [5]

(a)

$y = kx + 3 - - - - - - - (1)$

$xy + 20 = 5y - - - - - - (2)$

Substitute (1) into (2)

$x(kx + 3) + 20 = 5(kx + 3)$

$kx^2 + 3x + 20 = 5kx + 15$

$kx^2 + 3x - 5kx + 5 = 0$


Since line and curve does not intersect,

$(3 - 5k)^2 - 4(k)(5) < 0$

$9 - 30k + 25k^2 - 20k < 0$

$25k^2 - 50k + 9 < 0$


$(5k - 1)(5k - 9) < 0$



$\frac{1}{5} < k < \frac{9}{5}$ *or* $0.2 < k < 1.8$

www.testpapersfree.com

Downloaded from SingaKids.com


	Alternatively, $y = kx + 3$ $xy + 20 = 5y$ $20 = 5y - xy$ $y(5 - x) = 20$ $y = \frac{20}{5 - x}$ $kx + 3 < \frac{20}{5 - x}$ $(kx + 3)(5 - x) < 20$ $5kx - kx^2 + 15 - 3x < 20$ $-kx^2 + (5k - 3)x - 5 < 0$ $kx^2 + (3 - 5k)x + 5 > 0$ for all real values x. $b^2 - 4ac < 0$ $(3 - 5k)^2 - 4k(5) < 0$ $9 - 30k + 25k^2 - 20k < 0$ $25k^2 - 50k + 9 < 0$ $(5k - 1)(5k - 9) < 0$  $\frac{1}{5} < k < \frac{9}{5}$ *or* $0.2 < k < 1.8$
11(b)	For equal roots, $b^2 - 4ac = 0$ $(5n)^2 - 4m(9m) = 0$ $25n^2 = 36m^2$ $\frac{m^2}{n^2} = \frac{25}{36}$ $\frac{m}{n} = \frac{5}{6}$ or $\frac{m}{n} = -\frac{5}{6}$(rejected)

www.testpapersfree.com

Downloaded from SingaKids.com

	For real roots, $b^2-4ac\geq 0$ $(5n)^2-4m(9m)\geq 0$ $25n^2-36m^2\geq 0$ $36m^2-25n^2\leq 0$ $(6m-5n)(6m+5n)\leq 0$ $-5n\leq 6m\leq 5n$ $-\frac{5}{6}\leq\frac{m}{n}\leq\frac{5}{6}$ Since $\frac{m}{n}\geq 0$, $0\leq\frac{m}{n}\leq\frac{5}{6}$.

12



The picture above shows the path of a dolphin jumping out of the water to see clearly and to watch the surface of the oceans. The path is modelled by a quadratic function, $y=ax^2+bx+c$ as shown in the diagram. The dimensions of the path is given in the table below.

How high can the dolphin jump out of the water?	450 cm
How far can the dolphin jump out of the water?	600 cm

(i) Find the coordinates of the highest point in the path. [1]

(ii) Find the equation, in the form $y=ax^2+bx+c$, of the path of the dolphin, where a, b and c are real numbers. [4]

www.testpapersfree.com

Downloaded from SingaKids.com

12(i)	$x = \frac{600}{2}$ $= 300$ Coordinates of the turning point = $(300,\ 450)$
(ii)	$y = a(x-300)^2 + 450$ When $x = 0,\ y = 0$ $0 = a(-300)^2 + 450$ $a = -\frac{1}{200}$ or -0.005 Equation of the path is $y = -\frac{1}{200}(x-300)^2 + 450$ $= -\frac{1}{200}(x^2 - 600x + 90000) + 450$ $= -\frac{1}{200}x^2 + 3x - 450 + 450$ $= -\frac{1}{200}x^2 + 3x.$
12(ii)	Alternatively, When $x = 0,\ y = 0,\ y = ax^2 + bx + c$ becomes $0 = a(0)^2 + b(0) + c$ $c = 0$ $y = ax^2 + bx$ When $x = 300,\ y = 450,$ $450 = a(300)^2 + b(300)$ $b = \frac{3}{2} - 300a$ ----------(1) When $y = 0,\ x = 600,$ $0 = a(600)^2 + b(600)$ $b = -600a$ ----------(2) Equating (1) and (2), $-600a = \frac{3}{2} - 300a$ $a = -\frac{1}{200}$ From (2), $b = -600a$ $b = 3$ Equation of the path is $y = -\frac{1}{200}x^2 + 3x.$

www.testpapersfree.com

Downloaded from SingaKids.com

13(i) Factorise x^4-1 completely.

$$\begin{aligned} x^4-1 &= (x^2-1)(x^2+1) \\ &= (x-1)(x+1)(x^2+1) \end{aligned}$$

(ii) Given that $\dfrac{x^4}{x^4-1} = a + \dfrac{b}{x^4-1}$, find the values of a and of b.

$$\begin{aligned} \frac{x^4}{x^4-1} &= \frac{(x^4-1)+1}{x^4-1} \\ &= 1+\frac{1}{x^4-1} \end{aligned}$$

$a=1,\ b=1$

(iii) Express $\dfrac{x^4}{x^4-1}$ in partial fractions.

$$\frac{x^4}{x^4-1} = 1+\frac{1}{x^4-1}$$

$$\begin{aligned} \frac{1}{x^4-1} &= \frac{1}{(x-1)(x+1)(x^2+1)} \\ &= \frac{A}{x-1}+\frac{B}{x+1}+\frac{Cx+D}{x^2+1} \end{aligned}$$

$$1 = A(x+1)(x^2+1)+B(x-1)(x^2+1)+(Cx+D)(x-1)(x+1)$$

When $x=1$,
$4A=1$

$$A=\frac{1}{4}$$

When $x=-1$
$-4B=1$

$$B=-\frac{1}{4}$$

When $x=0$
$A-B-D=1$

$$D=\frac{1}{4}-\left(-\frac{1}{4}\right)-1=-\frac{1}{2}$$

By comparing coefficient of x^3

www.testpapersfree.com

Downloaded from SingaKids.com

$0 = A + B + C$

$C = -\frac{1}{4} - \left(-\frac{1}{4}\right) = 0$

$\frac{x^4}{x^4 - 1} = 1 + \frac{1}{4(x-1)} - \frac{1}{4(x+1)} - \frac{1}{2(x^2+1)}$

14 John deposits \$ P into a bank which pays a compound interest r % annually. The table shows the total amount \$ A in John's bank account over a period of 5 years.

n	1	2	3	4	5
A	8960	10035	11239	12588	14099

It is known that n and A are related by the equation $A = P\left(1 + \frac{r}{100}\right)^n$.

(i) Express this equation in a form suitable for drawing a straight line graph. [2]

(ii) Draw a straight line graph for the given data. [2]

(iii) Use your graph to estimate the value of
(a) P, correct to the nearest thousand, and [2]
(b) r, correct to 2 significant figures. [2]

14(i)

$A = P\left(1 + \frac{r}{100}\right)^n$

$\lg A = \lg P\left(1 + \frac{r}{100}\right)^n$


$\lg A = \lg P + \lg\left(1 + \frac{r}{100}\right)^n$

$\lg A = n\lg\left(1 + \frac{r}{100}\right) + \lg P$

(ii)

n	1	2	3	4	5
A	8960	10035	11239	12588	14099
$\lg A$	3.95	4.00	4.05	4.10	4.15

www.testpapersfree.com
Downloaded from SingaKids.com



(iii) (a) From the graph, the $\lg A$ – intercept is 3.9

Therefore, $\lg P = 3.9$

$P = 10^{3.9}$

$P = 7943.28$

$P \approx 8000$ (nearest thousand)

(b) Gradient of graph is $\lg\left(1+\frac{r}{100}\right) = \frac{4.15-9.95}{5-1}$

$\lg\left(1+\frac{r}{100}\right) = 0.05$

$1+\frac{r}{100} = 10^{0.05}$

$\frac{r}{100} = 1.122-1$

$r \approx 12$ (2 s.f.)

www.testpapersfree.com

Downloaded from SingaKids.com

NAME: _____ ( ) CLASS: 4 ( )

ANGLICAN HIGH SCHOOL
SECONDARY FOUR
PRELIMINARY EXAMINATIONS 2021

S4

ADDITIONAL MATHEMATICS

4049 / 02

Paper 2

31st August 2021 Tuesday

Candidates answer on the Question Paper.

2 hours 15 minutes

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your name, index number and class in the space at the top of this page.
Write in dark blue or black pen.
You may use a HB pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters and glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 100.

For Examiners' Use

Question	Marks	Question	Marks	Table of Penalties	
1		7			
2		8		Clarity/Logic	
3		9		Accuracy/Precision	
4		10		Units	
5		11		Total:	
6		12			
Parent's Name & Signature: Date:				90	

This document consists of **21** printed pages.

www.testpapersfree.com

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2\sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\text{Area of } \Delta = \frac{1}{2}ab\sin C$$

www.testpapersfree.com

Downloaded from SingaKids.com


1 Angles A and B are both obtuse and $\tan(A+B)=\frac{3}{4}$.

Without the use of calculators,


(i) state the range of angle $(A+B)$, [1]

(i)	$180° < (A+B) < 270°$

(ii) find the value of $\sin(A+B)$, [2]

(ii)	 $\sin(A+B)=-\frac{3}{5}$	

(iii) given that $\tan B=-\frac{12}{5}$, find the value of $\cos A$. [3]

(iii)	$\tan(A+B)=\frac{3}{4}$ $\frac{\tan A+\tan B}{1-\tan A\tan B}=\frac{3}{4}$ $\frac{\tan A-\frac{12}{5}}{1+\frac{12}{5}\tan A}=\frac{3}{4}$ $-\frac{16}{5}\tan A=\frac{63}{5}$ $\tan A=-\frac{63}{16}$ $\cos A=-\frac{16}{65}$ 	

www.testpapersfree.com

Downloaded from SingaKids.com

2 A slice of bread is toasted in an oven to a temperature of 154 °C. It is subsequently left to cool in such a way that its temperature, T °C, after removal from the oven, is given by $T = 26 + Ae^{-kt}$, where A and k are constants, and t is the time after removal from oven in minutes.

(i) Find the value of A. [1]

(i)	At $t = 0$, $T = 154$. $154 = 26 + Ae^{0}$ $A = 128$	 B1

(ii) What does the value "26" represent? [1]

(ii)	"26" represents the room temperature.	B1

When $t = 2$, the temperature of the bread is 116 °C.

(iii) Find the value of k. [2]

(iii)	At $t = 2$, $T = 116$. $116 = 26 + 128e^{-2k}$ $e^{-2k} = \frac{116 - 26}{128}$ $-2k = \ln\left(\frac{90}{128}\right)$ $k \approx 0.17611$ $k \approx 0.176$ (3 s.f.)	

The toasted bread is recommended to be consumed at a temperature of 45 °C.

(iv) Determine, with clear working, the waiting time, in minutes, after removal from the oven before consuming the toasted bread. [2]

(iv)	When $T = 45$, $45 = 26 + 128e^{-0.17611t}$ $e^{-0.17611t} = \frac{45 - 26}{128}$ $-0.17611t = \ln\left(\frac{19}{128}\right)$ $t \approx 10.8$ (1 d.p.) Waiting time is 10.8 min after removal from oven.	

www.testpapersfree.com

Downloaded from SingaKids.com

3 Given that the function $f(x) = 1 + 2\sin^2 3x + 4\cos^2 3x$,

(i) Express $f(x)$ in the form $a + b\cos 6x$. [2]

(i)	$f(x) = 1 + 2\sin^2 3x + 4\cos^2 3x$ $= 1 + 2(1 - \cos^2 3x) + 4\cos^2 3x$ $= 3 + 2\cos^2 3x$ $= 3 + \cos 6x + 1$ $= 4 + \cos 6x$	

(ii) Hence, find the exact value of $f(x) = 3.5$ for $0 \le x \le \frac{\pi}{2}$. [3]

(ii)	$f(x) = 3.5$ $4 + \cos 6x = 3.5$ $\cos 6x = -\frac{1}{2}$ basic angle $= \cos^{-1}\left(\frac{1}{2}\right)$ $= \frac{\pi}{3}$ $0 \le x \le \frac{\pi}{2}$ $0 \le 6x \le 3\pi$ $6x = \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{8\pi}{3}$ $x = \frac{\pi}{9}, \frac{2\pi}{9}, \frac{4\pi}{9}$	

www.testpapersfree.com

Downloaded from SingaKids.com

4 **(i)** Find the derivative of the functions $y = \ln x$ and $y = \ln 2x$ respectively. [1]

(i)	$\frac{dy}{dx} = \frac{d}{dx}\ln x$ $= \frac{1}{x}$ $\frac{dy}{dx} = \frac{d}{dx}\ln 2x$ $= \frac{1}{2x} \times 2$ $= \frac{1}{x}$	B1 B1

(ii) Explain why, in **(i)**, having the same derivative does not mean that the two functions are the same. [1]

(ii)	Since $y = \ln x$ $y = \log_e x$ $x = e^y$ And $y = \ln 2x$ $y = \log_e 2x$ *or* $y = \ln 2 + \ln x$ $2x = e^y$ $\ln x = y - \ln 2$ $x = \frac{1}{2}e^y$ $x = e^{y-\ln 2}$ $x = \frac{e^y}{e^{\ln 2}} = \frac{e^y}{2}$ Hence, they are different functions.	

5 **(i)** Using $R\cos(\theta - \alpha)$, where $R > 0$ and $0° \le \alpha \le 90°$, solve the equation

$$5\cos\theta + 8\sin\theta = 6 \text{ for } 0° \le \theta \le 180°.$$ [4]


(i)	$5\cos\theta + 8\sin\theta = 6$ $R\cos(\theta - \alpha) = 6$ $R = \sqrt{5^2 + 8^2}$ $\tan\alpha = \frac{8}{5}$ $= \sqrt{89}$ $\alpha = 57.9946°$	

www.testpapersfree.com

Downloaded from SingaKids.com

$\sqrt{89}\cos(\theta - 57.9946°) = 6$

$\cos(\theta - 57.9946°) = \frac{6}{\sqrt{89}}$



$\text{basic angle} = \cos^{-1}\left(\frac{6}{\sqrt{89}}\right)$

$= 50.5059°$

$\theta - 57.9946° = -50.5059°,\ 50.5059°$

$\theta = 7.4887°, 108.5005°$

$\theta = 7.5°, 108.5°$

(ii) State the largest value of $20 - (5\cos x + 8\sin x)^2$ and find the corresponding value of x, such that $0° \le x \le 180°$ [2]

(ii)

$20 - (5\cos x + 8\sin x)^2 = 20 - \left(\sqrt{89}\cos(x - 57.9946°)\right)^2$

$= 20 - 89\cos^2(x - 57.9946°)$

Largest value = 20

$\cos(x - 57.9946°) = 0$

$x - 57.9946° = 90°$


$x = 147.9946°$

$= 148.0°$

www.testpapersfree.com

Downloaded from SingaKids.com

6 **Solutions to this question by accurate drawing will NOT be accepted.**



The figure shows a right-angled triangle ABC, where points A, B and C are $(k,0)$, $(2,8)$ and $(-2,-4)$ respectively. BC cuts the y-axis at N. M is a point on AC.

(i) Given that $k > 0$, find the value of k. [3]

(i)	Gradient of $AB \times$ gradient of $AC = -1$ $\left(\frac{0-8}{k-2}\right)\times\left(\frac{0-(-4)}{k-(-2)}\right)=-1$ $(-8)(4)=-(k-2)(k+2)$ $-32=-k^2+4$ $k^2=36$ $k=\pm 6$ Since $k>0$, $k=6$	

www.testpapersfree.com

Downloaded from SingaKids.com

(ii) Find the coordinates of N and show that N is the midpoint of BC. [3]

(ii)	Let the coordinates of N be $(0,n)$. Gradient of BN = gradient of BC $\left(\frac{8-n}{2-0}\right)=\left(\frac{8-(-4)}{2-(-2)}\right)$ $\left(\frac{8-n}{2}\right)=3$ $8-n=6$ $n=2$ Coordinates of N is $(0,2)$. Midpoint of BC is $\left(\frac{2-2}{2},\frac{8-4}{2}\right)=(0,2)$. Hence, N is the midpoint of BC.	

www.testpapersfree.com

Downloaded from SingaKids.com

(iii) Given that the area of quadrilateral $ABNM$ is 25 units^2, find the coordinates of M. [6]

(iii)

Gradient of $AC = \dfrac{0-(-4)}{6-(-2)} = \dfrac{1}{2}$

Equation of AC is

$y - 0 = \dfrac{1}{2}(x-6)$

$y = \dfrac{1}{2}x - 3$

Let the coordinates of M be $\left(a, \dfrac{1}{2}a-3\right)$

Area of quadrilateral $ABNM = 25$ units^2

$$\frac{1}{2}\begin{vmatrix} 6 & 2 & 0 & a & 6 \\ 0 & 8 & 2 & \frac{1}{2}a-3 & 0 \end{vmatrix} = 25$$

$$\frac{1}{2}\left[(48+4)-\left(2a+6\left(\frac{1}{2}a-3\right)\right)\right] = 25$$

$$52 - 2a - 3a + 18 = 50$$

$$5a = 20$$

$$a = 4$$

Coordinates of M is $(4,-1)$

www.testpapersfree.com

Downloaded from SingaKids.com

7 **(i)** Show that $\frac{d}{dx}\ln(e^x\cos x)=1-\tan x$. [3]

(i)	$\frac{d}{dx}\ln(e^x\cos x) = \frac{d}{dx}\left(\ln e^x + \ln\cos x\right)$ $= \frac{d}{dx}x + \frac{1}{\cos x}\frac{d}{dx}\cos x$ $= 1 + \frac{-\sin x}{\cos x}$ $= 1-\tan x$ (shown) OR $\frac{d}{dx}\ln(e^x\cos x) = \frac{1}{e^x\cos x}\left[\left(e^x(-\sin x)\right)+\cos x(e^x)\right]$ $= \frac{e^x\left[(-\sin x)+\cos x\right]}{e^x\cos x}$ $= \frac{(-\sin x)+\cos x}{\cos x}$ $= 1-\tan x$ *(shown)*	

(ii) Using part (i), find $\int \tan x\,dx$. [2]

(ii)	$\frac{d}{dx}\ln(e^x\cos x)=1-\tan x$ $\tan x = 1-\frac{d}{dx}\ln(e^x\cos x)$ $\int\tan x\,dx = \int 1\,dx - \int\frac{d}{dx}\ln(e^x\cos x)\,dx$ $= x-\ln(e^x\cos x)+c$	

www.testpapersfree.com

Downloaded from SingaKids.com

(iii) Hence show that $\int_0^{\frac{\pi}{3}} \tan x \, dx = \ln 2$. [2]

(iii)	$\int_0^{\frac{\pi}{3}} \tan x \, dx = \left[x - \ln(e^x \cos x) \right]_0^{\frac{\pi}{3}}$ $= \left[\frac{\pi}{3} - \ln\left(e^{\frac{\pi}{3}} \cos\frac{\pi}{3} \right) \right] - \left[0 - \ln\left(e^0 \cos 0 \right) \right]$ $= \left[\frac{\pi}{3} - \ln\left(e^{\frac{\pi}{3}} \right) - \ln\left(\cos\frac{\pi}{3} \right) \right] - 0$ $= \left[\frac{\pi}{3} - \frac{\pi}{3} - \ln\left(\frac{1}{2} \right) \right]$ $= -\ln 2^{-1}$ $= \ln 2 \left(\text{shown} \right)$	

8 Point P lies on the curve $y = \frac{kx^2}{\sqrt{2-x}}$, where k is a constant. The x-coordinate of P is -2.

www.testpapersfree.com

Downloaded from SingaKids.com


(i) Given that the gradient of the normal at P is $\frac{1}{7}$, show that $k = 4$. [4]

(i)	$\frac{dy}{dx} = \frac{2kx\sqrt{2-x} - kx^2\left(\frac{1}{2}\right)(2-x)^{-\frac{1}{2}}(-1)}{2-x}$ $= \frac{2kx(2-x) + \frac{k}{2}x^2}{(2-x)^{\frac{3}{2}}}$ At $x = -2$, $\frac{dy}{dx} = -7$ $\frac{2k(-2)(2+2) + \frac{k}{2}(-2)^2}{(2+2)^{\frac{3}{2}}} = -7$ $\frac{-16k + 2k}{8} = -7$ $-14k = -56$ $k = 4$ (*shown*)	

www.testpapersfree.com

Downloaded from SingaKids.com

(i) The point P moves along the curve. Given that x is increasing at a rate of 0.9 units per second, find the rate of change of $(OP)^2$ at the point $x = 0.5$. [4]




(ii)	$(OP)^2 = x^2 + y^2$ $= x^2 + \left(\frac{4x^2}{\sqrt{2-x}}\right)^2$ $= x^2 + \frac{16x^4}{2-x}$ $\frac{d(OP)^2}{dx} = 2x + \frac{64x^3(2-x) - 16x^4(-1)}{(2-x)^2}$ $= 2x + \frac{128x^3 - 48x^4}{(2-x)^2}$ At $x = 0.5$, $\frac{d(OP)^2}{dx} = 2(0.5) + \frac{128(0.5)^3 - 48(0.5)^4}{(2-0.5)^2}$ $= \frac{61}{9}$ By chain rule, $\frac{d(OP)^2}{dt} = \frac{d(OP)^2}{dx} \times \frac{dx}{dt}$ $= \frac{61}{9} \times 0.9$ $= 6.1 \quad or \quad 6\frac{1}{10}\ units/s$ The rate of change of $(OP)^2 = 6.1$ units/s	

www.testpapersfree.com

Downloaded from SingaKids.com

9 The diagram shows the glass test tube of radius r cm and height $(h+r)$ cm with a hemisphere of radius r cm as the base. The surface area of the test tube is 12π.

(i) Show that $h=\dfrac{6-r^2}{r}$. [2]



(i)	$12\pi = 2\pi rh + 2\pi r^2$ $6 = rh + r^2$ $h = \dfrac{6-r^2}{r}$	

(ii) Show that the volume, V cm^3, of the test tube is $V = 6\pi r - \dfrac{1}{3}\pi r^3$. [2]

(ii)	$V = \pi r^2 h + \dfrac{2}{3}\pi r^3$ $V = \pi r^2\left(\dfrac{6-r^2}{r}\right) + \dfrac{2}{3}\pi r^3$ $V = 6\pi r - \pi r^3 + \dfrac{2}{3}\pi r^3$ $V = 6\pi r - \dfrac{1}{3}\pi r^3$	

www.testpapersfree.com

Downloaded from SingaKids.com

(iii) Calculate the stationary value of V and determine if the volume is a maximum or minimum. [5]

(iii)

$$\frac{dV}{dr} = 6\pi - \pi r^2$$

For stationary value of V, $\frac{dV}{dr} = 0$

$$6\pi - \pi r^2 = 0$$

$$\pi r^2 = 6\pi$$

$$r = \sqrt{6}, \quad \text{reject } r = -\sqrt{6}$$

$$= 2.4495$$

$$\text{Stationary volume of } V = 6\pi(2.4495) - \frac{1}{3}\pi(2.4495)^3$$

$$= 30.781$$

$$= 30.8\,\text{cm}^3$$

$$\frac{d^2V}{dr^2} = -2\pi r$$

$$= -2\sqrt{6}\,\pi < 0$$

Alternatively,

r	2.4495-	2.4495	2.4495+
$\frac{dV}{dr}$	+ve	0	-ve

Therefore volume is maximum.

www.testpapersfree.com

Downloaded from SingaKids.com

10 The equation of a circle, C_1, is $x^2+y^2-4x+8y+11=0$.

(i) Find the coordinates of the centre of C_1 and its radius. [3]

(i)	$x^2+y^2-4x+8y+11=0$ $(x-2)^2-4+(y+4)^2-16+11=0$ $(x-2)^2+(y+4)^2=3^2$ Centre $\equiv(2,-4)$ Radius $=3$ units	

(ii) Find the equation of the circle, C_2, which is a reflection of C_1 in the line $y=1$. [2]

(ii)	New centre $=(2,6)$ Equation of new circle is $(x-2)^2+(y-6)^2=9$ or $x^2+y^2-4x-12y+31=0$	

(iii) Given that the line $y-x+3=0$ intersects the circle, C_1, at two points, R and S, find the equation of the perpendicular bisector of the line RS. [4]

(iii)	$x^2+y^2-4x+8y+11=0$ ----------(1) Sub $y-x+3=0$ into (1) $x^2+(x-3)^2-4x+8(x-3)+11=0$ $x^2-x-2=0$ $(x+1)(x-2)=0$ $x=2$ or $x=-1$ when $x=2$, $y=-1$ when $x=-1$, $y=-4$ Gradient of $RS=\dfrac{-1+4}{2+1}=1$ Midpoint of $RS=\left(\dfrac{2-1}{2},\dfrac{-1+4}{2}\right)=\left(\dfrac{1}{2},-\dfrac{5}{2}\right)$ Gradient of normal of $RS=-1$ Equation of the perpendicular bisector of the line RS: $y+\dfrac{5}{2}=-\left(x-\dfrac{1}{2}\right)$ $y=-x-2$	

www.testpapersfree.com

Downloaded from SingaKids.com

11 A particle moves in a straight line, so that, t seconds after leaving a fixed point O its velocity, v is given by $v = \frac{36}{(t-3)^2} - 9$. Find

(i) the initial acceleration of the particle. [2]

(i)	$v = \frac{36}{(t-3)^2} - 9$ $= 36(t-3)^{-2} - 9$ $\frac{dv}{dt} = 36(-2)(t-3)^{-3}$ $= \frac{-72}{(t-3)^3}$ Initial acceleration is when $t = 0$ Initial acceleration $= \frac{-72}{(0-3)^3}$ $= \frac{8}{3}$ *or* $2\frac{2}{3}$ ms^{-2}	

(ii) the values of t when the particle is at instantaneous rest. [2]

(ii)	At instantaneous rest, $v = 0$, $\frac{36}{(t-3)^2} - 9 = 0$ $\frac{36}{(t-3)^2} = 9$ $(t-3)^2 = \frac{36}{9}$ $(t-3) = \pm\sqrt{4}$ $t = 3 + 2$ *or* $t = 3 - 2$ $t = 1$ *or* 5	

www.testpapersfree.com

Downloaded from SingaKids.com

(iii) the total distance travelled by the particle in the first 2 seconds. [6]

(iii)

$$s = \int \left(\frac{36}{(t-3)^2} - 9 \right) dt$$

$$= \int 36(t-3)^{-2} dx - \int 9\, dt$$

$$= \frac{36(t-3)^{-2+1}}{(-2+1)} - 9t + c$$

$$= -36(t-3)^{-1} - 9t + c$$

$$s = \frac{-36}{(t-3)} - 9t + c$$

At the fixed point, $t = 0, s = 0$

$$0 = \frac{-36}{-3} - 9(0) + c$$

$$c = -12$$

Therefore, $s = \frac{-36}{(t-3)} - 9t - 12$

Displacement, s from $t = 0$ to

$t = 1$ (first instantaneous rest)

$$s = \frac{-36}{(1-3)} - 9(1) - 12$$

$$= 18 - 9 - 12$$

$$= -3$$

Displacement, s when $t = 2$

$$s = \frac{-36}{(2-3)} - 9(2) - 12$$

$$= 36 - 30$$

$$= 6$$

Therefore total distance travelled

= 3 (on the left of fixed point O when $t = 1$)

+3 (back to fixed point from the left)

+6 (from fixed point when $t = 2$


= 12 m



www.testpapersfree.com

Downloaded from SingaKids.com

12



The diagram shows part of the curve $y = \sqrt{3x+4}$. The line AB is tangent to the curve at A. The points B and C lie on the x-axis, and C is a point on the curve. The gradient of the tangent of the curve at A is $\frac{3}{2}$. Find

(i) the coordinates of A and equation of the line AB. [5]

(i)

$$\frac{dy}{dx} = \frac{1}{2}(3x+4)^{\frac{1}{2}-1}(3)$$

$$= \frac{3}{2}(3x+4)^{-\frac{1}{2}}$$

$$= \frac{3}{2\sqrt{3x+4}}$$

Grad at A, $\frac{dy}{dx} = \frac{3}{2}$

$$\frac{3}{2} = \frac{3}{2\sqrt{3x+4}}$$

$$x = -1$$

When $x = -1$, $y = \sqrt{3(-1)+4}$

$$= 1$$

Coordinates of A are $(-1,1)$

Equation of AB is $y - 1 = \frac{3}{2}(x-(-1))$

Equation of AB is $y = \frac{3}{2}x + \frac{5}{2}$

www.testpapersfree.com

Downloaded from SingaKids.com

(ii) the coordinates of B and of C. [2]

(ii)	y-coordinate of B is 0, x-coordinate of B is $0 = \frac{3}{2}x + \frac{5}{2}$ $x = -\frac{5}{3}$ Coordinates of B are $\left(-\frac{5}{3}, 0\right)$ y-coordinate of C is 0, x-coordinate of C is $0 = \sqrt{3x+4}$ $x = -\frac{4}{3}$ Coordinates of C are $\left(-\frac{4}{3}, 0\right)$	

(iii) the area of the shaded region ABC. [3]

(iii)	Area of ABC = Area under line - Area under curve $= \int_{-\frac{5}{3}}^{-1}\left(\frac{3}{2}x + \frac{5}{2}\right)dx - \int_{-\frac{4}{3}}^{-1}\sqrt{3x+4}\,dx$ $= \left[\frac{3x^2}{4} + \frac{5}{2}x\right]_{-\frac{5}{3}}^{-1} - \left[\frac{(3x+4)^{\frac{3}{2}}}{\frac{3}{2}\times 3}\right]_{-\frac{4}{3}}^{-1}$ $= \left[\frac{3}{4} - \frac{5}{2} - \left(\frac{\frac{25}{3}}{4} - \frac{25}{6}\right)\right] - \frac{2}{9}\left[(3(-1)+4)^{\frac{3}{2}} - \left(3\times -\frac{4}{3} + 4\right)^{\frac{3}{2}}\right]$ $= \frac{1}{3} - \frac{2}{9}$ $= \frac{1}{9}$ units^2	

www.testpapersfree.com

Downloaded from SingaKids.com

	Alternatively, Area of ABC = Area of triangle - Area under curve $= \frac{1}{2}\left(\frac{5}{3}-1\right)\times 1 - \int_{-\frac{4}{3}}^{-1} \sqrt{3x+4}\, dx$ $= \frac{1}{3} - \frac{2}{9}\left[(3(-1)+4)^{\frac{3}{2}} - \left(3\times -\frac{4}{3}+4\right)^{\frac{3}{2}}\right]$ $= \frac{1}{9}$ units^2	

www.testpapersfree.com

Downloaded from SingaKids.com