More papers at www.testpapersfree.com

GAN ENG SENG SCHOOL
PRELIM 2022 S4EXP AM1_OLP

Marking Scheme
2022 Sec 4 Express PRELIM AM Paper 1

			Marker's Comments
1	Volume $= \frac{1}{3} \times$ Base Area $\times$ height $34\sqrt{5}+32 = \frac{1}{3} \times (2\sqrt{5}+7) \times$ height height $= \frac{3(34\sqrt{5}+32)}{2\sqrt{5}+7} \times \frac{2\sqrt{5}-7}{2\sqrt{5}-7}$ ---M1 $= \frac{3(68(5)-238\sqrt{5}+64\sqrt{5}-224)}{4(5)-49}$ ---M1 $= \frac{3(116-174\sqrt{5})}{-29}$ $= -12+18\sqrt{5}$ ---A1	**Note: To award up to 2 method marks for correct methods shown.** **3 marks**	* Read qn carefully: Qn wants to find height! Not p & q!
2	$\frac{3}{4} \times \left(\frac{\sqrt{3}}{2}\right)^4 - \frac{1}{3} \times (1)^3 + c = \frac{\sqrt{3}}{4} - \frac{1}{3}$ $c = \frac{\sqrt{3}}{4} - \frac{1}{3} - \frac{3\sqrt{3}}{8} + \frac{1}{3}$ $= -\frac{\sqrt{3}}{8}$ Equation of curve is $y = \frac{3}{4}\sin 4x - \frac{1}{3}\tan 3x - \frac{\sqrt{3}}{8}$ ---A1	**4 marks**	

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

3(a) 1.3 m ---B1 Subst. $x=0$ — **1 mark**

(b)(i)


$y=-0.01x^2+3.5x+1.3$

$=-0.01\left[x^2-350x\right]+1.3$ ---M1

$=-0.01\left[\left(x-\frac{350}{2}\right)^2-\left(\frac{350}{2}\right)^2\right]+1.3$ ---M1

$=-0.01\left[(x-175)^2-306.25\right]+1.3$

$=-0.01(x-175)^2+307.55$ ---A1



3 marks

(ii) 307.55 m ---B1 max pt's y — **1 mark** 5/5

4 ☆

$y=\frac{\ln(x+3)^3}{3x+9}$ ALWAYS Simplify ★ first where possible

$=\frac{3\ln(x+3)}{3x+9}$

$=\frac{\ln(x+3)}{x+3}$

$\frac{dy}{dx}=\frac{(\ \ +3)\frac{d}{dx}\ln(x+3)-\ln(x+3)\frac{d}{dx}(x+3)}{(x+3)^2}$

$=\frac{(x+3)\times\frac{1}{x+3}-\ln(x+3)(1)}{(x+3)^2}$ ---M1 (all terms correct)

OR $\frac{dy}{dx}=\frac{(3x+9)\left(\frac{3}{x+3}\right)-[\ln(x+3)^3](3)}{(3x+9)^2}$

$=\frac{3(x+3)\left(\frac{3}{x+3}\right)-3\ln(x+3)^3}{\text{or}}$

$=\frac{9-9\ln(x+3)}{9(x+3)^2}$

or $\frac{9-3\ln(x+3)^3}{(3x+9)^2}$

$=\frac{1-\ln(x+3)}{(x+3)^2}$

For y Given

Any $f(x)$, $\frac{dy}{dx}>0$

For $\frac{dy}{dx}>0$, $1-\ln(x+3)>0$ ---M1

$-\ln(x+3)>-1$

$\ln(x+3)<1$

$(x+3)<e$

$x<e-3$

$\therefore -3<x<e-3$ ---A1

take note that $x>-3$ is given in q2 & include it in $\in$ ans!!

4 marks

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

5

$$x^3-x^2+4x-4\overline{)\,5x^3-4x^2+15x-21}$$

$$\underline{-(5x^3-5x^2+20x-20)}\quad ---\text{M1}$$

$$x^2-5x-1$$

$$\frac{5x^3-4x^2+15x-21}{x^3-x^2+4x-4}$$

$$=5+\frac{x^2-5x-1}{(x^2+4)(x-1)}$$

$$\frac{ax+b}{x^2+4}$$

$$(ax+b)(x-1)+c(x^2+4)=x^2-5x-1---\text{M1}$$

Sub $x=1$

$5c=1-5-1$

$c=-1$ //

Sub $x=0$

$-b+4c=-1$

$-b+4(-1)=-1$

$-b-4=1$

$-b=3$

$b=-3$ //

Compare coefficient of x^2 :

$a+c=1$

$a-1=1$


$a=2$ //

$a=2, b=-3, c=-1---\text{A2,1,0}$

$$\frac{5x^3-4x^2+15x-21}{x^3-x^2+4x-4}$$

$$=5+\frac{2x-3}{x^2+4}-\frac{1}{x-1}---\text{A1}$$

Note:
To assign 0 mark if long division is not done.



Downloaded from SingaKids.com

More papers at www.testpapersfree.com

GAN ENG SENG SCHOOL
PRELIM 2022 S4EXP AM1 OLP

6(a)

$f(x) = 2(x-\frac{3}{2})(x+1)(x^2+bx+c)$ ---M1

$f(1) = -8$

$2(1-\frac{3}{2})(1+1)(1^2+b(1)+c) = -8$

$-2(1+b+c) = -8$

$(1+b+c) = 4$

$b+c = 3$ ---(1)---A1

$f(-2) = 28$

$2(-2-\frac{3}{2})(-2+1)((-2)^2+b(-2)+c) = 28$

$7(4-2b+c) = 28$

$4-2b+c = 4$

$-2b+c = 0$ ---(2)---A1

(1) − (2)

$(b+c)-(-2b+c) = 3$

$3b = 3$

$b = 1$

$c = 2$

$f(x) = 2(x-\frac{3}{2})(x+1)(x^2+x+2)$ ---A1

4 marks

* important to form f(x) first using its factors!!

(b)

$\frac{3}{2}$ and $x = -1$

2 marks

(a) $f(x) = (2x-3)(x+1)(x^2+bx+c)$

OR $f(1) = -8$

$(-1)(2)(1+b+c) = -8$

$1+b+c = 4$

$b+c = 3$ // -(1)

$f(-2) = 28$

$(-7)(-1)(4-2b+c) = 28$

Subst (2) into (1):

$\therefore f(x) = (2x-3)(x+1)(x^2+x+2)$

$2x^2-x-3$

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

middle

7(a)

$h = -25\cos\frac{2}{3}t + 35$

$\frac{2\pi}{b} = 3\pi$

$b = \frac{2}{3}$ ---B1

$a = -25$ ---B1


$c = 35$ ---B1

- Centre is 35m ⇒ axis of curve is ⇒ $c = 35$
- blade is 25m long ⇒ amplitude is 25 ⇒ $a = -25$ because height from ground is then drop
- period $= \frac{2\pi}{b} = 3\pi$ | $b = \frac{2\pi}{3\pi} = \frac{2}{3}$

3 marks

35

(b)



$a = -25$, $b = \frac{2}{3}$, $c = 35$ *

$h = -25\cos\frac{2}{3}t + 35$

Points labelled clearly on the h axis – 10 and 60 and t axis - 3π and 6π. (B1)

2 marks

* mark will crosses key points for an accurate curve!

(c)

$-25\cos\frac{2}{3}t + 35 = 42$

$-25\cos\frac{2}{3}t = 7$

$\cos\frac{2}{3}t = -\frac{7}{25}$

basic angle $\frac{2}{3}t = \cos^{-1}\left(\frac{7}{25}\right)$ ---M1

$= 1.287$ rad

$\frac{2}{3}t = \pi - 1.287$ ---M1

$t = \frac{3}{2}(\pi - 1.287)$

$t = 2.78$ seconds ---A1

one 4

* first bar 12m above

Note:
To award 0 mark if the equation is incorrect

3 marks

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

8(a)

Length of $QA = \sqrt{x^2 + 550^2}$

$= \sqrt{x^2 + 302500}$ ---B1

Cost of laying under river $= 8k\sqrt{x^2 + 302500}$

Cost of laying under ground $= k \times (1200 - x)$ ---B1

$\therefore C = 8k\sqrt{x^2 + 302500} + k(1200 - x)$

2 marks

(b)(i)

$$\frac{dC}{dx} = 8k \times \frac{1}{2}\left(x^2 + 302500\right)^{-\frac{1}{2}}(2x) - k$$

$$= \frac{8kx}{\sqrt{x^2 + 302500}}$$

* k is a constant! No need to use product rule!

$\frac{d}{dx}(kx) = k$

2 marks

(ii)

Let $\frac{dC}{dx} = 0$

$$\frac{8kx}{\sqrt{x^2 + 302500}} - k = 0$$

$$\frac{8kx}{\sqrt{x^2 + 302500}} = k$$

$$\sqrt{x^2 + 302500} = 8x$$

$x^2 + 302500 = 64x^2$ ---M1 (correct sq)

$63x^2 = 302500$

$x^2 = \frac{302500}{63}$ / 4801.5873

$= 69.2935$

$x = 69.3$ ---A1

Method 1: using 2nd derivative test

$$\frac{dC}{dx} = \frac{8kx}{\sqrt{x^2 + 302500}} - k$$

$$\frac{d^2C}{dx^2} = \frac{\left(\sqrt{x^2 + 302500} \times 8k\right) - 8kx\left(\frac{1}{2}\right)\left(x^2 + 302500\right)^{-\frac{1}{2}}(2x)}{\left(\sqrt{x^2 + 302500}\right)^2}$$

$$= \frac{(x^2 + 302500) \times 8k - 8kx^2}{\left(x^2 + 302500\right)^{\frac{3}{2}}} \text{---M1}$$

$$= \frac{2420000k}{\left(x^2 + 302500\right)^{\frac{3}{2}}}$$

At $x = 69.3$,

$$\frac{d^2C}{dx^2} = \frac{2420000k}{\left(69.3^2 + 302500\right)^{\frac{3}{2}}}$$

$= 0.0142k > 0$ ---A1

Since $\frac{d^2C}{dx^2} > 0$, total cost is minimum at $x = 69.3$

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(c)	Method 2 : 1st derivative test x | 68 | 69.3 | ~~69~~ 70 $\frac{dC}{dx}$ | $\frac{8k(68)}{\sqrt{68^2+302500}}-k$ | 0 | $\frac{8k(70)}{\sqrt{70^2+302500}}-k$ $=0.0153k-k$ | $=1.01k-k$ $-0.9847k<0$ | $0.01k>0$ Shape | \ | — | / M1 for attempt to find gradient on either side of $x=69.2$ and A1 for 2 correct values of $\frac{dC}{dx}$.	**Note:** **To award 0 mark if gradient values are found but only a sketch of the shape is produced** **4 marks**	
(c)	$C=8k\sqrt{69.2934^2+302500}+(1200-69.2934)k$ $=4434.783k+1130.7066k$ $=5565.4869k$ $=5570k$ ---B1	**1 mark**	
9(a)	$\text{LHS}=\frac{\sin 2A+\cos A}{1-\cos 2A+\sin A}$ $=\frac{2\sin A\cos A+\cos A}{1-(1-2\sin^2 A)+\sin A}$ ---M1,M1 $=\frac{\cos A(2\sin A+1)}{2\sin^2 A+\sin A}$ $=\frac{\cos A(2\sin A+1)}{\sin A(1+2\sin A)}$ ---A1 $=\cot A$	**3 marks**	

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

GAN ENG SENG SCHOOL
PRELIM 2022 S4EXP AM1 OLP

(b) ☆

$$\frac{1-\cos 4x+\sin 2x}{\sin 4x+\cos 2x}=5-2\sec^2 2x$$

$$\tan 2x=5-2\sec^2 2x$$

$$\tan 2x=5-2\left(1+\tan^2 2x\right)$$

$$2\tan^2 2x+\tan 2x-3=0 \text{ ----M1}$$

$$\tan 2x=\frac{-1\pm\sqrt{(1)^2-4(2)(-3)}}{2(2)}$$


$$=\frac{-1\pm\sqrt{25}}{4}$$

$$=\frac{-1\pm 5}{4} \text{ ---M1}$$

$$=-\frac{3}{2},1$$

which

x



$\pi, -(\pi+0.982794)$

$x=1.08, 2.65, -0.49, -2.06$

negative range

which

$\tan 2x=1$, basic ∠ 1.185398


$2x=\frac{\pi}{4}, \pi+\frac{\pi}{4}, -\left(\pi-\frac{\pi}{4}\right)$ $\pi-\frac{\pi}{4}$, $-\left(\pi-\frac{\pi}{4}\right)$, $-\left(2\pi-\frac{\pi}{4}\right)$

$-\frac{3\pi}{4}$, $-\frac{7\pi}{4}$

$x=0.39, 1.96, -2.36, -0.785$ -1.18, -2.75

A1 for correct positive angles, A1 for 2 correct negative angles ea

5 marks



10(a) See the △!!

S

$\angle CAB = \angle BCA$ (Base angles of isosceles triangle) B1 ★

$\therefore \angle BCT = \angle BCA$ (BC bisects $\angle ACT$) B1

use (longer note)
$= \angle BDC$ (alt. segment thm)
$= \angle CAB$
(As in same segment)

(b)

3 marks

wrong order but correct reasons & AA test
−1m

☆ → only use the same letters from these 2 △s !!
→ must match the order !!

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(c)

$\angle BCT + \angle BTC + \angle CBT = 180^\circ$

$\angle BCT + \angle BTC = 180^\circ - \angle CBT$ ---B1

From (b) $\angle CBT = \angle ACT$

$= 2 \times \angle BCA$

$= 2 \times \angle BAC$

$\angle BAC = \angle PDC$ (Angles in the same segment)---B1

$\therefore \angle BCT + \angle BTC = 180^\circ - 2 \times \angle PDC$

2 marks

11(a)

$\ln 3^{5x+1} = \ln 9^{x+5} + \log_2 16^{1-2x}$

$(5x+1)\ln 3 = \ln 3^{2(x+5)} + \log_2 (2^4)^{1-2x}$

$(5x+1)\ln 3 = 2(x+5)\ln 3 + 4(1-2x)$---M1

$5x\ln 3 + \ln 3 = 2x\ln 3 + 10\ln 3 + 4 - 8x$---M1 (expansion)

$3x\ln 3 + 8x = 9\ln 3 + 4$

$x(3\ln 3 + 8) = 9\ln 3 + 4$

$x = \dfrac{9\ln 3 + 4}{3\ln 3 + 8}$---A1

$\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y$

3 marks

(b)

$\log_4(x-3) + \dfrac{1}{2}\log_2(2x+1) = 1$

$\dfrac{\log_2(x-3)}{\log_2 4} + \dfrac{1}{2}\log_2(2x+1) = 1$---< Change base - M1>

$\dfrac{\log_2(x-3)}{2} + \dfrac{1}{2}\log_2(2x+1) = 1$

$\log_2(x-3) + \log_2(2x+1) = 2$

$\log_2(x-3)(2x+1) = 2$---M1

$(x-3)(2x+1) = 2^2$

$2x^2 + x - 6x - 3 - 4 = 0$

$2x^2 - 5x - 7 = 0$

$(2x-7)(x+1) = 0$

$x = \dfrac{7}{2}, -1$(reject)---A1

Note: Reject to be shown to be given 1 answer mark

3 marks

OR

change to base 10

$\dfrac{\lg(x-3)}{\lg 4} + \dfrac{\frac{1}{2}\lg(2x+1)}{\lg 2} = 1$

$\dfrac{\lg(x-3)}{\lg 4} + \dfrac{\lg(2x+1)}{2\lg 2} = 1$

$\lg(x-3)(2x+1) = \lg 4$

$(x-3)(2x+1) = 4$

↓

OR

$\dfrac{\log_4(x-3) + \log_4(2x+1)^{\frac{1}{2}}}{\log_4 2} = 1$

$\log_4(x-3) + \dfrac{\log_4(2x+1)^{\frac{1}{2}}}{\frac{1}{2}} = 1$

$\log_4(x-3)(2x+1) = 1$

↓

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

12(a)

At $t = 60$

Volume $= 30 \times 60$

$= 1800 \text{ cm}^3$

$\pi \times r^2 \times 200 = 1800$ ---M1

$r = \sqrt{\frac{9}{\pi}} = 1.69257$

$= 1.69 \text{ cm}$ ---A1

2 marks

(b)

$V = \pi r^2 h$

$= \pi r^2 (200)$ $\quad \frac{dV}{dt} = 30$

$= 200\pi r^2$

$\frac{dV}{dr} = 400\pi r$ ---M1

$\frac{dV}{dr} \times \frac{dr}{dt} = \frac{dV}{dt}$

$400\pi r \times \frac{dr}{dt} = 30$ ---M1

$\frac{dr}{dt} = \frac{30}{400\pi r}$

At $r = 2$ cm, $\frac{dr}{dt} = \frac{30}{400\pi(2)} = \frac{3}{80\pi}$

$= 0.0119 \text{ cm/s}$ ---A1

3 marks

13(a)

Since gradient of $OB = \frac{7}{4}$

$B(x, \frac{7}{4}x)$ ---M1

Gradient of AB = Gradient of DC

$\frac{\frac{7}{4}x - 1}{x + 4} = \frac{-1-(-4)}{3-(-1)}$ $\quad A(-4,1)$ $\quad C(3,-1)$ & $D(-1,-4)$

$\frac{\frac{7}{4}x - 1}{x + 4} = \frac{3}{4}$ ---M1

$7x - 4 = 3x + 12$

$4x = 16$

$x = 4$

$\therefore B(4,7)$ ---A1

3 marks

wrong concept!

gradient of $OB = \frac{7}{4}$

Subst $C(0,0)$ & $B(x,y)$

$\frac{y}{x} = \frac{7}{4}$

$\therefore y = 7$ & $x = 4$

But $\frac{y}{x} = \frac{7}{4} = \frac{14}{8} = \frac{21}{12}$ ...

Cannot use ratio here!

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

must be in * anti-clockwise direction

a)	Area of quadrilateral $ABCD$ $=\frac{1}{2}\begin{vmatrix}-4 & -1 & 3 & 4 & -4\\ 1 & -4 & -1 & 7 & 1\end{vmatrix}$ ---M1 $=\frac{1}{2}((16+1+21+4)-(-1-12-4-28))$ $=\frac{1}{2}(42+45)$ $=43.5$ units^2 ---A1	2 marks	
13(c) Mtd 1		2 marks	
13(d)	$C'(4+3,7-5)$---M1 Fastest mtd. $C'(7,2)$---A1 or B1,B1	2 marks	

B(c) Mtd 2 D(-1, -4) & C(3, -1)

Midpoint of DC $= \left(\frac{-1+3}{2}, \frac{-4-1}{2}\right)$

$= \left(1, -\frac{5}{2}\right)$

Gradient of ⊥ bisector of DC is $-\frac{4}{3}$.

$y+\frac{5}{2} = -\frac{4}{3}(x-1)$ M1

$y = -\frac{4}{3}x - \frac{7}{6}$

Subst. $x = 2.5$,

$y = -\frac{4}{3}(2.5) - \frac{7}{6}$

∴ * A1

13(d) midpoint of BD

Mtd 2 $= \left(\frac{4-1}{2}, \frac{7-4}{2}\right)$

$= \left(\frac{3}{2}, \frac{3}{2}\right)$

Let C be (x, y)

midpoint of AC'

$= \left(\frac{-4+x}{2}, \frac{1+y}{2}\right)$

$\frac{-4+x}{2} = \frac{3}{2}$, $\frac{1+y}{2} = \frac{3}{2}$

$-4+x = 3$, $1+y = 3$

$x = 7$ $y = 2$

∴ C(7, 2)

Mtd 3

eqn of BC' is $y = -\frac{5}{3}x + \frac{41}{3}$ ①

eqn of OC' is $y = \frac{3}{4}x - \frac{13}{4}$ ②

① = ②

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

130

14(a)

$$y=\left(\frac{3}{4}x-5\right)^{-2}+1$$

$$\frac{dy}{dx}=-2\left(\frac{3}{4}x-5\right)^{-3}\left(\frac{3}{4}\right)$$

$$=-\frac{3}{2}\left(\frac{3}{4}x-5\right)^{-3}---\text{M1}$$

At $x=6$, $\frac{dy}{dx}=-\frac{3}{2}\left(\frac{3}{4}(6)-5\right)^{-3}$

$$=-\frac{3}{2}\left(-\frac{1}{2}\right)^{-3}$$

$$=12---\text{A1}$$

At $x=6$, $y=\left(\frac{3}{4}(6)-5\right)^{-2}+1$

$$=\left(-\frac{1}{2}\right)^{-2}+1$$

$$=5---\text{B1}$$

Equation of PQ:

$y=12x+c$

Sub $(6,5)$

$5=12(6)+c---\text{M1}$

$c=-67$

$y=12x-67$

At Q, $y=0$

$x=\frac{67}{12}$ =

$\therefore Q(\frac{67}{12},0)---\text{A1}$

5 marks

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

Shaded area

$$= \int_0^6 \left[\left(\frac{3}{4}x - 5 \right)^{-2} + 1 \right] dx - \frac{1}{2}\left(6 - \frac{67}{12} \right)(5) \text{---M1,M1}$$

$$= \left[\frac{\left(\frac{3}{4}x - 5 \right)^{-1}}{-1\left(\frac{3}{4} \right)} + x \right]_0^6 - \frac{25}{24} \text{---<Correct integration - M1>}$$

$$= \frac{\left(\frac{3}{4}(6) - 5 \right)^{-1}}{-1\left(\frac{3}{4} \right)} + 6 - \frac{\left(\frac{3}{4}(0) - 5 \right)^{-1}}{-1\left(\frac{3}{4} \right)} - \frac{25}{24}$$

$$= \frac{26}{3} - \frac{4}{15} - \frac{25}{24} \text{---M1}$$

$$= 7\frac{43}{120} \text{ units}^2 \text{---A1}$$

5 marks

$$\left[\frac{1}{\left(\frac{3}{4}x - 5 \right)} \right]$$

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

O

GAN ENG SENG SCHOOL
Preliminary Examination 2022

CANDIDATE NAME: Marking Scheme

CLASS: ____ INDEX NUMBER: ____

ADDITIONAL MATHEMATICS
Paper 2
Sec 4 Express

4049/02
29 August 2022
2 hours 15 min

Candidates answer on the Question Paper.
No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your class, index number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft HB pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of a scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.
The total marks for this paper is 90.

	For Examiner's Use
Total	90

This paper consists of 20 printed pages including the cover page.

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \cdots + \binom{n}{r}a^{n-r}b^r + \cdots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}ab\sin C$$

PartnerInLearning
143

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

1. Show that the equation $9^{x+1} - 5(3^{x+1}) = 10$ has only one solution and find its value correct to two decimal places. [6]

Solution

$9^{x+1} - 5(3^{x+1}) = 10$

$9^x \times 9 - 5(3^x \times 3) = 10$

$3^{2x} \times 9 - 15(3^x) - 10 = 0$ M1

Let $u = 3^x$

$9u^2 - 15u - 10 = 0$

M1

M1

M1

B1

A1

6/6

OR

$3^{2(x+1)} - 5(3^{x+1}) = 10$

$(3^{x+1})^2 - 5(3^{x+1}) = 10$

Let $u = 3^{x+1}$

$u^2 - 5u - 10 = 0$

$u = \frac{5 \pm \sqrt{(-5)^2 - 4(1)(-10)}}{2(1)}$

$= \frac{5 \pm \sqrt{65}}{2}$

$= 6.53$

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

2 Solve the equation $2x^3 - 7x + 2 = 0$, leaving non-rational roots in the form $a \pm b\sqrt{2}$, where a and b are rational numbers. [5]

Solution

Let $f(x) = 2x^3 - 7x + 2$

Try $f(-2) = 2(-2)^3 - 7(-2) + 2$

$= -16 + 14 + 2$

$= 0$

Therefore $x + 2$ is a factor. M1 M1

By inspection $2x^3 - 7x + 2 = (x + 2)(2x^2 + ax + 1)$, where a is a constant.

Compare terms in x: $-7x = 2ax + x$

$-7 = 2a + 1$

$2a = -8$

$a = -4$ M1 M1

mtd 2 long division

$2x - 4x + 1$

Therefore $2x^3 - 7x + 2 = 0$

$(x + 2)(2x^2 - 4x + 1) = 0$

Either $x = -2$ or $2x^2 - 4x + 1 = 0$

If $2x^2 - 4x + 1 = 0$

$$x = \frac{4 \pm \sqrt{(-4)^2 - 4 \times 2 \times 1}}{4}$$ M1

$$x = \frac{4 \pm \sqrt{8}}{4}$$

$$x = 1 \pm \frac{1}{2}\sqrt{2}$$

Ans: $x = -2$ or $x = 1 \pm \frac{1}{2}\sqrt{2}$ A1, 1

A1 A1

PartnerInLearning
145

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

3 A particle moves from rest at A and comes to rest at B. Its speed, in m/s, when travelling from A to B is given by the equation $v = 10t - \frac{1}{2}t^2$, where t is the time in seconds starting from A.

Show that the particle has a speed of 5 m/s or more for $6\sqrt{10}$ s. [4]

$v \geq 5$

Solution

$v \geq 5$

$10t - \frac{1}{2}t^2 \geq 5$

$20t - t^2 - 10 \geq 0$

$t^2 - 20t + 10 \leq 0$ m1

If $t^2 - 20t + 10 = 0$

$t = \frac{20 \pm \sqrt{(-20)^2 - 4 \times 1 \times (-10)}}{2}$ m1

$t = \frac{20 \pm \sqrt{360}}{2}$

$t = \frac{20 \pm 6\sqrt{10}}{2}$

$t = 10 \pm 3\sqrt{10}$

$t_1 = 10 - 3\sqrt{10}$ and $t_2 = 10 + 3\sqrt{10}$ s m1

Interval of time is $10 + 3\sqrt{10} - (10 - 3\sqrt{10})$ s

$= 6\sqrt{10}$ s A1

$-\frac{1}{2}t^2 + 10t - 5 = 0$ M1

OR

$t = \frac{-10 \pm \sqrt{(10^2) - 4(-\frac{1}{2})(-5)}}{2(-\frac{1}{2})}$ M1

$= \frac{-10 \pm \sqrt{90}}{-1}$


$= \frac{-10 \pm 3\sqrt{10}}{-1}$ M1

A1

$= \frac{-10 + 3\sqrt{10}}{-1}$ or $\frac{-10 - 3\sqrt{10}}{-1}$

$= 10 - 3\sqrt{10}$ $= 10 + 3\sqrt{10}$


$v = -\frac{1}{2}t^2 + 10t$



Downloaded from SingaKids.com

More papers at www.testpapersfree.com

4



The curve $y = \ln x$ cuts the x-axis at P. The area, A units, is enclosed by the curve, the x-axis and the line $x = e^2$.

Explain why $e^2 - 1 < \int_1^{e^2} y\,dx < 2(e^2 - 1)$. [4]

Solution

If $y = \ln x$ meets $y = 0$,

$\ln x = 0$,

$x = 1$

Therefore P is (1, 0). m1 M1

Draw QR parallel to the y-axis through $(e^2, 0)$ to meet the curve at R.
Then draw rectangle $PQRS$.

If $x = e^2$, $y = \ln e^2 = 2$

Therefore $Q = (e^2, 0)$ and R is at $(e^2, 2)$ m1 M1

Area of $\Delta PQR < \int_1^{e^2} y\,dx <$ Area of $PQRS$ M1

$\frac{1}{2}(e^2 - 1) \times 2 < \int_1^{e^2} y\,dx < 2(e^2 - 1)$ m1

$e^2 - 1 < \int_1^{e^2} y\,dx < 2(e^2 - 1)$ A1 A1

* DO NOT


$\int \ln x\,dx = \frac{1}{x} + C$

Wrong concept !!

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(5) In ΔABC, $AC = 12$ cm, $\angle ABC$ is a right angle, $\angle ACB = \frac{\pi}{6}$ radians and M is the mid-point of BC. Without the use of a calculator, find the value of the integer k, such that $\angle CAM = \sin^{-1}\left(\frac{\sqrt{k}}{14}\right)$.



Solution [5]

$\sin\left(\frac{\pi}{6}\right) = \frac{AB}{12}$

$AB = 12\sin\left(\frac{\pi}{6}\right) = 6$ cm (m1) M1

$\cos\left(\frac{\pi}{6}\right) = \frac{BC}{12}$

$BC = 12\cos\left(\frac{\pi}{6}\right)$

$BC = 12 \times \frac{\sqrt{3}}{2} = 6\sqrt{3}$ cm

(m is midp't) $MC = BM = 3\sqrt{3}$ cm (m1)

$AM^2 = AB^2 + BM^2$

$= 6^2 + (3\sqrt{3})^2$

$= 36 + 27 = 63$

$AM = 3\sqrt{7}$ cm (m1) M1

OR

$AM^2 = AC^2 + MC^2 - 2(AC)(MC)\cos\frac{\pi}{6}$ M1

$= 12^2 + \left(6\cos\frac{\pi}{6}\right)^2 - 2(12)\left(6\cos\frac{\pi}{6}\right)\cos\frac{\pi}{6}$

$= 63$

By the Sine Rule, $\frac{\sin\angle CAM}{MC} = \frac{\sin\left(\frac{\pi}{6}\right)}{AM}$ M1

$\frac{\sin\angle CAM}{\frac{1}{2}\sqrt{108}} = \frac{\sin\left(\frac{\pi}{6}\right)}{\sqrt{63}}$

$\frac{\sin\angle CAM}{3\sqrt{3}} = \frac{\frac{1}{2}}{3\sqrt{7}}$ (m1) M1

$\sin\angle CAM = \frac{\frac{1}{2} \times 3\sqrt{3}}{3\sqrt{7}}$

$\sin\angle CAM = \frac{\sqrt{3}}{2\sqrt{7}} = \frac{\sqrt{21}}{14}$

$\angle CAM = \sin^{-1}\left(\frac{\sqrt{21}}{14}\right)$

Therefore $k = 21$ Ans (A1) A1

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

$T_{r+1} = \binom{n}{r} a^{n-r} b^r$

6 (a) (i) In the binomial expansion of $\left(x+\frac{1}{ax^2}\right)^8$ where a is a positive integer, the coefficient of x^2 and $\frac{1}{x}$ are equal. Find the value of a. [4]

Solution

$$\left(x+\frac{1}{ax^2}\right)^8 = x^8 + \binom{8}{1}(x^7)\left(\frac{1}{ax^2}\right) + \binom{8}{2}(x^6)\left(\frac{1}{ax^2}\right)^2 + \binom{8}{3}(x^5)\left(\frac{1}{ax^2}\right)^3 + \ldots\ldots$$

M1

$$= x^8 + \frac{8}{a}x^5 + 28x^6 \times \frac{1}{a^2x^4} + 56x^5 \times \frac{1}{a^3x^6} + \ldots\ldots\ldots\ldots$$

$$= x^8 + \frac{8}{a}x^5 + \frac{28}{a^2}x^2 + \frac{56}{a^3}\left(\frac{1}{x}\right) + \ldots\ldots\ldots\ldots$$ M1

$\frac{28}{a^2} = \frac{56}{a^3}$ m1 M1

$28a^3 = 56a^2$ * do not cancel a !!

$a^3 - 2a^2 = 0$

$a^2(a-2) = 0$

Since $a \neq 0, a = 2$ Ans A1 A1

(ii) With the value of a found in part (i), show that there is no term independent of x in the expansion of $\left(x-\frac{1}{x^2}\right)\left(x+\frac{1}{ax^2}\right)^8$. [3]

Solution

$$\left(x-\frac{1}{x^2}\right)\left(x+\frac{1}{ax^2}\right)^8 = \left(x-\frac{1}{x^2}\right)\left(x+\frac{1}{2x^2}\right)^8$$

$$\left(x-\frac{1}{x^2}\right)\left(x+\frac{1}{2x^2}\right) = \left(x-\frac{1}{x^2}\right)\left(x^8 + \frac{8x^5}{2} + \frac{28x^2}{2^2} + \frac{56}{2^3}\left(\frac{1}{x}\right) + \ldots\right)$$ M1

$$= \left(x-\frac{1}{x^2}\right)\left(x^8 + 4x^5 + 7x^2 + \frac{7}{x} + \ldots..\right)$$

Term independent of x is $\left(x \times \frac{7}{x}\right) - \left(\frac{1}{x^2} \times 7x^2\right) + \ldots$ M1

$= 7 - 7$

$= 0$

Therefore, there is no term independent of x. A1

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(b) Calculate the term independent of x in the binomial expansion of $\left(x-\frac{1}{2x^5}\right)^{18}$. [3]

Solution

General term $= \binom{18}{r}\left(x^{18-r}\right)\left(-\frac{1}{2x^5}\right)^r$

$= \binom{18}{r}\left(x^{18-r}\right)\frac{(-1)^r}{2^r\left(x^{5r}\right)}$ M1

For term independent of x, $x^{18-r} = x^{5r}$

$18 - r = 5r$

$6r = 18$

$r = 3$ M1

Term independent of $x = \binom{18}{3} \times \frac{(-1)^3}{2^3}$

$= -102$ **Ans** A1

OR

$\left(x-\frac{1}{2x^5}\right)^{18}$

$= x^{18} + \binom{18}{1}(x^{17})\left(-\frac{1}{2x^5}\right)^1 + \binom{18}{2}(x^{16})\left(-\frac{1}{2x^5}\right)^2 + \binom{18}{3}(x^{15})\left(-\frac{1}{2x^5}\right)^3 + \cdots$

Term independent of $x = \binom{18}{3}(x^{15})\left(-\frac{1}{2x^5}\right)^3$

$= 816\left(-\frac{1}{8}\right)$

$= -102$.

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

7 (i) Find the values of p for which the line $y = x + 1$ is a tangent to the curve $y = x^2 + (2p + 3)x + p + 4$. [4]

Solution

If $y = x^2 + (2p + 3)x + p + 4$ meets $y = x + 1$

$x^2 + (2p + 3)x + p + 4 = x + 1$

$x^2 + (2p + 3)x + p + 4 - x - 1 = 0$

$x^2 + (2p + 2)x + p + 3 = 0$ M1

For equal roots, discriminant = 0

$(2p + 2)^2 - 4(1)(p + 3) = 0$ M1

$4p^2 + 4p - 8 = 0$

$p^2 + p - 2 = 0$

$(p + 2)(p - 1) = 0$

$p = -2$ or $p = 1$ Ans A1, A1

(ii) With the values of p found in part (i), find the coordinates of the points where the line meets the curve. [4]

Solution

If $p = 1$, $x^2 + (2\times1+2)x + 1 + 3 = 0$

$x^2 + 4x + 4 = 0$

$(x + 2)^2 = 0$

$x = -2$ M1

If $x = -2$, $y = -2 + 1 = -1$

Therefore, one point is $(-2, -1)$. A1

If $p = -2$, $x^2 + (2(-2) + 2)x + (-2) + 3 = 0$

$x^2 - 2x + 1 = 0$

$(x - 1)^2 = 0$

$x = 1$ M1

If $x = 1$, $y = 1 + 1 = 2$.


Therefore, the other point is (1, 2) A1

PartnerInLearning
151

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

8



The figure shows a piece of cardboard in the shape of a trapezium in which $\angle BAD$ and $\angle ABC$ are right angles. $CD = 20$ cm, $BC = 8$ cm and $\angle ADC = \theta$ radians.

(i) Show that the perimeter, P cm, is given by $P = 20\cos\theta + 20\sin\theta + 36$. [2]

Solution

Draw CE perpendicular to AD

In $\Delta\ CDE$, $\cos\theta = \dfrac{DE}{20}$

$DE = 20\cos\theta$ cm

$\sin\theta = \dfrac{CE}{20}$

$CE = 20\sin\theta$

Therefore $P = AB + BC + CD + AD$

$= 20\sin\theta + 8 + 20 + 20\cos\theta + 8$

$P = 20\cos\theta + 20\sin\theta + 36$ Shown A1

M1

(ii) Express P in the form $R\cos(\theta - \alpha) + k$, where α is acute. [3]

Solution

$R^2 = 20^2 + 20^2$

$= 800$

$R = \sqrt{800}$

$= 20\sqrt{2}$ M1

M1

$\tan\alpha = \dfrac{20}{20}$

$\alpha = \dfrac{\pi}{4}$ M1

M1

Therefore $P = 20\sqrt{2}\cos\left(\theta - \dfrac{\pi}{4}\right) + 36$ Answer A1

$P = 20\sqrt{2}\cos(\theta - 0.785) + 36$

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(iii) If θ, can vary, find the maximum value of P and the corresponding value of θ. [2]

Solution

Maximum value of P is when $\cos\left(\theta-\frac{\pi}{4}\right)=1$

Therefore $P = 20\sqrt{2}\times 1+36$

64.3 / $P = 20\sqrt{2}+36$ Ans simplest & exact value B1

As θ is acute, $\theta-\frac{\pi}{4}=0$

$\theta = \frac{\pi}{4}$ 0.785 rad B1

(iv) [2]

Solution

If $P = 60$ cm

$20\sqrt{2}\cos\left(\theta-\frac{\pi}{4}\right)+36=60$

$20\sqrt{2}\cos\left(\theta-\frac{\pi}{4}\right)= 24$

$\cos\left(\theta-\frac{\pi}{4}\right)=\frac{24}{20\sqrt{2}}$ M1

Basic angle = 0.5576 00 rad (6 s.f.)

$\theta-\frac{\pi}{4}=0.557600$ or $\theta-\frac{\pi}{4} = -0.557600$


$\theta = \frac{\pi}{4}+0.5576$ $\theta = \frac{\pi}{4} - 0.5576$

$\theta =$ 1.34 radian (3 s.f.) Answer $= 0.228$ rad A1

$0 < \theta < \frac{\pi}{2}$

$-\frac{\pi}{4} < \theta-\frac{\pi}{4} < \frac{\pi}{2}-\frac{\pi}{4}$

$-0.785 < \theta-\frac{\pi}{4} < 0.785$



$\therefore \theta = 1.34$ rad, 0.228 rad (A1)

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

Show ⇒ show every step!

9 (i) Show that $\frac{d}{dx}\left(x(3x-2)^{\frac{4}{3}}\right)=(7x-2)(3x-2)^{\frac{1}{3}}$ [3]

Solution

$$\frac{d}{dx}\left\{x(3x-2)^{\frac{4}{3}}\right\}=(3x-2)^{\frac{4}{3}}\frac{d}{dx}(x)+x\frac{d}{dx}(3x-2)^{\frac{4}{3}}$$

$$=(3x-2)^{\frac{4}{3}}\times 1+x\times\frac{4}{3}(3x-2)^{\frac{1}{3}}\times 3 \quad \text{M1}$$

$$=(3x-2)^{\frac{1}{3}}[3x-2+4x] \quad \text{M1}$$

$$=(3x-2)^{\frac{1}{3}}(7x-2) \quad \text{Proven} \quad \text{A1}$$

(ii) Hence evaluate $\int x(3x-2)^{\frac{1}{3}}dx$. [4]

Solution

$\int(3x-2)^{\frac{1}{3}}(7x-2)dx=x(3x-2)^{\frac{4}{3}}+c_1$ where c_1 is a constant M1

$\int 7x(3x-2)^{\frac{1}{3}}dx-\int 2(3x-2)^{\frac{1}{3}}dx=x(3x-2)^{\frac{4}{3}}+c_1$ M1

$\int 7x(3x-2)^{\frac{1}{3}}dx=\int 2(3x-2)^{\frac{1}{3}}dx+x(3x-2)^{\frac{4}{3}}+c_1$

$\int 7x(3x-2)^{\frac{1}{3}}dx=\frac{2(3x-2)^{\frac{4}{3}}}{\frac{4}{3}\times 3}+c_2+x(3x-2)^{\frac{4}{3}}+c_1$ where c_2 is a constant M1

$\int 7x(3x-2)^{\frac{1}{3}}dx=\frac{(3x-2)^{\frac{4}{3}}}{2}+c_2+x(3x-2)^{\frac{4}{3}}+c_1$

$\int x(3x-2)^{\frac{1}{3}}dx=\frac{1}{14}(3x-2)^{\frac{4}{3}}+\frac{1}{7}x(3x-2)^{\frac{4}{3}}+c_1+c_2$ * Factorise

$\int x(3x-2)^{\frac{1}{3}}dx=\frac{1}{14}(3x-2)^{\frac{4}{3}}(1+2x)+c$ where c is a constant. Ans A1

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(iii) Find the value of $\int_{-\frac{1}{2}}^{\frac{2}{3}} x(3x-2)^{\frac{1}{3}}dx$ and explain what the result implies about the curve $y=x(3x-2)^{\frac{1}{3}}$. [3]

Solution

$$\int_{-\frac{1}{2}}^{\frac{2}{3}} x(3x-2)^{\frac{1}{3}}dx = \frac{1}{14}\left[(3x-2)^{\frac{4}{3}}(2x+1)\right]_{-\frac{1}{2}}^{\frac{2}{3}}$$

must show subst^{n} of x values

$$= \frac{1}{14}\left(3\times\frac{2}{3}-2\right)^{\frac{4}{3}}\left(2\times\frac{2}{3}+1\right)-\left(3\left(-\frac{1}{2}\right)-2\right)^{\frac{4}{3}}\left(2\left(-\frac{1}{2}\right)+1\right)$$ M1

$$= 0$$ A1

The area enclosed by the curve and the x-axis from $x=-\frac{1}{2}$ to $x=0$ is equal to the area enclosed by the curve and the x-axis from $x=0$ to $x=\frac{2}{3}$. B1

& below

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

10 A sports car driven along a straight road passes a traffic junction A at p m/s. A little later, it passes a second traffic junction B with a speed of 40 m/s. Between A and B, the speed of the car, v m/s. is given by $v = 5e^{0.05t} + 10$ where t is the time in seconds after passing A.

(i) State the value of p. [1]

Answer
When $t = 0$, $p = 5e^0 + 10$
$p = 5 + 10$
$= 15$ Ans B1

(ii) Show that the time taken to travel from A to B is $20\ln 6$ seconds. [3]

Solution
$5e^{0.05t} + 10 = 40$
$5e^{0.05t} = 30$
$e^{0.05t} = 6$ m1 M1
$\ln e^{0.05t} = \ln 6$
$0.05t = \ln 6$ m1
$t = \frac{\ln 6}{0.05}$ $\frac{1}{20}t = \ln 6$ M1
$t = 20\ln 6$ $t = 20\ln 6$
Time taken to travel from A to $B = 20\ln 6$ seconds A1

(iii) Calculate the distance AB. [3]

Solution
Displacement, $s = \int_0^{20\ln 6} v\,dt$
$= \int_0^{20\ln 6} (5e^{0.05t} + 10)\,dt$ m1 M1
$= \left[\frac{5e^{0.05t}}{0.05} + 10t\right]_0^{20\ln 6}$ m1 M1
$= \left[100e^{0.05t} + 10t\right]_0^{20\ln 6}$
$= 100e^{0.05\times 20\ln 6} + 10\times 20\ln 6 - 100e^0$
$= 600 + 358.35 - 100$
$= 835.35$ m
$= 858$ m (3 s.f.) A1 A1

Mtd 2

$s = 100e^{0.05t} + 10t + C$

when $t = 0$, $s = 0$,

$0 = 100 + C$

$C = -100$

$s = 100e^{0.05t} + 10t - 100$

At $t = 20\ln 6$,

$s = 858$

PartnerInLearning 156

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(iv) Find the acceleration of the car when $t = 30$ s. [2]

Solution

Acceleration, $a = \frac{dv}{dt}$

$$a = \frac{d}{dt}(5e^{0.05t} + 10) \quad \text{M1}$$

$$= 5 \times 0.05e^{0.05t}$$

When the time is 30 s, $a = 5 \times 0.05e^{0.05 \times 30}$

$$= 1.12 \text{ m/s}^2 \quad \text{A1}$$

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

11 Points $A(8, 1)$ and $B(1, 2)$ lie on a circle whose centre is C. The line $4y = 3x - 20$ is tangent to the circle at A.

(i) Find the equation of the normal to the circle at A. [2]

Solution

Equation of tangent is $y = \frac{3}{4}x - 5$.

Gradient of normal at A is $-\frac{4}{3}$. m1 M1

Equation of normal $y - 1 = -\frac{4}{3}(x - 8)$

$3y - 3 = -4x + 32$

$3y + 4x = 35$ $y = -\frac{4}{3}x + \frac{35}{3}$ A1 $11\frac{2}{3}$

(ii) Find the equation of the circle. [6]

Solution

Gradient of $AB = \frac{1-2}{8-1} = -\frac{1}{7}$

Gradient of perpendicular bisector of $AB = 7$. m1 M1

Mid-point of $AB = \left(\frac{8+1}{2}, \frac{2+1}{2}\right)$

$= \left(\frac{9}{2}, \frac{3}{2}\right)$

Equation of perpendicular bisector of AB is

$y - \frac{3}{2} = 7\left(x - \frac{9}{2}\right)$

$y - \frac{3}{2} = 7x - \frac{63}{2}$ ★

$y = 7x - 30$ m1 M1

★ ⊥ bisector of chord AB cuts through centre.

If $y = 7x - 30$ meets $3y + 4x = 35$

$3(7x - 30) + 4x = 35$

$21x - 90 + 4x = 35$

$25x = 125$

$x = 5$ m1 M1

If $x = 5$, $y = 7 \times 5 - 30 = 5$

Therefore centre of circle $C(5, 5)$. m1 M1

$r^2 = (8-5)^2 + (1-5)^2$

$r^2 = 3^2 + 4^2$

$r^2 = 25$ m1 M1

Equation of circle is $(x-5)^2 + (y-5)^2 = 25$. A1 A1

$x^2 - 10x + y^2 - 10y + 25 = 0$

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(iii) Explain why the coordinate axes are tangents to the circle. [1]

Answer
Since the centre is at (5, 5), the centre is 5 units from each axis and the radius of the circle is 5 units. Therefore, the coordinate axes are tangents to the circle. B1

12 The table shows experimental values of x and y.

x	2	3	4	5	6
y	11.8	16.2	23.5	35.5	55.2

It is known that x and y are related by the equation $y = Ae^{kx} + 5$, where A and k are constants.

(i) Explain how a straight line graph may be drawn to estimate the values of A and of k. [2]

Solution
$y = Ae^{kx} + 5$
$y - 5 = Ae^{kx}$
$\ln(y-5) = \ln(Ae^{kx})$ M1
$\ln(y-5) = \ln A + kx$
Plot $\ln(y-5)$ against x. A1

PartnerInLearning
159

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

(ii) Draw a straight line graph to obtain estimated values of A and k. [5]

Solution

x	2	3	4	5	6
$\ln(y-5)$	1.92	2.42	2.92	3.42	3.92

M1

Answers

From the graph, $\ln A \approx 0.95$

$A \approx 2.59$ B1

$k \approx \dfrac{3-1.5}{4.1-1.1}$

$k \approx 0.5$ B1

(iii) Use your graph to find the value of x when $y = 30$. [1]

When $y = 30$, $\ln(30-5) = 3.22$

$x \approx 4.6$ Ans Accept 4.6 – 4.7) B1

(iv) By drawing a suitable straight line on the same axes, solve the equation [3]

$Ae^{kx} - 30(2^{-x})$

Given $y = Ae^{kx} + 5$

Solution

$Ae^{kx} + 5 = 30(2^{-x}) + 5$

$y = 30(2^{-x}) + 5$

$y - 5 = 30(2^{-x})$

$\ln(y-5) = \ln 30 + \ln(3^{-x})$

$\ln(y-5) = \ln 30 - x\ln 2$

Draw $\ln(y-5) = \ln 30 - x\ln 2$ M1 M1


x	0	5
$\ln(y-5)$	3.4	− 0.06

From the graph, $x \approx 2.05$ Accept (2.0 to 2.1) A1

PartnerInLearning
160

Downloaded from SingaKids.com

More papers at www.testpapersfree.com



END OF PAPER

PartnerInLearning
161

Downloaded from SingaKids.com

More papers at www.testpapersfree.com

Downloaded from SingaKids.com