Full Name	Class Index No	Class

Anglo-Chinese School
(Barker Road)

PRELIMINARY EXAMINATION 2023
SECONDARY FOUR EXPRESS / FIVE NORMAL (ACADEMIC)

ADDITIONAL MATHEMATICS
4049
PAPER 2

2 HOURS 15 MINUTES

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your index number and name on all the work you hand in.
Write in dark blue or black pen.

Answer **all** questions.
If working is needed for any question it must be shown with the answer.
Omission of essential working will result in loss of marks.
The use of an approved scientific calculator is expected, where appropriate.
If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.
For π, use either your calculator value or 3.142, unless the question requires the answer in terms of π.

The number of marks is given in brackets [ ] at the end of each question or part question.
The total of the marks for this paper is 90.

For Examiner's Use

This question paper consists of **20** printed pages and **2** blank pages.

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 Mr Tan invested some money in stock in May.
The value of a stock, \$*y*, varies with time, *x*, can be modelled by the equation
$y = -\frac{1}{3}x^2 + 2x + 1$, where *y* is in thousands and *x* is in months.

(a) State the initial value of the stock. [1]

(b) Express y in the form $a(x-b)^2 + c$. [2]

(c) Determine the best time to sell off his stocks to make a maximum profit.
Find the value of the stock at this time. [2]

Downloaded from SingaKids.com

2 It is given that $f(x) = ax^3 + 2bx^2 - 34x + 12$, where a and b are constants, has a factor of $x - 3$ and leaves a remainder of 32 when divided by $x + 1$.

(a) Find the value of a and of b. [4]

(b) Find the range of values of k for which the line $y + 9 = kx$ intersects the curve $x = \sqrt{y - 3x}$ at two distinct points. [4]


Downloaded from SingaKids.com

3 **(a)** Show that $\dfrac{2\cos^2\theta - \sin\theta\cos\theta + 1}{\sin^2\theta} = 3\cot^2\theta - \cot\theta + 1$. [3]

(b) Hence solve the equation $2\cos^2\theta - \sin\theta\cos\theta + 1 = \sin^2\theta$ for $-\pi \le \theta \le \pi$. [4]

Downloaded from SingaKids.com

4 A coffee filter is in the shape of a right circular cone. The cone has a base radius of 3 cm and a height of 12 cm. The coffee powder, as seen in the cross-section of the filter below, has a depth of 4 cm. Let h be the height of the hot water in the filter at time t seconds. Assume that the amount of water in the coffee powder is negligible.



(a) Show that the volume of hot water, V, in the filter at the time t seconds is given by $V = \frac{\pi}{48}(h+4)^3 - \frac{4\pi}{3}$. [3]

[The volume of a cone of height h and base radius r is $\frac{1}{3}\pi r^2 h$]


Downloaded from SingaKids.com

Initially, the filter contains only coffee powder. A machine dispenses hot water into the filter at a constant rate of 5 cm^3 per second and hot water drips out of the filter at a constant rate of 2 cm^3 per second.

(b) Find the exact rate of change of the depth of water when $h = 8$. [4]

Downloaded from SingaKids.com

5



The diagram shows an isosceles triangle PQR with vertices at $P(0, k)$, $Q(3, 9)$ and $R(h, 6)$, where h and k are positive constants and $h>k$.

(a) Show that $h + k = 12$. [3]

Downloaded from SingaKids.com


(b) It is now given that $h = 7$.
The point S is such that $PQRS$ is a kite. Given that S lies on the line $21y = 7x - 8$, find the coordinates of S. [4]

Downloaded from SingaKids.com

(c) Find the area of the kite. [2]

Downloaded from SingaKids.com

6



In the diagram above, *CDEF* is a circle and the line *AB* is a tangent to the circle at *E*. *ADC* and *CFB* are straight lines and *AE* = *EB* and *CF* = *FB*.

(a) Prove that *AC* is parallel to *EF*. [1]

(b) Prove that angle *AED* = angle *CEF* . [2]

(c) Prove that triangle *AED* is similar to triangle *CEF* . [2]

Downloaded from SingaKids.com

7 Theresa bought a house on 2 January 1970. The house was valued by a local estate agent on the same date every 10 years up to 2010.

The valuation price, V, of the house is related to t, the number of years since 1970. The variables V and t can be modelled by the equation $V = pq^t$, where p and q are constants.

The table below gives the values of V and t for some of the years 1970 to 2010.

Year	1970	1980	1990	2000	2010
t	0	10	20	30	40
V (\$)	8000	17500	38000	83000	190000


(a) Plot lg V against t and draw a straight line graph on the grid on the next page. [2]

(b) Use your graph to estimate the values of p and q. [3]

(c) Determine the year in which Theresa's house will first be worth a million dollars. [3]

(d) Explain whether your answer to **part (c)** is likely to be reliable. [1]

Downloaded from SingaKids.com



Downloaded from SingaKids.com


8 It is given that $\log_3\left(4-x^2\right)-\log_{\sqrt{3}}(x-1)=1$.

(a) Explain clearly why $1<x<2$. [4]

(b) Hence solve $\log_3\left(4-x^2\right)-\log_{\sqrt{3}}(x-1)=1$ and show that it has only one solution. [5]

Downloaded from SingaKids.com

9 A beverage company makes closed cylindrical cans, each with volume 400cm^3. The materials for the curved surface and base of the cans cost \$0.025 per cm^2 and \$0.03 per cm^2 respectively. Assume that the cost of wasted materials is negligble.



(a) Show that the cost in dollars, C, of the materal required to make the container is $C = \frac{3}{50}\pi r^2 + \frac{20}{r}$. [3]


Downloaded from SingaKids.com

(b) Given that r can vary, find the value of r for which the cost of material is stationary. [3]

(c) Justify, why the beverage company should choose to use the value of r found in **part (b)** in producing the beverage can. [3]

Downloaded from SingaKids.com

10



The diagram shows part of the curve $y = 3\sin\frac{x}{2}$ that cuts the x – axis at $x = 0$ and $x = 2\pi$. The normal to the curve at $x = \frac{4\pi}{3}$ cuts the x-axis at Q.

(a) Find the coordinates of Q, leaving your answer in exact form. [6]

Downloaded from SingaKids.com

(b) Find the area of the shaded region bounded by the curve, the normal PQ and the coordinate axes. [4]

Downloaded from SingaKids.com

11 A particle moves in a straight line, so that, t seconds after passing a fixed point A, its velocity, v m/s, is given by $v = 10e^{-0.1t} - 5$. The particle comes to instantaneous rest at the point B.

(a) Find the value of t when the particle reaches B. Express your answer in the form of $p \ln q$, where p and q are integers. [3]

(b) Calculate the distance AB. [4]

Downloaded from SingaKids.com

(c) Find the acceleration of the particle when $t = 3$. [2]

(d) Show that the particle is again at A at some instant during the sixteenth second after first passing through A. [3]

Downloaded from SingaKids.com

BLANK PAGE

Downloaded from SingaKids.com

BLANK PAGE

Downloaded from SingaKids.com