| Name _____Solutions_____ ( ) | Class 4 _____ |
|---|---|

# CRESCENT GIRLS' SCHOOL
# SECONDARY FOUR
# PRELIMINARY EXAMINATION

**ADDITIONAL MATHEMATICS** **4049/01**
**Paper 1**

**24 August 2023**
**2 hours 15 minutes**

Candidates answer on the Question Paper.
No Additional Materials are required.

## READ THESE INSTRUCTIONS FIRST

Write your name, class and index number in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to three significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total of the marks for this paper is **90**.

***For Examiner's Use***

| Question | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Marks | 3 | 4 | 4 | 3 | 4 | 7 | 8 | 6 | 11 | 11 | 9 | 8 | 8 | 4 |

| Table of Penalties | | Qn. No. | | |
|---|---|---|---|---|
| ***Presentation*** | –1 | | | |
| ***Accuracy/ Units*** | –1 | | **Parent's/ Guardian's Signature** | **90** |

**This question paper consists of 20 printed pages.**

Downloaded from SingaKids.com

*Mathematical Formulae*

## 1. ALGEBRA

*Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

*Binomial expansion*

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + ... + \binom{n}{r}a^{n-r}b^r + ... + b^n,$$

where $n$ is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-r)...(n-r+1)}{r!}$

## 2. TRIGONOMETRY

*Identities*

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$
$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$
$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$
$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$
$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

*Formulae for $\Delta ABC$*

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2}bc \sin A$$

Downloaded from SingaKids.com

**1** Given that $k = 2\sqrt{2} - \sqrt{3}$, **without using the calculator**, express $3k - \frac{2}{k}$ in the form $\frac{a\sqrt{2} - b\sqrt{3}}{c}$, where *a*, *b* and *c* are integers. [3]

| | |
|---|---|
| $k = 2\sqrt{2} - \sqrt{3}$<br>$3k - \frac{2}{k} = 3\left(2\sqrt{2} - \sqrt{3}\right) - \frac{2}{2\sqrt{2} - \sqrt{3}}$<br>$= 6\sqrt{2} - 3\sqrt{3} - \frac{2\left(2\sqrt{2} + \sqrt{3}\right)}{8 - 3}$<br>$= 6\sqrt{2} - 3\sqrt{3} - \frac{4}{5}\sqrt{2} - \frac{2}{5}\sqrt{3}$<br>$= \frac{26\sqrt{2} - 17\sqrt{3}}{5}$<br>$= \frac{26}{5}\sqrt{2} - \frac{17}{5}\sqrt{3}$ | |

---

**2** The straight line $y = kx + 20$ intersects the curve $3y = 2kx^2 - 21$ at the points *A* and *B* whose *x*-coordinates are –3 and 4.5 respectively. Find the value of *k*. [4]

| | |
|---|---|
| $3(kx + 20) = 2kx^2 - 21$<br>$2kx^2 - 3kx - 81 = 0$<br>$2kx^2 - 3kx - 81 = x^2 - 1.5x - \frac{81}{2k} = 0$<br>–3 and 4.5 are solutions<br>$(x + 3)(x - 4.5) = 0$<br>$x^2 - 1.5x - 13.5 = 0$<br>by comparison<br>$-\frac{81}{2k} = -13.5$<br>$k = 3$ | |

Downloaded from SingaKids.com

**3** Express $-3x^2+12x-4$ in the form $a(x-h)^2+k$, where $a$, $h$ and $k$ are integers.

Hence state the coordinates of the turning point of the curve $y=-3x^2+12x-4$. [4]

| | |
|---|---|
| $-3x^2+12x-4=-3(x^2-4x)-4$<br>$-3(x^2-4x+2^2-2^2)-4=-3(x-2)^2+8$<br><br>Turning point (2, 8) | |

---

**4** Integrate $\tan^2 2x$ with respect to $x$. [3]

| | |
|---|---|
| $\int \tan^2 2x \, dx = \int \sec^2 2x - 1 \, dx$<br>$= \frac{\tan 2x}{2} - x + c$ ($c$ is a constant) | |

Downloaded from SingaKids.com

**5** Express $\frac{6x^2-5x+5}{(x-1)(x^2+2)}$ in partial fractions. [4]

| | |
|---|---|
| $\frac{6x^2-5x+5}{(x-1)(x^2+2)} = \frac{A}{x-1}+\frac{Bx+C}{x^2+2}$<br>$6x^2-5x+5 = A(x^2+2)+(Bx+C)(x-1)$<br>Sub $x=1$,<br>$6=3A$<br>$A=2$<br>Sub $x=0$,<br>$5=2A-C$<br>$C=-1$<br>Sub $x=2$,<br>$2B=8$<br>$B=4$<br>$\therefore \frac{6x^2-5x+5}{(x-1)(x^2+2)} = \frac{2}{x-1}+\frac{4x-1}{x^2+2}$ | |

Downloaded from SingaKids.com

**6** $f(x) = x^{2n} - (p+1)x^2 + p$, where $n$ and $p$ are positive integers.

**(a)** Show that $(x+1)$ is a factor of $f(x)$ for all values of $p$. [2]

| | |
|---|---|
| $f(-1) = (-1)^{2n} - (p+1)(-1)^2 + p$<br>$= 1 - (p+1)(1) + p$<br>$= 0$<br>Therefore $(x + 1)$ is a factor | |

**(b)** Given $p = 4$,

**(i)** find the value of $n$ for which $(x-2)$ is a factor, [2]

| | |
|---|---|
| $f(2) = 0$<br>$0 = 2^{2n} - (4+1)(2)^2 + 4$<br>$0 = 2^{2n} - 16$<br>$n = 2$ | |

**(ii)** hence, solve $f(x) = 0$. [3]

| | |
|---|---|
| $x^4 - 5x^2 + 4 = (x+1)(x-2)(x^2 + ax + b)$<br>by observation<br>$b = -2$<br>substitute $x = 1$,<br>$1 - 5 + 4 = 2(-1)(1 + a - 2)$<br>$a = 1$<br>$(x+1)(x-2)(x^2 + x - 2) = 0$<br>$(x+1)(x-2)(x-1)(x+2) = 0$<br>$x = -1, 1, -2, 2$ | |

Downloaded from SingaKids.com

**7** For $0 \le x \le \pi$, $f(x) = 3\sin nx$, where $n$ is a positive integer, and $g(x) = 4\cos 2x + 1$.

**(i)** Given that $\frac{\pi}{6}$ satisfies the equation $f(x) = g(x)$, show that smallest value for $n = 3$.


[2]

| | |
|---|---|
| $3\sin n\left(\frac{\pi}{6}\right) = 4\cos 2\left(\frac{\pi}{6}\right) + 1$<br>$\sin n\left(\frac{\pi}{6}\right) = 1$<br>smallest $n = 3$ | |

**(ii)** State the amplitude of $g(x)$. [1]

| | |
|---|---|
| Amplitude = 4 | |

**(iii)** Sketch, on the axes below, the graphs of $y = f(x)$ and $y = g(x)$. [4]





| | |
|---|---|
| f($x$)<br>g($x$) | |

**(iv)** State, in terms of $\pi$, the other roots of the equation $f(x) = g(x)$ for $0 \le x \le \pi$. [1]

| | |
|---|---|
| $\frac{\pi}{2}, \frac{5\pi}{6}$ | |

Downloaded from SingaKids.com

**8** A piece of wire, 100 cm in length, is bent to form the figure as shown.




Given that angle $ABC$ = angle $EFG = 60°$, angle $CDE$ = angle $GHA = 90°$,

$AB = BC = EF = FG = x$ cm and $CD = DE = GH = HA = y$ cm.

**(a)** Show that the area of the figure, $P$ cm$^2$, is given by

$$P = \left(\frac{\sqrt{3}}{2} + 1 - \sqrt{2}\right)x^2 + (25\sqrt{2} - 50)x + 625. \qquad [4]$$

$4(x + y) = 100 \Rightarrow$

$x + y = 25$

$y = 25 - x$

$CE = \sqrt{y^2 + y^2} = \sqrt{2}y$

$P =$ $2 \times$ Area of $\Delta ABC +$
$2 \times$ Area of $\Delta CDE +$
Area of rectangle $ACEG$

$= x^2 \sin 60° + y^2 + x(\sqrt{2}y)$

$= \frac{\sqrt{3}}{2}x^2 + (25 - x)^2 + \sqrt{2}x(25 - x)$

$= \frac{\sqrt{3}}{2}x^2 + 625 - 50x + x^2 + 25\sqrt{2}x - \sqrt{2}x^2$

$= \left(\frac{\sqrt{3}}{2} + 1 - \sqrt{2}\right)x^2 + (25\sqrt{2} - 50)x + 625$


Downloaded from SingaKids.com

**(b)** Find the value of $x$ for which $P$ has a stationary value. [2]

| | |
|---|---|
| $\frac{\mathrm{d}P}{\mathrm{d}x} = 2\left(\frac{\sqrt{3}}{2}+1-\sqrt{2}\right)x + 25\sqrt{2} - 50$<br>$\frac{dP}{dx} = 0$<br>$2\left(\frac{\sqrt{3}}{2}+1-\sqrt{2}\right)x = 50 - 25\sqrt{2}$<br>$x = \frac{50-25\sqrt{2}}{2\left(\frac{\sqrt{3}}{2}+1-\sqrt{2}\right)} = 16.2 \text{ (3 s.f.)}$ | |

Downloaded from SingaKids.com

**9**




The diagram shows a kite $ABCD$ with $AB = AD$ and $CB = CD$.

The diagonals intersect at $M$. The point $A$ lies on the $y$-axis, the point $B$ is (–9, 2) and the equation of $AC$ is $2x + y = 9$.

**(i)** State the coordinates of $A$. [1]

| | |
|---|---|
| $A$ (0, 9) | |

**(ii)** Find the equation of $BD$. [2]

| | |
|---|---|
| Gradient of $AC = -2$<br>Gradient of $BD = \frac{1}{2}$<br>Equation $BD$ :<br>$y - 2 = \frac{1}{2}(x - (-9))$<br>$y = \frac{1}{2}x + 6\frac{1}{2}$ | |

Downloaded from SingaKids.com

**(iii)** Find the coordinates of $M$ and of $D$. [4]

| | |
|---|---|
| By solving simultaneous equations<br>$y = \frac{1}{2}x + 6\frac{1}{2}$ and $2x + y = 9$<br><br>$9 - 2x = \frac{1}{2}x + 6\frac{1}{2}$<br>$x = 1$<br>$y = 7$<br>$M\ (1, 7)$<br><br>$M$ is mid-point of $BD$<br>$\frac{-9 + x_D}{2} = 1, \frac{2 + y_D}{2} = 7$<br>$D\ (11, 12)$ | |

Given further that the area of the triangle $ABD$ is $\frac{1}{4}$ of the area of the triangle $CBD$, find

**(iv)** the coordinates of $C$, [2]

| | |
|---|---|
| $MC = 4\ AM$<br>$x_C = 1 + 4(1) = 5, y_C = 7 - 4(2) = -1$<br>$C(5, -1)$ | |

**(v)** the area of the kite $ABCD$. [2]

| | |
|---|---|
| $\frac{1}{2}\begin{vmatrix} 0 & -9 & 5 & 11 & 0 \\ 9 & 2 & -1 & 12 & 9 \end{vmatrix}$<br><br>$= 125$ units$^2$ | |

Downloaded from SingaKids.com

**10** **(a)** Find in radians, the principal value of $\tan^{-1}\left(-\frac{1}{\sqrt{3}}\right)$. [2]

| | |
|---|---|
| Principal value of $\tan^{-1}\left(-\frac{1}{\sqrt{3}}\right)$ is in the 4th quadrant<br><br>Since $\tan\left(\frac{\pi}{3}\right)=\frac{1}{\sqrt{3}}$,<br><br>The principal value is $-\frac{\pi}{6}$ or –0.524. | |

**(b)** Given $\sin 3\theta + \sin\theta = 2\sin 2\theta\cos\theta$. Prove that

**(i)** $\cos 3\theta + \cos\theta = 2\cos 2\theta\cos\theta$, [3]

| | |
|---|---|
| $\cos 3\theta + \cos\theta = \cos 2\theta\cos\theta - \sin 2\theta\sin\theta + \cos\theta$<br>$= \left(1-2\sin^2\theta\right)\cos\theta - 2\sin^2\theta\cos\theta + \cos\theta$<br>$= 2\cos\theta - 4\sin^2\theta\cos\theta$<br>$= 2\cos\theta(1-2\sin^2\theta)$<br>$= 2\cos 2\theta\cos\theta$ | |

**(ii)** $\frac{\sin\theta - \sin 2\theta + \sin 3\theta}{\cos\theta - \cos 2\theta + \cos 3\theta} = \tan 2\theta$. [3]

| | |
|---|---|
| $\frac{\sin\theta - \sin 2\theta + \sin 3\theta}{\cos\theta - \cos 2\theta + \cos 3\theta} = \frac{\sin 3\theta + \sin\theta - \sin 2\theta}{\cos 3\theta + \cos\theta - \cos 2\theta}$<br>$= \frac{2\sin 2\theta\cos\theta - \sin 2\theta}{2\cos 2\theta\cos\theta - \cos 2\theta}$<br>$= \frac{\sin 2\theta\left(2\cos\theta - 1\right)}{\cos 2\theta\left(2\cos\theta - 1\right)}$<br>$= \tan 2\theta$ | |


Downloaded from SingaKids.com

**(c)** Hence solve the equation $\dfrac{\cos\theta - \cos 2\theta + \cos 3\theta}{\sin\theta - \sin 2\theta + \sin 3\theta} = -\dfrac{1}{2}$ for $0 \le \theta \le \pi$. [3]

| | |
|---|---|
| $\dfrac{\sin\theta - \sin 2\theta + \sin 3\theta}{\cos\theta - \cos 2\theta + \cos 3\theta} = -2$<br>$\tan 2\theta = -2$<br>$\alpha = 1.1071$<br>$2\theta = 2.0345..., 5.1761...$<br>$\theta = 1.02, 2.59$ (3 s.f.) | |

Downloaded from SingaKids.com

**11** In the diagram, not to scale, $BC$ and $CE$ are diameters of the circles, $S_1$ and $S_2$, respectively. $CE$ is tangent to $S_1$ at $C$, $CF$ and $BD$ meet at $G$, and $G$ lies on the circumference of $S_1$. $F$ lies on the circumference of $S_2$. $CB$ produced and $EF$ produced meet at $A$.




Show that

**(i)** triangles $CBG$ and $DCG$ are similar, [3]



| | |
|---|---|
| $\angle CGB = 90°$ (Angle in a semi-circle)<br>$\angle CGD = 90°$ (adjacent angles on a straight line)<br>$\therefore \angle CGB = \angle CGD$<br><br>$\angle CBG = \angle DCG$<br>(Alternate Segment Theorem/ Tangent-Chord Theorem)<br><br>Therefore, by AA pty, triangles $CBG$ and $DCG$ are similar | |

Downloaded from SingaKids.com

**(ii)** lines $BGD$ and $AFE$ are parallel, [2]

| | |
|---|---|
| $\angle CFE = 90°$ (Angle in a semi-circle)<br>Since $\angle CGD = 90°$ as well,<br>lines $BGD$ and $AFE$ are parallel by means of corresponding angles | |

**(iii)** $CE^2 = AE \times EF$. [4]

| | |
|---|---|
| In triangles $CEF$ and $AEC$,<br>$\angle EFC = 90°$ (Angle in a semicircle)<br>$\angle ECA = 90°$ (Tangent perpendicular to radius)<br>Therefore $\angle EFC = \angle ECA$.<br><br>$\angle CEF = \angle AEC$ (Common angle)<br>Triangles $CEF$ and $AEC$ are similar (AA).<br><br>Comparing triangles $AEC$ and $CEF$,<br>$\frac{CE}{AE} = \frac{FE}{CE}$<br><br>$CE \times CE = FE \times AE$<br>$CE^2 = AE \times EF$ | <br> |

Downloaded from SingaKids.com

**12** Solve the following equations:

**(a)** $\log_3(2x-1) - \log_{\sqrt{3}} 3 = \log_2 4$ [4]

$$\log_3(2x-1) - \log_{\sqrt{3}} 3 = \log_2 4$$

$$\log_3(2x-1) - \frac{\log_3 3}{\log_3 \sqrt{3}} = \log_2 4$$

$$\log_3(2x-1) - \frac{1}{\log_3 3^{\frac{1}{2}}} = \log_2 2^2$$

$$\log_3(2x-1) - \frac{1}{\frac{1}{2}\log_3 3} = 2\log_2 2$$

$$\log_3(2x-1) - 2 = 2$$

$$\log_3(2x-1) = 4$$

$$2x-1 = 3^4$$

$$x = \frac{3^4+1}{2} = 41$$

Downloaded from SingaKids.com

**(b)** $\dfrac{1}{\sqrt{25^{2x+1}}}+\dfrac{10}{5^{x+2}}=3$. [4]

$$\frac{1}{\sqrt{25^{2x+1}}}+\frac{10}{5^{x+2}}=3$$

$$\left(25^{2x+1}\right)^{-\frac{1}{2}}+10\left(5^{-x-2}\right)=3$$

$$\left(5^{2}\right)^{-x-\frac{1}{2}}+10\left(5^{-x}\right)\left(5^{-2}\right)=3$$

$$5^{-2x-1}+\frac{10}{25}\left(5^{-x}\right)=3$$

$$\left(5^{-2x}\right)\left(5^{-1}\right)+\frac{2}{5}\left(5^{-x}\right)=3$$

$$5^{-2x}+2\left(5^{-x}\right)=15$$

$$\left(5^{-x}\right)^{2}+2\left(5^{-x}\right)=15$$

Let $u = 5^x$

$$15u^{2}-2u-1=0$$

$$(3u-1)(5u+1)=0$$


$$u=\frac{1}{3} \quad \text{or} \quad u=-\frac{1}{5}$$

$$5^{x}=\frac{1}{3} \quad \text{or} \quad 5^{x}=-\frac{1}{5}\text{(N.A.)}$$

$$x=\frac{\ln\frac{1}{3}}{\ln 5}=-0.683$$

Downloaded from SingaKids.com

**13** The sketch shows the graphs of the curve, $y = \frac{1}{x-1} - 3$ , the lines $3y = 4x - 9$ and $y = 3$. The curve and the line $3y = 4x - 9$ intersect at $P$. The curve cuts the $x$-axis at $R\left(\frac{4}{3}, 0\right)$. The line $3y = 4x - 9$ cuts the $x$-axis at $Q\left(2\frac{1}{4}, 0\right)$.




The region $A$ is bounded by the curve, $y = \frac{1}{x-1} - 3$, the line $y = 3$ and the $y$-axis.

The region $B$ is bounded by the curve, the line $3y = 4x - 9$, and the $x$-axis.

**(i)** Verify that the coordinates of $P$ are $\left(\frac{3}{2}, -1\right)$. [2]

| | |
|---|---|
| Solve simultaneous,<br>$3\left(\frac{1}{x-1} - 3\right) = 4x - 9$<br>$\frac{3}{x-1} = 4x$<br>$4x^2 - 4x - 3 = 0$<br>$(2x-3)(2x+1) = 0$<br>$2x - 3 = 0$ or $2x + 1 = 0$<br>$x = \frac{3}{2}$ or $x = -\frac{1}{2}$ (N.A.)<br>$x = \frac{3}{2}$, $3y = 4\left(\frac{3}{2}\right) - 9$, $y = -1$<br>$P\left(\frac{3}{2}, -1\right)$ | |

Downloaded from SingaKids.com

**(ii)** Find the area of $A$ and of $B$. [6]

$$\text{Area } A = \int_0^3 \left( \frac{1}{y+3} + 1 \right) dy$$

$$= \left[ \ln(y+3) + y \right]_0^3$$

$$= \left( \ln(3+3) + 3 \right) - \left( \ln(0+3) + 0 \right)$$

$$= \ln 6 - \ln 3 + 3$$

$$= \ln 2 + 3$$

$$= 3.69 \text{ units}^2 \text{ (3 s.f.)}$$

$$\text{Area } B = \left| \int_{\frac{4}{3}}^{\frac{3}{2}} \left( \frac{1}{x-1} - 3 \right) dx \right| + \frac{1}{2} \times \left( 2\frac{1}{4} - \frac{3}{2} \right) \times 1$$

$$= \left| \left[ \ln(x-1) - 3x \right]_{\frac{4}{3}}^{\frac{3}{2}} \right| + \frac{3}{8}$$

$$= \left| \left( \ln\left(\frac{3}{2} - 1\right) - 3\left(\frac{3}{2}\right) \right) - \left( \ln\left(\frac{4}{3} - 1\right) - 3\left(\frac{4}{3}\right) \right) \right| + \frac{3}{8}$$

$$= \left| \left( \ln\left(\frac{1}{2}\right) - \left(\frac{9}{2}\right) \right) - \left( \ln\left(\frac{1}{3}\right) - 4 \right) \right| + \frac{3}{8}$$


$$= \left| \left( \ln\left(\frac{3}{2}\right) - \frac{1}{2} \right) \right| + \frac{3}{8}$$

$$= 0.470 \text{ units}^2 \text{ (3 s.f.)}$$

Downloaded from SingaKids.com

**14** A vessel is in the shape of a right circular cone.
The radius of cone is 8 cm and the height is 32 cm.
Water is poured into the vessel at a rate of 10 $cm^3/s$.

Calculate the rate at which the water level is rising when the vessel is $\frac{1}{8}$ full. [4]




| | |
|---|---|
| Using similar triangles<br>$\frac{r}{h}=\frac{8}{32}$<br>$r=\frac{1}{4}h$<br>$V=\frac{1}{3}\pi r^2 h$<br>$=\frac{1}{3}\pi\left(\frac{h}{4}\right)^2 h$<br>$V=\frac{1}{48}\pi h^3$<br>$\frac{dV}{dh}=\frac{1}{16}\pi h^2$<br>$\frac{v}{V}:\left(\frac{h}{H}\right)^3$<br>$\frac{1}{8}:\left(\frac{h}{32}\right)^3$<br>$h=16$<br>$\frac{dV}{dt}=\frac{dV}{dh}\times\frac{dh}{dt}$<br>$10=\frac{1}{16}\pi(16)^2\times\frac{dh}{dt}$<br>$\frac{dh}{dt}=10\div 16\pi$<br>$\frac{dh}{dt}=\frac{5}{8\pi}$ cm/ s or 0.199 cm/ s | |

**END OF PAPER**

Downloaded from SingaKids.com