CEDAR GIRLS' SECONDARY SCHOOL
Preliminary Examination
Secondary Four

CANDIDATE NAME										
CLASS	4			INDEX NUMBER						
CENTRE/ INDEX NO					/					

ADDITIONAL MATHEMATICS

4049/02

Paper 2

26 August 2024

2 hours 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 90.

For Examiner's Use
90

This document consists of **21** printed pages and 1 blank page.

[Turn over

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A\cos B \pm \cos A\sin B$$

$$\cos(A \pm B) = \cos A\cos B \mp \sin A\sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A\tan B}$$

$$\sin 2A = 2\sin A\cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

Answer **all** the questions.

1 **(a)** The curve $y = mx^2 - 12x + 2(x^2 + m - 1)$ lies entirely below the x-axis for all real values of x. Find the largest integer value of m. [5]

(b) Show that the roots of the equation $2x^2 - 3(1-x) = -p$ are real if $p \leq 4\frac{1}{8}$. [4]

Downloaded from SingaKids.com

2 Let $\mathrm{f}(x) = \dfrac{3-3x^2}{(2x+1)(x+2)^2}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

Downloaded from SingaKids.com

(b) Hence find the value of $\int_0^4 \mathrm{f}(x)\,\mathrm{d}x$, giving your answer in the form $a + b\ln c$, where a, b and c are integers. [5]

Downloaded from SingaKids.com

3 **(a)** **(i)** Prove $\operatorname{cosec} 2\theta - \cot 2\theta = \tan\theta$. [3]

(ii) Hence, solve $\operatorname{cosec} 4\theta - \cot 4\theta = -\sqrt{3}$ for $0 < \theta < \pi$. [2]

Downloaded from SingaKids.com

(b) The angles A and B are such that

$$\sin(A+45°) = \left(2\sqrt{2}\right)\cos A \quad \text{and} \quad 4\sec^2 B + 5 = 12\tan B.$$

Without using a calculator, find the exact value of $\tan(A-B)$. [5]

Downloaded from SingaKids.com

4 The fifth term in the expansion of $\left(px - \frac{q}{x} \right)^n$, where p and q are positive numbers, is independent of x.

(a) Show that $n = 8$. [2]

(b) Hence, explain why the fifth term is a positive constant. [1]

Downloaded from SingaKids.com

It is given that $p = 3$ and $q = 1$.


(c) Find the term independent of x in $\left(2 + \frac{1}{x^2}\right)\left(px - \frac{q}{x}\right)^n$ for $n = 8$. [4]

Downloaded from SingaKids.com

5 The diagram shows a line DE which cuts the y-axis at P and a line through F meets the y-axis at Q such that angle FPQ = angle FQP.

The coordinates of D and E are $(-4, 0)$ and $(10, 7)$ respectively.

F is a point on the line DE such that $DF : FE = 4 : 3$.



(a) Find the coordinates of F. [3]


Downloaded from SingaKids.com

(b) Find the equation of the straight line FQ. [2]

(c) Find the area of the triangle QFE. [3]

Downloaded from SingaKids.com

6



The diagram shows a figure *PQRST* which consists of a rectangle *PQST* and an isosceles triangle *QRS*.
It is given that $PQ = 4$ cm and $QR = 8$ cm.

(a) Give angle SQR = angle $QSR = x$ radians and the area of *PQRST* is given by A cm^2, show that $A = 64\cos x + 32\sin 2x$. [4]

Downloaded from SingaKids.com

(b) Find the value of x for which A has a stationary value. [3]

(c) Hence find the exact stationary value of A and determine whether it is a maximum or a minimum. [2]

Downloaded from SingaKids.com

7 It is given that $f(x) = x^3 + ax^2 - 5x + b$, where a and b are constants, has a factor of $x+1$ and leaves a remainder of -24 when divided by $(x+3)$.

(a) Show that $a = -1$ and $b = -3$. [4]

Downloaded from SingaKids.com


(b) Hence factorise f (x) completely and write down the solutions for $f(x) = 0$. [3]

Downloaded from SingaKids.com

From part **(a)**, it is known that $f(x) = x^3 - x^2 - 5x - 3$.

The diagram below shows part of the curve $y = f(x)$ and a line that cuts the curve at B and C. B has coordinates $(1, -6)$.

A and C are points on the curve and the x-axis.



(c) Find the area of the shaded region. [5]

Downloaded from SingaKids.com

(d) Show that the equation of the tangent at the minimum point of the curve is $y = -\frac{256}{27}$. [3]

Downloaded from SingaKids.com

8 A circle has a diameter AB. The point A has coordinates $(1, -6)$ and the equation of the tangent to the circle at B is $3x + 4y = k$.

(a) Show that the equation of the normal to the circle at the point A is $4x - 3y = 22$. [3]

It is also given also that the line $x = -1$ touches the circle at the point $(-1, -2)$.

(b) Find the coordinates of the centre and the radius of the circle. [4]

Downloaded from SingaKids.com

Continuation of working space for question **8 (b).**

(c) Find the value of k. [3]

Downloaded from SingaKids.com

9 Mr Chan was driving a car along a straight road. He was 35 m away from the stop line when he applied his brakes near a traffic light. His acceleration, a m/s^2, after applying the brakes was given by $a = -7.5e^{-\frac{t}{2}}$, where t is the time in seconds after he applied the brakes.

(a) Explain mathematically why $a < 0$ for all $t \geq 0$ and the significance of $a < 0$. [2]

(b) Mr Chan's car was travelling at 14 m/s just before he applied his brakes. Express the velocity of his car, v m/s, in terms of t. [3]

Downloaded from SingaKids.com

(c) Hence find the time taken for his car to come to a complete stop. [2]

(d) Obtain an expression, in terms of t, for the displacement of Mr Chan's car from the point he applied the brakes [3]

(e) Determine if Mr Chan's car was able to come to a complete stop before reaching the stop line. Explain your answer. [2]

End of Paper

Downloaded from SingaKids.com

BLANK PAGE

Downloaded from SingaKids.com