General Certificate of Education Ordinary Level
JUYING SECONDARY SCHOOL, SINGAPORE
Secondary Four Express/Five Normal Academic
Preliminary Examination

CANDIDATE NAME									
CENTRE NUMBER	S					INDEX NUMBER				

ADDITIONAL MATHEMATICS

4049/01

Paper 1

22 August 2024

2 hours 15 minutes

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions.
The number of marks is given in brackets [ ] at the end of each question or part question.

If working is needed in any question it must be shown with the answer.
Omission of essential working will result in loss of marks.
The total number of marks for this paper is 90.

The use of an approved scientific calculator is expected, where appropriate.
If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.
For π, use either your calculator value or 3.142.

This document consists of **18** printed pages.

Set by: Mr Albert Lui
Vetted by: Mdm Norhafiani Bte Abdul Majid

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \ldots + \binom{n}{r} a^{n-r} b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

Downloaded from SingaKids.com

Answer **ALL** the questions

1 **(a)** The function f is defined, for all values of x, by $f(x) = (2x - x^2)e^x$.


Find the range of values of x such that $f(x)$ is a decreasing function. [4]

(b) The gradient function of the curve is $2(p + 1)x + 2$, where p is a constant.

Given that the tangent to the curve at $(2\ , -2)$ is parallel to $y + 2x - 5 = 0$, find the value of p. [3]

Downloaded from SingaKids.com

2 The diagram shows a chocolate bar in the form of a triangular prism and the cross-section of the chocolate bar is an isosceles triangle with $AB = AC$.
$MC = \left(\sqrt{2} + \frac{1}{2}\right)$ cm and $\angle ACB = 45°$.




(a) Find the exact length of AC. [3]

Downloaded from SingaKids.com

(b) Given that the volume of the chocolate bar is $\left(25 + 22\sqrt{2}\right)\text{cm}^3$, find the length of AD in the form $\left(a + b\sqrt{2}\right)$ cm, where a and b are integers. [4]

Downloaded from SingaKids.com

3



The diagram shows a circle, centre *O*, with diameter *AB*. The points *D* and *F* lie on the circle. The point *E* is such that *EB* and *EF* are tangents to the circle.

(a) Given that the points C and D are midpoints of BE and AE respectively, prove that angle $DCE = 90°$. [3]

(b) Given that triangle *BEF* is equilateral, prove that$\angle BEF = \angle BAF$. [2]

Downloaded from SingaKids.com

4 **(a)** Find the remainder when $6x^3 - 13x^2 + 17x - 6$ is divided by $2x - 1$. [2]

(b) Show that there is only one real root of the equation

$6x^3 - 13x^2 + 17x - 6 = 0$. [3]

Downloaded from SingaKids.com

5 Solve the following equations.

(a) $5^x - 5^{\frac{x}{2}+1} = 6,$ [3]

(b) $2\lg(x-3) - \lg(x+7) = \frac{1}{\log_{100} 10}.$ [4]

Downloaded from SingaKids.com

6 **(a)** State the values between which the principal value of $\sin^{-1} x$ must lie. [1]

(b) Find the principal value of $\tan^{-1} 1$ in radian in exact form. [1]

7 Given that $\cot\theta = -\frac{3}{4}$ and that $\tan\theta$ and $\cos\theta$ have opposite signs, without evaluating θ, find the exact values of each of the following.

(a) $\cos(-\theta)$, [2]

(b) $\sin 2\theta$ [2]


Downloaded from SingaKids.com

8. The approximate mean distance x (in millions of kilometres) from the centre of the Sun and the period of the orbit T (in Earth years) are recorded in the table.

	Mercury	**Venus**	**Mars**	**Uranus**
x	58	108	228	2871
T	0.24	0.62	1.88	84.11

It is believed that the planets orbiting around the Sun obey a law of the form $T = kx^n$, where k and n are constants.

(a) Express the equation in a form suitable for drawing a straight line graph and draw the graph using appropriate scaling on both axes. [4]



Downloaded from SingaKids.com

(b) Use your graph to estimate the value of k and of n, to two significant figures. [3]

(c) Using the graph, find the orbital period of the Earth, if the distance between the Earth and the Sun is about $149.6 \times 10^6 \, km$. Give your answer correct to the nearest integer. [2]

(d) If the orbital period of the Jupiter is 11.86 Earth years, estimate the distance of the Jupiter from the Sun in km using your graph. [2]

Downloaded from SingaKids.com

9. **(a)** Express $\frac{2x^3+2x^2-7x+4}{x(x-1)^2}$ in partial fractions. [5]

(b) Hence evaluate $\int_2^4 \frac{4x^3+4x^2-14x+8}{3x(x-1)^2}\,dx$. [4]

Downloaded from SingaKids.com

10. **(a)** Find the range of values of k for which the line $2x - y = 5$ intersects the curve $xy = kx - 2$ at two distinct points. [4]

(b) Find the smallest integer value of h for which the graph $y = 2x^2 - 4x + h$ lies entirely above the line $y = 3$ for all values of x. [3]

Downloaded from SingaKids.com

11. **(a)** Prove the identity $\frac{1+\cos\theta}{\sin\theta} + \frac{\sin\theta}{1+\cos\theta} = 2\text{cosec}\,\theta$. [4]

(b) Hence, find all the angles from $0° \leq \theta \leq 360°$ which satisfy the equation $\frac{1+\cos 2\theta}{\sin 2\theta} + \frac{\sin 2\theta}{1+\cos 2\theta} = \tan 75°$. [3]

Downloaded from SingaKids.com

12. Find the derivatives of each of the following, simplifying your answer.


(a) $y = 3\left(1 - \frac{x}{3}\right)^4$ [1]

(b) $f(x) = (2 - 3x)\left(\sqrt{1 - 4x}\right)$ [3]

(c) $\frac{dy}{dx} = \frac{2(3x-2)}{4+x}$ [2]

Downloaded from SingaKids.com

13.



The diagram shows a glass window $ABCDEF$, consisting of a rectangle $ABEF$ of height $3x$ cm and width y cm and a trapezium $BCDE$ in which $CD = x$ cm and $BC = 2y$ cm. ABC is a straight line and $DE = 10\sqrt{2}$ cm.

Given that x can vary,

(a) show that the area of the glass window $S = 7x\left(\sqrt{50 - x^2}\right)$, [3]

Downloaded from SingaKids.com

(b) find the value of x for which S has a stationary value and determine whether this value of S is a maximum or a minimum. [5]

Downloaded from SingaKids.com

14. It is given that $f(x)$ is such that $f'(x) = \cos 4x - \sin 2x$. Given also that $f\left(\frac{\pi}{2}\right) = \frac{1}{4}$, show that $f''(x) + 4f(x) = 3 - 3\sin 4x$. [5]

Downloaded from SingaKids.com