KENT RIDGE SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2024

ADDITIONAL MATHEMATICS **4049/02**
PAPER 2

SECONDARY 4 EXPRESS/ 5 NORMAL ACADEMIC

Monday 26 August 2024 **2 hour 15 minutes**

KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE
KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE
KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE
KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE SECONDARY SCHOOL KENT RIDGE

Name: _____ ( ) **Class: Sec** _____

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your name, index number and class in the spaces at the top of this page.

Do not open this question paper until you are told to do so.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue, correction fluid or correction tape.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 90.

For Examiner's Use	
Total	90

This Question Paper consists of 19 printed pages, including this page.

[Turn over

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}ab\sin C$$

Downloaded from SingaKids.com

1 Express $\frac{4}{(x^2+1)(x+1)}$ in partial fractions. [5]

Downloaded from SingaKids.com


2 (a) Radiation intensity, R, varies inversely with the square of d, the distance from the source of radiation such that $R = \frac{k}{d^2}$, where k is a constant.

Values of R for different values of d have been collected and tabulated.
Explain how a straight-lined graph can be drawn to determine the formula connecting R and d. [4]

(b) The number of particles present in a room, t minutes after turning on the air filter is y. When corresponding values of lg y and t are plotted on a lg y against t axes, the points form a straight line that passes through (0,3) and (10,1.5) as drawn on the axes on the next page.

(i) Find y in terms of t. [4]


Downloaded from SingaKids.com



(ii) Use the graph to estimate the time taken for the number of particles in the room to be halved. [3]

Downloaded from SingaKids.com

3 (a) The diagram below shows the graph of the curve $y = \cos x + \frac{x}{2}$ for $x \geq 0$ radians.
The tangent to the curve when $x = 0$ at C, is drawn to D which is vertically above point B, the minimum point of the curve.
Points A and B are the first two stationary points of the curve.
Find x_b, the x coordinates of point B.
You do not need to show that it is a minimum point. [4]



(b) (i) Find the equation of CD. [2]

Downloaded from SingaKids.com

(ii) Find the area shaded that is bounded by the tangent to the curve $y = \cos x + \frac{x}{2}$ at $x = 0$, the curve and the line $x = x_b$. [5]

Downloaded from SingaKids.com

4 (a) $2y = 16x + k$ is a tangent to the curve $y = \frac{1}{2x} + 2kx$. Find the value of constant k. [4]

Downloaded from SingaKids.com

(b) Find the range of values of a such that $ax^2 + \sqrt{8}x + (a-1) < 0$ for all values of x. [4]

Downloaded from SingaKids.com

5 Given $y = e^{2x} \sin 3x$.

(a) Find $\frac{dy}{dx}$. [2]

(b) Find $\frac{d^2y}{dx^2}$. [2]

Downloaded from SingaKids.com

(c) Given that $\frac{dy}{dx} + \frac{d^2y}{dx^2} + ay = be^{2x}\cos 3x$, form 2 equations involving a and b and use them to find the value of a and of b. [4]

Downloaded from SingaKids.com

6 (a) Show that $\frac{d}{dx}\left(\frac{x-2}{\sqrt{3x+1}}\right) = \frac{3x+8}{2\sqrt{(3x+1)^3}}$. [4]


(b) Hence evaluate $\int_0^5 \frac{3x+7}{2\sqrt{(3x+1)^3}}\,dx$. [5]

Downloaded from SingaKids.com

For continuation of working for question 6 part (b)

Downloaded from SingaKids.com

7 AF is a tangent to the circle $ABCD$ at A.
E is on AF such that EDC is parallel to AB.



(a) Prove that triangle ABD and triangle DAE are similar. [3]

(b) Show that if triangle BCD and triangle DAB are similar, BD must be the diameter of the circle. [3]

Downloaded from SingaKids.com

8 (a) Solve $3(3^{x+1}) = 10 - 3^{-x}$. [4]


(b) Given $\log_{100} x + \lg y = 3$, express y in terms of x. [4]

Downloaded from SingaKids.com

9 The chord AB of a circle C has equation $y = -\frac{1}{2}x + 10$, where the x coordinate of A is smaller than the x coordinate of B.

The circle C has equation $(x - 2.5)^2 + (y - 5)^2 = \frac{365}{4}$ with centre E.

(a) Find the coordinates of A. [4]



Downloaded from SingaKids.com

(b) State the centre of circle C, and use it to show that the perpendicular bisector of AB passes through the origin. [4]

(c) The chord AB is extended to cut the x-axis at point D. Show that the mid-point of AD lies inside circle C. [4]

Downloaded from SingaKids.com

10 A particle starts moving in a straight line when it is 6 metres from a fixed point O, such that its velocity, t seconds after the start of the motion is given by $v = 4e^{-2t} + t - 3$ m/s.

(a) Find the initial velocity of the particle. [2]

(b) Show that the minimum velocity is negative, and it happens when $t = \frac{1}{2}\ln 8$. [4]

(c) Using your answer from part (a) and part (b), explain if the particle changes its direction of motion. [2]

Downloaded from SingaKids.com

(d) Find the displacement of the particle from O, 2 seconds after the start of the motion. [4]

End of Paper

Downloaded from SingaKids.com

Downloaded from SingaKids.com

Kent Ridge Secondary School
Secondary 4 Express/5 Normal Academic Preliminary Examination 2024
Add Math Prelim 2024 P2 Mark scheme

Qn	Solutions	Marks	
1a	$\frac{4}{(x^2+1)(x+1)} = \frac{Ax+B}{x^2+1} + \frac{C}{x+1}$ $\frac{4}{(x^2+1)(x+1)} = \frac{(Ax+B)(x+1)}{(x^2+1)(x+1)} + \frac{C(x^2+1)}{(x^2+1)(x+1)}$ $4 = (Ax+B)(x+1) + C(x^2+1)$ Sub x = -1 $4 = C((-1)^2+1)$ $C = 2$ Sub x = 0 $4 = (B)(1) + C(1)$ $B = 2$ Compare coef of x^2: $A + C = 0$ $A = -2$ $\frac{4}{(x^2+1)(x+1)} = \frac{2-2x}{x^2+1} + \frac{2}{x+1}$	A1 M1 M1 M1 A1	
2a	Plot points of **corresponding values of R and** $\frac{1}{d^2}$ Draw best fit line through points and the **origin** Find **gradient** of the line gives the **value of *k***	B1 B1 B1 B1	
2bi	Gradient of line $= \frac{1.5}{-10} = -0.15$ Lg y intercept = 3 $\lg y = -0.15t + 3$ $y = 10^{-0.15t+3}$	M1 B1 M1 A1	
2bii	Initial number of particles : $\lg y = 3$ $y = 10^3 = 1000$ Find the point on the straight line when $\lg y = \lg 500 = 2.69$ The time taken is the t value of the point	 M1 M1 B1 – their t value (± 0.4)	
3a	$\frac{dy}{dx} = -\sin x + \frac{1}{2}$ $-\sin x + \frac{1}{2} = 0$ $\sin x = \frac{1}{2}$ $\alpha = \frac{\pi}{6}$ $x = \frac{\pi}{6}, \frac{5\pi}{6}$ $x = \frac{5\pi}{6}$	M1 A1 M1 A1	
3bi	Gradient of tangent at x = 0 $\frac{dy}{dx} = -\sin 0 + \frac{1}{2} = \frac{1}{2}$ $y = \cos 0 + 0 = 1$ Equation of tangent $y = \frac{1}{2}x + 1$	M1 M1	

Downloaded from SingaKids.com

Qn	Solutions	Marks	
3bii	$Area = \int_0^{\frac{5\pi}{6}} \frac{1}{2}x + 1 - \cos x - \frac{x}{2}\, dx$	M1 – mtd to find area trap under tangent M1 – definite integral of curve from 0 to x_b	
	$= [x - \sin x]_0^{\frac{5\pi}{6}}$	A1 – correct expr of their integrals	
	$= \frac{5\pi}{6} - \sin\left(\frac{5\pi}{6}\right)$	A1 – correct sub of limits	
	$= \frac{5\pi}{6} - \frac{1}{2} = 2.12\ (3\ s.f.)$	A1	
4a	$\frac{k}{2} + 8x = \frac{1}{2x} + 2kx$	M1	
	$kx + 16x^2 = 1 + 4kx^2$ $(4k - 16)x^2 - kx + 1 = 0$	M1	
	$k^2 - 4(4k - 16)(1) = 0$ $k^2 - 16k + 64 = 0$ $k = 8$	M1 A1	
4b	Discriminant: $\sqrt{8}^2 - 4a(a-1) < 0$ $8 - 4a^2 + 4a < 0$ $a^2 - a - 2 > 0$ $(a-2)(a+1) > 0$ Since $a < 0$, $a < -1$	M1 – expr for D M1 – condition for D B1 A1	
5a	$y = e^{2x}\sin 3x$ $\frac{dy}{dx} = 2e^{2x}\sin 3x + 3e^{2x}\cos 3x$	M1 either term seen A1 use of product rule and final ans	
5b	$\frac{d^2y}{dx^2} = 2(2e^{2x}\sin 3x + 3e^{2x}\cos 3x) + 3(2e^{2x}\cos 3x - 3e^{2x}\sin 3x$	M1 use of at one correct product rule of their dy/dx	
5c	$\frac{d^2y}{dx^2} = -5e^{2x}\sin 3x + 12e^{2x}\cos 3x$	A1	
	$2e^{2x}\sin 3x + 3e^{2x}\cos 3x - 5e^{2x}\sin 3x + 12e^{2x}\cos 3x + ae^{2x}\sin 3x = be^{2x}\cos 3x$	M1	
	$2 - 5 + a = 0$	M1	
	$3 + 12 = b$	M1	
	$a = 3, b = 15$	A1	

Downloaded from SingaKids.com

Qn	Solutions	Marks	
6	$\frac{d}{dx}\left(\frac{x-2}{\sqrt{3x+1}}\right)=\frac{\sqrt{3x+1}-\frac{3(x-2)}{2\sqrt{3x+1}}}{3x+1}$ $=\frac{\frac{2(3x+1)}{2\sqrt{3x+1}}-\frac{3(x-2)}{2\sqrt{3x+1}}}{3x+1}$ $=\frac{3x+8}{2\sqrt{3x+1}(3x+1)}$ $=\frac{3x+8}{2\sqrt{(3x+1)^3}}$	M1 – quotient rule seen with positive sq root or product seen with negative sq root M1 – simplify with common denominator or taking out common factor M1 – all factors in denominator collected B1	
6b	$\int_{x_1}^{x_2}\frac{3x+8}{2\sqrt{(3x+1)^3}}dx=\left[\frac{x-2}{\sqrt{3x+1}}\right]_{x_1}^{x_2}$ $\int_{x_1}^{x_2}\frac{3x+7}{2\sqrt{(3x+1)^3}}dx+\int_{x_1}^{x_2}\frac{1}{2\sqrt{(3x+1)^3}}dx=\left[\frac{x-2}{\sqrt{3x+1}}\right]_{x_1}^{x_2}$ $\int_{x_1}^{x_2}\frac{3x+7}{2\sqrt{(3x+1)^3}}dx$ $=\left[\frac{x-2}{\sqrt{3x+1}}\right]_{x_1}^{x_2}-\int_{x_1}^{x_2}\frac{1}{2\sqrt{(3x+1)^3}}dx$ $=\left[\frac{x-2}{\sqrt{3x+1}}\right]_{x_1}^{x_2}-\frac{1}{2}\int_{x_1}^{x_2}(3x+1)^{-\frac{3}{2}}dx$ $=\left[\frac{x-2}{\sqrt{3x+1}}\right]_{0}^{5}-\left[\frac{1}{2}\frac{(3x+1)^{-\frac{1}{2}}}{3\left(-\frac{1}{2}\right)}\right]_{0}^{5}$ $=\frac{3}{4}-\frac{-2}{1}-\left(-\frac{1}{3(4)}+\frac{1}{3(1)}\right)$ $=2\frac{1}{2}$	M1 – seen or implied M1 – any equivalent form To show 7 = 8-1 or 8 = 7+ 1 M1 – standard integral M1 – show the correct limits substituted into a valid integral A1	
7a	$\angle DAE=\angle ABD$ (angles in alternate segment) $\angle ADE=\angle BAD$ (alternate angles of parallel lines) triangle ABD is similar to triangle DAE (AA similarity)	B1 B1 B1	
7b	$\angle BAD=\angle DCB$ (corresponding angles of similar triangles) $\angle BAD+\angle DCB=180°$ (angles in opposite segment)	B1 B1	

Downloaded from SingaKids.com

Qn	Solutions	Marks	
	$\angle BAD = \angle DCB = 90°$ BD is diameter (angle in semicircle = 90°)	 B1	
8a	$3(3^{x+1}) = 10 - 3^{-x}$ Let $u = 3^x$ $3(3u) = 10 - \frac{1}{u}$ $9u^2 - 10u + 1 = 0$ $3^x = \frac{1}{9}$ or $3^x = 1$ $x = -2$ *or* 0	 M1 – breakdown 3^{x+1} M1 – general QE M1 – eqn in x A1	
8b	$\log_{100} x + \lg y = 3$ $\frac{\lg x}{\lg 100} + \lg y = 3$ $\frac{\lg x}{2} + \lg y = 3$ $\lg\sqrt{x} + \lg y = 3$ $\lg \sqrt{x}\, y = 3$ $\sqrt{x}y = 10^3$ $y = \frac{1000}{\sqrt{x}}$	M1 – change base M1 – step before simplifying to one log term M1 – one log term A1	
9a	$(x - 2.5)^2 + (-\frac{1}{2}x + 5)^2 = \frac{365}{4}$ $x^2 - 5x + 6.25 + \frac{1}{4}x^2 - 5x + 25 = \frac{365}{4}$ $\frac{5}{4}x^2 - 10x + 31.25 = \frac{365}{4}$ $5x^2 - 40x + 125 = 365$ $5x^2 - 40x - 240 = 0$ $x = 12, x = -4$ $A(-4, 12)$	M1v - substitution M1 – general QE A1 A1	
9b	centre of circle (2.5,5) $y = 2x + c$ Sub centre of circle (2.5,5) $5 = 2(2.5) + c$ $c = 0$	B1 M1 – grad ⊥ seen B1 A1	
9c	Sub y = 0 into AB $0 = -\frac{1}{2}x + 10$ $x = 20$ D(20,0) M, Mid point AD = (8,6) Distance ME² = $(8 - 2.5)^2 + (6 - 5)^2 = \frac{125}{4} < \frac{365}{4}$	 M1 M1 M1 M1	
10a	Sub t = 0, v = 1	B1, B1	
10b	$a = -8e^{-2t} + 1 = 0$ $e^{-2t} = \frac{1}{8}$	M1 M1	

Downloaded from SingaKids.com

Qn	Solutions	Marks	
	$-2t = \ln\frac{1}{8}$ $-2t = \ln 1 - \ln 8 = -\ln 8$ $t = \frac{1}{2}\ln 8$ $v = 4e^{-\ln 8} + \frac{1}{2}\ln 8 - 3 = -1.46$	B1 A1	
10c	Since **velocity changes from positive to negative**, the particle **did change** its direction of motion	B1 B1	
10d	$s = -2e^{-2t} + \frac{t^2}{2} - 3t + c$ Sub t= 0, s = 6 $6 = -2 + c$ $c = 8$ Sub t= 2 $s = -2e^{-4} + 2 - 6 + 8 = 3.96\text{m}$	M1 integrate exp term M1 integrate power term M1 A1	

Downloaded from SingaKids.com

Downloaded from SingaKids.com