# NCHS
## PAPER-2

1 (a) Given that there is a term that is independent of $x$ in the expansion of $\left(5x^2 - \frac{1}{\sqrt{x}}\right)^n$, where $n$ is a positive integer, find the smallest possible value of $n$. [3]

Downloaded from SingaKids.com

(b) Using the value of $n$ found in part (a), explain if there is any term independent of $x$ in the expansion of $\left(1-\frac{1}{50x^3}\right)\left(5x^2-\frac{1}{\sqrt{x}}\right)^n$. [4]

Downloaded from SingaKids.com

2 The expression $10\,f(x) + 3\,f'(x) - f''(x) + 7\sin 2x + 3\cos 2x$, may be written as $10x + 43$, when $f'(x) = e^{5x} + 2\sin^2 x$. Find $f(x)$.

[6]

Downloaded from SingaKids.com

3 **Do not use a calculator in this question.**

It is given that $\tan A = \frac{5}{12}$ and that $\frac{\pi}{2} < A < \frac{3\pi}{2}$.

**(a)** By expressing $\cos 3A = \cos(2A + A)$, find the exact value of $\cos 3A$. [4]


Downloaded from SingaKids.com

(b) Find the exact value of $\tan\frac{A}{2}$. [4]

Downloaded from SingaKids.com

4 Coffee is poured into an empty cup. At time $t$ minutes after the coffee is poured, its temperature exceeds room temperature by $y$°C. The room temperature is 25°C.

(a)




The variables $t$ and $y$ are related by the equation $y = e^{kt+c}$, where $k$ and $c$ are constants. The diagram above shows the straight line graph obtained by plotting ln $y$ against $t$. The line passes through the points (5, 2.69) and (20, 0.88).
Find the value of $k$ and of $c$. [3]

(b) Calculate the time which the temperature of the coffee would drop to half of its initial temperature. [3]

Downloaded from SingaKids.com

(c) At the same time, when the coffee was poured into the cup, coffee of the same volume is also poured into an empty tumbler.

Similarly, at time $t$ minutes after the coffee is poured into the tumbler, its temperature exceeds room temperature by $y$°C and is modelled by another equation.

The solution to the equation $e^{(k+0.2)t} = e^{5-c}$ is the timing where the temperature of the coffee in both the cup and tumbler are equivalent. By using the diagram in part (a), outline the steps to find this timing. [4]

Downloaded from SingaKids.com

5





Diagram 1




Diagram 2

Diagram 1 shows a piece of square cardboard of side 80 cm.
Four small identical isosceles triangles, $P$, $Q$, $R$ and $S$ and four big identical isosceles triangles, $W$, $X$, $Y$ and $Z$ are removed. The remaining cardboard is folded along the dotted lines to form an open container as shown in Diagram 2.

(a) Let the height of the open container be $h$ cm, show that the total exterior area, $A$ cm$^2$, of the open container is $3200 - 80\sqrt{2}h + h^2$. [4]


Downloaded from SingaKids.com

(b) Find the value of $h$ for which the total exterior area of the open container is a maximum. [3]

(c) Hence, find the maximum total exterior area of the open container that can be obtained from the piece of square cardboard. [2]

Downloaded from SingaKids.com

6




The diagram shows part of the curve $y = 3 - 5\cos 2x$, which cuts the $x$-axis at $M$ and $N$. The tangent to the curve at $P$ is $-5$ and this tangent cuts the $x$-axis at $Q$. Find the area of the shaded region $PNQ$. [9]

Downloaded from SingaKids.com

Continuation of working space for Question 6.

Downloaded from SingaKids.com

7 (a) Show that $x + 2y$ is a factor of $4x^3 + x^2y - 11xy^2 + 6y^3$ and hence factorise $4x^3 + x^2y - 11xy^2 + 6y^3$ completely. [3]

Downloaded from SingaKids.com

(b) Hence, solve the equation $4^{p+1} + 2^{p+1} = 44 - 48(2^{-p})$. [5]

Downloaded from SingaKids.com

8 (a) The graph of $y = a \sin bx + c$ has one minimum point at $\left(\frac{\pi}{6}, 1\right)$ and the next maximum point after this has coordinates $\left(\frac{\pi}{2}, 9\right)$. Find the values of the constants $a$, $b$ and $c$. [3]

(b) A particle, travelling in a straight line, passes through a fixed point $O$. The velocity, $v$ m/s, at time t seconds, is given by $v = 3t^2 - 5t + 7$ for $0 \leq t \leq 5$.

(i) Find the acceleration of the particle when $t = 4$. [2]


Downloaded from SingaKids.com

After $t > 5$ seconds, the particle travels at a velocity, in m/s, where $v = -4t + 77$.

(ii) Find the total distance travelled by the particle in the first 30 seconds. [7]

Downloaded from SingaKids.com

9




The diagram shows a metal structure $ABCD$ consisting of five metal rods of different lengths. The length of $BC$ and $CD$ are 15 m and 8 m respectively. Angle $ADC = \theta$ for $0° < \theta < 90°$.

**(a)** Show that the total lengths, $P$ m, of the five metal rods used is $40 + 23 \sin\theta + 7 \cos\theta$. [3]

**(b)** Express $P$ in the form $40 + R \sin(\theta + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. [2]


Downloaded from SingaKids.com

(c) Find the value of $\theta$ if the total length of the five metal rods is 60 m. [3]

(d) State the minimum value of $\frac{1}{40+(R\sin(\theta+\alpha))^2}$ and the corresponding value of $\theta$ for which it occurs. [3]

Downloaded from SingaKids.com

10




The diagram shows a circle, centre $F$, with $BD$ as diameter. The point $E$ lies on the circle. The tangent at a point $B$ on the circle meets $DE$ extended at the point $A$. Point $C$ lies on $AB$ extended and $ED = AE$.

(a) State the relationship between the length of $DF$ and $AB$. Give reasons to support your answer. [2]

Downloaded from SingaKids.com

**(b)** Prove that

**(i)** triangle $ABE$ is similar to triangle $EDF$, [2]

**(ii)** $AD^2 - BD^2 = 2 \times BE \times ED$. [3]

Downloaded from SingaKids.com

Point $G$ is on minor arc $EB$ such that $BG$ bisects angle $EBA$.
Point $H$ is on the circle such that $EGHD$ is a cyclic quadrilateral.

(c) Prove that angle $GHD = 90° -$ angle $GBE$. [3]

**-End of Paper-**

Downloaded from SingaKids.com

**1(a)**

$$T_{r+1} = \binom{n}{r}(5x^2)^{n-r}\left(-\frac{1}{\sqrt{x}}\right)^r$$
$$= \binom{n}{r}(5)^{n-r}(-1)^r x^{2(n-r)}x^{-\frac{1}{2}r}$$
$$= \binom{n}{r}(5)^{n-r}(-1)^r x^{2n-\frac{5}{2}r}$$

Since there is a term independent of $x$,
$$x^{2n-\frac{5}{2}r} = x^0$$
$$2n - \frac{5}{2}r = 0$$
$$n = \frac{5}{4}r$$

Since both $n$ and $r$ are positive integers,
Smallest $n = 5$, when $r = 4$.

**1(b)**

For the term $\left(5x^2 - \frac{1}{\sqrt{x}}\right)^n$,

$T_{r+1} = \binom{n}{r}(5)^{n-r}(-1)^r x^{2n-\frac{5}{2}r}$

For constant, $r = 4$,

Constant $= \binom{5}{4}(5)^{5-4}(-1)^4$
$= 25$

For $x^5$,
$x^{2n-\frac{5}{2}r} = x^5$
$2n - \frac{5}{2}r = 5$
$r = 2$

Coefficient of $x^5 = \binom{5}{2}(5)^{5-2}(-1)^2$
$= 1250$

Term independent of $x$
$= (1)(125) + \left(-\frac{1}{50}\right)(1250)$
$= 0$

Hence, there are no term independent of $x$.

**2**

Method 1

$f'(x) = e^{5x} + 2\sin^2 x$
$= e^{5x} + 2(\sin x)^2$
$= e^{5x} + 1 - \cos 2x$

$f''(x) = \frac{d}{dx}(f'(x))$
$= \frac{d}{dx}(e^{5x} + 2(\sin x)^2)$
$= 5e^{5x} + 4\sin x\cos x$
$= 5e^{5x} + 2\sin 2x$

$f(x) = \int f'(x)\,dx$
$= \int e^{5x} + 1 - \cos 2x\,dx$
$= \frac{e^{5x}}{5} + x - \frac{\sin 2x}{2} + c$

Given
$10f(x) + 3f'(x) - f''(x) + 7\sin 2x + 3\cos 2x = 10x + 43$

$10\left(\frac{e^{5x}}{5} + x - \frac{\sin 2x}{2} + c\right) + 3(e^{5x} + 2\sin^2 x) - (5e^{5x} + 2\sin 2x) + 7\sin 2x + 3\cos 2x = 10x + 43$

$2e^{5x} + 10x - 5\sin 2x + 10c + 3e^{5x} + 6\sin^2 x - 5e^{5x} - 2\sin 2x + 7\sin 2x + 3\cos 2x = 10x + 43$

$10c + 6\sin^2 x + 3\cos 2x = 43$
$10c + 3(1 - \cos 2x) + 3\cos 2x = 43$
$c = 4$

$\therefore f(x) = \frac{e^{5x}}{5} + x - \frac{\sin 2x}{2} + 4$

Method 2

$f'(x) = e^{5x} + 2\sin^2 x$

$f''(x) = \frac{d}{dx}(f'(x))$
$= \frac{d}{dx}(e^{5x} + 2(\sin x)^2)$
$= 5e^{5x} + 4\sin x\cos x$
$= 5e^{5x} + 2\sin 2x$

Given
$10f(x) + 3f'(x) - f''(x) + 7\sin 2x + 3\cos 2x = 10x + 43$

$10f(x) + 3(e^{5x} + 2\sin^2 x) - (5e^{5x} + 2\sin 2x) + 7\sin 2x + 3\cos 2x = 10x + 43$


$10f(x) + 3e^{5x} + 6\sin^2 x - 5e^{5x} - 2\sin 2x + 7\sin 2x + 3\cos 2x = 10x + 43$

$10f(x) - 2e^{5x} + 5\sin 2x + 3(1 - \cos 2x) + \cos 2x = 10x + 43$

$10f(x) - 2e^{5x} + 5\sin 2x + 3 - 3\cos 2x + \cos 2x = 10x + 43$

$10f(x) = 2e^{5x} - 5\sin 2x + 10x + 40$
$f(x) = \frac{e^{5x}}{5} + x - \frac{\sin 2x}{2} + 4$

**3(a)**




$\sin A = -\frac{5}{13}$
$\cos A = -\frac{12}{13}$

Method 1

$\cos 3A = \cos(2A + A)$
$= \cos 2A\cos A - \sin 2A\sin A$
$= (1 - 2\sin^2 A)\cos A - (2\sin A\cos A)\sin A$
$= \left(1 - 2\left(-\frac{5}{13}\right)^2\right)\left(-\frac{12}{13}\right) - \left(2\left(-\frac{5}{13}\right)\left(-\frac{12}{13}\right)\right)\left(-\frac{5}{13}\right)$
$= -\frac{828}{2197}$

Method 2

$\cos 3A = \cos(2A + A)$
$= \cos 2A\cos A - \sin 2A\sin A$
$= (2\cos^2 A - 1)\cos A - (2\sin A\cos A)\sin A$
$= \left(2\left(-\frac{12}{13}\right)^2 - 1\right)\left(-\frac{12}{13}\right) - \left(2\left(-\frac{5}{13}\right)\left(-\frac{12}{13}\right)\right)\left(-\frac{5}{13}\right)$
$= -\frac{828}{2197}$

**3(b)**

$\tan A = \frac{2\tan\frac{A}{2}}{1 - \tan^2\frac{A}{2}}$

$\frac{5}{12} = \frac{2\tan\frac{A}{2}}{1 - \tan^2\frac{A}{2}}$

$5 - 5\tan^2\frac{A}{2} = 24\tan\frac{A}{2}$

$0 = 5\tan^2\frac{A}{2} + 24\tan\frac{A}{2} - 5$

$\tan\frac{A}{2} = \frac{-(24) \pm \sqrt{(24)^2 - 4(5)(-5)}}{2(5)}$
$= -5$ or $\frac{1}{5}$

Since $\pi < A < \frac{3\pi}{2}$,
$\frac{\pi}{2} < \frac{A}{2} < \frac{3\pi}{4}$ (2nd Quadrant),
$\therefore \tan\frac{A}{2} = -5$

Downloaded from SingaKids.com

**4(a)**

$$\text{Gradient} = \frac{0.88 - 2.69}{20 - 5} = -\frac{181}{1500}$$

$$Y - 2.69 = -\frac{181}{1500}(X - 5)$$

$$Y = -\frac{181}{1500}X + \frac{247}{75}$$

$$\ln y = -\frac{181}{1500}t + \frac{247}{75}$$

$$y = e^{-\frac{181}{1500}t + \frac{247}{75}}$$

$$\therefore k = -\frac{181}{1500}, c = \frac{247}{75}$$

**4(b)**

Initial Temperature

When $t = 0$,

$$y = e^{\frac{247}{75}}$$

$$\text{Initial Temperature} = 25 + e^{\frac{247}{75}} = 51.9325°\text{C}$$

Temperature to drop to half

$$y = \frac{51.9325}{2} - 25 = 0.966245$$

When $y = 0.966245$,

$$\ln(0.966245) = -\frac{181}{1500}t + \frac{247}{75}$$

$$\therefore t = 27.577 = 27.6 \text{ min (to 3sf)}$$

**4(c)**

$$e^{(k+0.2)t} = e^{5-c}$$

$$e^{kt}e^{0.2t} = \frac{e^5}{e^c}$$

$$e^{kt}e^{c} = \frac{e^5}{e^{0.2t}}$$

$$e^{kt+c} = e^{5-0.2t}$$

Tumbler Mode

$$y = e^{5-0.2t}$$

$$\ln y = 5 - 0.2t$$

Step 1:

Draw a straight-line graph in (a), where the gradient of the straight line is $-0.2$ and the $\ln y$-intercept is 5.

Step 2:

The intersection between the 2 straight line graphs will be the timing where the temperatures are equivalent.

**5(a)**

Consider $\Delta P$,

$$\text{hyp} = \sqrt{h^2 + h^2} = \sqrt{2}h$$

Consider $\Delta Z$,

$$\text{Side} = \frac{80 - \sqrt{2}h}{2}$$

$$= 2\left(\frac{80 - \sqrt{2}h}{2}\right)^2$$

$$= \frac{(80 - \sqrt{2}h)^2}{2}$$

$$\text{base} = \text{hyp} = \frac{80 - \sqrt{2}h}{\sqrt{2}}$$

$\therefore$ Total exterior area, $A$

$= 4 \times \text{base} \times \text{height} + \text{base} \times \text{base}$

$$= 4h\left(\frac{80 - \sqrt{2}h}{\sqrt{2}}\right) + \left(\frac{80 - \sqrt{2}h}{\sqrt{2}}\right)^2$$

$$= \frac{320}{\sqrt{2}}h - 4h^2 + \frac{6400 - 160\sqrt{2}h + 2h^2}{2}$$

$$= \frac{320\sqrt{2}}{2}h - 4h^2 + 3200 - 80\sqrt{2}h + h^2$$

$$= 3200 + 80\sqrt{2}h - 3h^2$$

Method 2

Consider $\Delta Z$,

$$\text{Side} = \frac{80 - \sqrt{2}h}{2}$$

$\therefore$ Total exterior area, $A$

$= 80 \times 80 - 4 \times \text{small triangle} - 4 \times \text{big triangle}$

$$= 6400 - 4 \times \left(\frac{1}{2}\right)(\sqrt{2}h)^2 - 4 \times \left(\frac{1}{2}\right)\left(\frac{80 - \sqrt{2}h}{2}\right)^2$$

$$= 6400 - 4h^2$$

$$- \frac{6400 - 160\sqrt{2}h + 2h^2}{2}$$

$$= 6400 - 4h^2 - 3200 + 80\sqrt{2}h - h^2$$

$$= 3200 + 80\sqrt{2}h - 3h^2$$

**5(b)**

When the total exterior area is a maximum, $\frac{dA}{dh} = 0$,

$$A = 3200 + 80\sqrt{2}h - 3h^2$$

$$\frac{dA}{dh} = 80\sqrt{2} - 6h$$

$$0 = 80\sqrt{2} - 6h$$

$$h = \frac{80\sqrt{2}}{6} = \frac{40\sqrt{2}}{3} \text{ cm}$$

$$\frac{d^2A}{dh^2} = -6 (< 0)$$

$\therefore A$ is maximum.

**5(c)**

When $h = \frac{40\sqrt{2}}{3}$,

$$A = 3200 + 80\sqrt{2}\left(\frac{40\sqrt{2}}{3}\right) - 3\left(\frac{40\sqrt{2}}{3}\right)^2$$

$$= 4266\frac{2}{3} \text{ cm}^2$$

**6**

$$y = 3 - 5\cos 2x$$

Point N

When $y = 0$,

$$3 - 5\cos 2x = 0$$

$$\cos 2x = \frac{3}{5}$$

$$\alpha = \cos^{-1}\frac{3}{5} = 0.927295$$

$2x$ lies in 1st (rej)/ 4th quad.

$$2x = 5.35589$$

$$x = 2.67795$$

Point P

$$y = 3 - 5\cos 2x$$

$$\frac{dy}{dx} = -5(-\sin 2x)(2) = 10\sin 2x$$

When $\frac{dy}{dx} = -5$,

$$10\sin 2x = -5$$

$$\sin 2x = -\frac{1}{2}$$

$$\alpha = \sin^{-1}\frac{1}{2} = \frac{\pi}{6}$$

$2x$ lies in 3rd/ 4th (rej) quad.

$$2x = \frac{7\pi}{6}$$

$$x = \frac{7\pi}{12}$$

When $x = \frac{7\pi}{12}$,

$$y = 3 - 5\cos 2\left(\frac{7\pi}{12}\right) = 7.330127$$

$$\therefore P\left(\frac{7\pi}{12}, 7.330127\right)$$

Equation PQ, Point Q

$$y - 7.330127 = -5\left(x - \frac{7\pi}{12}\right)$$

$$y = -5x + 16.4931$$

When $y = 0$,

$$-5x + 16.4931 = 0$$

$$x = 3.29862$$

$$\therefore Q(3.29862, 0)$$

Area below the curve from x to Q

$$\text{area} = \int_{\frac{7\pi}{12}}^{2.67795} 3 - 5\cos 2x \, dx$$

$$= \left[3x - \frac{5\sin 2x}{2}\right]_{\frac{7\pi}{12}}^{2.67795}$$

$$= 10.0338 - 6.74779$$

$$= 3.28601 \text{ units}^2$$

$\therefore$ shaded area

= area of triangle − area below curve

$$= \frac{1}{2}(7.330127)\left(3.29862 - \frac{7\pi}{12}\right) - 3.28601$$

$$= 5.37307 - 3.28601$$

$$= 2.0871$$

$$= 2.09 \text{ units}^2$$

Downloaded from SingaKids.com

**7(a)** To prove $(x + 2y)$ is a factor:

Method 1

Let $f(x) = 4x^3 + x^2y - 11xy^2 + 6y^3$,

$f(-2y) = 4(-2y)^3 + (-2y)^2y - 11(-2y)y^2 + 6y^3$

$= -32y^3 + 4y^3 + 22y^3 + 6y^3$

$= 0$

Since $f(-2y)$, by factor theorem, $(x + 2y)$ is a factor.

Method 2

$$\begin{array}{r} 4x^2 - 7yx + 3y^2 \\ x + 2y \overline{)\,4x^3 + x^2y - 11xy^2 + 6y^3} \\ -(4x^3 + \ldots) \\ \hline \ldots - 11\ldots \\ \ldots \\ \hline -(3y^2x + 6y^3) \\ \hline 0 \end{array}$$

Since the remainder is 0, $(x + 2y)$ is a factor.

Hence factorise:

Method 1 – Compare coefficients

$f(x) = (x + 2y)(4x^2 + bx \ldots)$

Comparing coefficients of,

$x^2$ : $\ldots = -7y$

Method 2 – Division

As shown earlier

$\therefore f(x) = (x + 2y)(4x^2 - 7yx + 3y^2)$

$= (x + 2y)(x - y)(4x - 3y)$

**7(b)**

$4^{p+1} + 2^{p+1} = 44 - 48(2^{-p})$

$2^{2(p+1)} + 2^{p+1} = 44 - \dfrac{48}{2^p}$

$2^{2p}\,2^2 + 2^p\,2 = 44 - \dfrac{48}{2^p}$

Let $u = 2^p$,

$4u^2 + 2u = 44 - \dfrac{48}{u}$

$(\times u)$

$4u^3 + 2u^2 - 44u + 48 = 0$

From (a),

$48 = 6y^3$

$y^3 = 8$

$y = 2$

From (a), $x = u$

From (a),

$f(x) = (x + 2y)(x - y)(4x - 3y)$


$f(x) = (x + 4)(x - 2)(4x - 6) = 0$

| $x = -4$ | $x = 2$ | $x = \frac{3}{2}$ |
|---|---|---|
| $2^p$ | $2^p = 2$ | $2^p = \frac{3}{2}$ |
| $= -4$ (rej) | $\therefore p = 1$ | |

$\therefore p = \dfrac{\lg \frac{3}{2}}{\lg 2}$

$= 0.585$ ($t$ o 3sf)

**8(a)**




| $y = a \sin bx + c$ | | |
|---|---|---|
| $c = \frac{9+1}{2} = 5$ | $\frac{1}{2}$ period $= \frac{\pi}{2} - \frac{\pi}{6}$; period $= \frac{2\pi}{3}$; $\frac{2\pi}{b} = \frac{2\pi}{3}$; $b = 3$ | Amplitude $= 9 - 5 = 4$; $a = -4$ |

**8(b) (i)**

$a = \dfrac{dv}{dt}$

$= 6t - 5$

When $t = 4$,

$a = 6(4) - 5$

$= 19$ m/s$^2$

**8(b) (ii)**

For $0 \le t \le 5$

$v = 3t^2 - 5t + 7$

Displacement expression:

$s = \int 3t^2 - 5t + 7\ dt$

$= \dfrac{3t^3}{3} - \dfrac{5t^2}{2} + 7t + c$

When $t = 0$, $s = 0$,

$c = 0$

$\therefore s = \dfrac{3t^3}{3} - \dfrac{5t^2}{2} + 7t$

Turning point:

When $v = 0$,

$3t^2 - 5t + 7 = 0$

$b^2 - 4ac = (-5)^2 - 4(3)(7)$

$= -59\ (< 0)$

Since $b^2 - 4ac < 0$, there is no turning point.

When $t = 0$, $s = 0$ m

$t = 5$, $s = 97.5$ m

For $t \ge 5$,

$v = -4t + 77$

Displacement expression:

$s = \int -4t + 77\ dt$

$= -\dfrac{4t^2}{2} + 77t + d$

When $t = 5$, $s = 97.5$,

$97.5 = -2(5)^2 + 77(5) + d$

$d = -\dfrac{475}{2}$

$\therefore s = -2t^2 + 77t - \dfrac{475}{2}$

Turning point:


When $v = 0$,

$-4t + 77 = 0$

$t = \dfrac{77}{4} = 19\dfrac{1}{4}$

When $t = 19\frac{1}{4}$, $s = 503.625$ m

When $t = 30$, $s = 272.5$ m




$\therefore$ total distance

$= 503.625 + (503.625 - 272.5)$

$= 734.75$ m

**9(a)**

Consider $\Delta BCD$, using Pythagoras theorem,

$BD = \sqrt{8^2 + 15^2} = 17$

Consider $\Delta BXC$,

$CX = 8\cos\theta$

$BX = 8\sin\theta$

Consider $\Delta CYD$,

$CY = 15\sin\theta$

$YD = 15\cos\theta$

Total lengths

$= AB + 8 + 15 + AD + BD$

$= (CY - CX) + 8 + 15 + (BX + YD) + BD$

$= 15\sin\theta - 8\cos\theta + 8 + 15 + 8\sin\theta + 15\cos\theta + 17$

$= 40 + 23\sin\theta + 7\cos\theta$ (shown)

**9(b)**

$P = 40 + 23\sin\theta + 7\cos\theta$

$= 40 + R\sin(\theta + \alpha)$

$R = \sqrt{23^2 + 7^2}$

$= \sqrt{578}$

$= 24.04163 = 24.0$ (to 3sf)

Downloaded from Singakids.com

| | |
|---|---|
| | $\alpha = \tan^{-1}\frac{7}{23} = 16.9275°$<br>$= 16.9°$ (to 1 dp)<br><br>$\therefore P = 40 + 24.0\sin(\theta + 16.9°)$ |
| 9(c) | When $P = 60$,<br>$40 + 24.04163\sin(\theta + 16.9275°) = 60$<br>$\sin(\theta + 16.9275°) = 0.83189$<br>$\alpha = \sin^{-1}0.83189$<br>$= 56.2934°$<br>$(\theta + 16.9275°)$ lies in 1st / 2nd (rej) quad.<br>$\theta + 16.9275° = 56.2934°$<br>$\theta = 39.3659°$<br>$= 39.4°$ (to 1 dp) |
| 9(d) | $\max(R\sin(\theta + \alpha))^2 = \left(\sqrt{578}\right)^2 = 578$<br><br>$\therefore \min\frac{1}{40 + (R\sin(\theta + \alpha))^2}$<br>$= \frac{1}{40 + 578} = \frac{1}{618}$<br><br>$\theta + 16.9275° = 90°$<br>$\theta = 73.0725°$<br>$= 73.1°$ (to 1 dp) |
| 10(a) | Since $E$ is the midpoint of $AD$ (given),<br>$F$ is the midpoint of $DB$ (given),<br>By midpoint theorem,<br><br>$FE = \frac{1}{2}AB$ |
| 10(b) (i) | $\angle ABE$<br>$= \angle EDF$ (alternate segment theorem)<br><br>From (a), by midpoint theorem,<br>$EF \parallel AB$<br>$\angle DEF$<br>$= \angle BAE$ (corresponding angles, $EF \parallel AB$) |

| | |
|---|---|
| | $\therefore \Delta ABE$ is similar to $\Delta EDF$ (AA similarity) |
| 10(b) (ii) | From (i),<br>$\frac{AB}{ED}$<br>$= \frac{BE}{DF}$ (corresponding sides of similar $\Delta$s)<br>$AB \times DF = BE \times ED$<br>$AB \times \frac{1}{2}AB = BE \times ED$<br>$AB^2 = 2 \times BE \times ED$<br><br>Since $\angle DBA = 90°$ (tangent $\perp$ radius),<br>Using Pythagoras' theorem,<br>$AD^2 - BD^2 = 2 \times BE \times ED$ |
| 10(c) | Method 1<br>Let $\angle GBE$ be $\alpha$,<br>$\angle GBA = \alpha$ ($BG$ bisects $\angle EBA$)<br>$\angle GEB$<br>$= \alpha$ (alternate segment theorem)<br>$\angle DEB = 90°$ (rt $\angle$ in a semicircle)<br><br>$\angle GHD$<br>$= 180°$<br>$- (90° + \alpha)$ ($\angle$ in opp segment)<br>$= 90° - \alpha$<br>$= 90° - \angle GBE$<br><br>Method 2<br>Let $\angle GBE$ be $\alpha$,<br>$\angle GBA = \alpha$ ($BG$ bisects $\angle EBA$)<br>$\angle DBA = 90°$ (tangent $\perp$ radius)<br>$\angle GBD = 90° - \alpha$<br><br>$\angle GHD$<br>$= \angle GBD$ ($\angle$ in the same segment)<br>$= 90° - \alpha$<br>$= 90° - \angle GBE$ |

Downloaded from SingaKids.com