1 (a) Given that there is a term that ~~is~~ independent of x in the expansion of $\left(5x^2 - \frac{1}{\sqrt{x}}\right)^n$, where n is a positive integer, find the smallest possible value of n. [3]

Scanned with CamScanner
Downloaded from Stuvia.com

(b) Using the value of n found in part **(a)**, explain if there is any term independent of x in the expansion of $\left(1-\frac{1}{50x^5}\right)\left(5x^2-\frac{1}{\sqrt{x}}\right)^n$. [4]

Scanned with CamScanner
Downloaded from Stuguides.com

2 The expression $10\,f(x) + 3\,f'(x) - f''(x) + 7\sin 2x + 3\cos 2x$, may be written as $10x + 43$, when $f'(x) = e^{5x} + 2\sin^2 x$. Find $f(x)$.

[6]

Scanned with CamScanner
Downloaded from Stuvia.com

3 **Do not use a calculator in this question.**

It is given that $\tan A = \frac{5}{12}$ and that $\frac{\pi}{2} < A < \frac{3\pi}{2}$.

(a) By expressing $\cos 3A = \cos(2A + A)$, find the exact value of $\cos 3A$. [4]


Scanned with CamScanner
Downloaded from Stuvia.com

(b) Find the exact value of $\tan\frac{A}{2}$. [4]

Scanned with CamScanner
Downloaded from Stuguides.com

4 Coffee is poured into an empty cup. At time t minutes after the coffee is poured, its temperature exceeds room temperature by y°C. The room temperature is 25°C.

(a)



The variables t and y are related by the equation $y = e^{kt+c}$, where k and c are constants. The diagram above shows the straight line graph obtained by plotting ln y against t. The line passes through the points (5, 2.69) and (20, 0.88).
Find the value of k and of c. [3]

(b) Calculate the time which the temperature of the coffee would drop to half of its initial temperature. [3]

Scanned with CamScanner
Downloaded from Stuvia.com

(c) At the same time, when the coffee was poured into the cup, coffee of the same volume is also poured into an empty tumbler.

Similarly, at time t minutes after the coffee is poured into the tumbler, its temperature exceeds room temperature by y°C and is modelled by another equation.

The solution to the equation $e^{(k+0.2)t} = e^{5-c}$ is the timing where the temperature of the coffee in both the cup and tumbler are equivalent. By using the diagram in part **(a)**, outline the steps to find this timing. [4]

Scanned with CamScanner
Downloaded from Stuguides.com

5




Diagram 1




Diagram 2

Diagram 1 shows a piece of square cardboard of side 80 cm.
Four small identical isosceles triangles, P, Q, R and S and four big identical isosceles triangles, W, X, Y and Z are removed. The remaining cardboard is folded along the dotted lines to form an open container as shown in Diagram 2.

(a) Let the height of the open container be h cm, show that the total exterior area, A cm^2, of the open container is $3200 - 80\sqrt{2}h + h^2$. [4]


Scanned with CamScanner
Downloaded from SingaMath.com

(b) Find the value of h for which the total exterior area of the open container is a maximum. [3]

(c) Hence, find the maximum total exterior area of the open container that can be obtained from the piece of square cardboard. [2]

Scanned with CamScanner
Downloaded from Stuguides.com

6



The diagram shows part of the curve $y = 3 - 5\cos 2x$, which cuts the x-axis at M and N. The tangent to the curve at P is -5 and this tangent cuts the x-axis at Q. Find the area of the shaded region PNQ. [9]

Scanned with CamScanner
Downloaded from Stuguides.com

Continuation of working space for Question 6.

Scanned with CamScanner
Downloaded from Stuvia.com

7 (a) Show that $x + 2y$ is a factor of $4x^3 + x^2y - 11xy^2 + 6y^3$ and hence factorise $4x^3 + x^2y - 11xy^2 + 6y^3$ completely. [3]

Scanned with CamScanner
Downloaded from Stuvia.com

(b) Hence, solve the equation $4^{p+1} + 2^{p+1} = 44 - 48(2^{-p})$. [5]

Scanned with CamScanner
Downloaded from Stuguides.com

8 **(a)** The graph of $y = a \sin bx + c$ has one minimum point at $\left(\frac{\pi}{6}, 1\right)$ and the next maximum point after this has coordinates $\left(\frac{\pi}{2}, 9\right)$. Find the values of the constants a, b and c. [3]

(b) A particle, travelling in a straight line, passes through a fixed point O. The velocity, v m/s, at time t seconds, is given by $v = 3t^2 - 5t + 7$ for $0 \leq t \leq 5$.

(i) Find the acceleration of the particle when $t = 4$. [2]


Scanned with CamScanner
Downloaded from Stuguides.com

After $t > 5$ seconds, the particle travels at a velocity, in m/s, where $v = -4t + 77$.

(ii) Find the total distance travelled by the particle in the first 30 seconds. [7]

Scanned with CamScanner
Downloaded from Stuvera.com

9



The diagram shows a metal structure $ABCD$ consisting of five metal rods of different lengths. The length of BC and CD are 15 m and 8 m respectively. Angle $ADC = \theta$ for $0° < \theta < 90°$.

(a) Show that the total lengths, P m, of the five metal rods used is $40 + 23\sin\theta + 7\cos\theta$. [3]

(b) Express P in the form $40 + R\sin(\theta + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. [2]


Scanned with CamScanner
Downloaded from Stuvia.com

(c) Find the value of θ if the total length of the five metal rods is 60 m. [3]

(d) State the minimum value of $\frac{1}{40+(R\sin(\theta+\alpha))^2}$ and the corresponding value of θ for which it occurs. [3]

Scanned with CamScanner
Downloaded from Stuguides.com

10



The diagram shows a circle, centre F, with BD as diameter. The point E lies on the circle. The tangent at a point B on the circle meets DE extended at the point A. Point C lies on AB extended and $ED = AE$.

(a) State the relationship between the length of DF and AB. Give reasons to support your answer. [2]

Scanned with CamScanner
Downloaded from Stuvia.com

(b) Prove that

(i) triangle ABE is similar to triangle EDF, [2]

(ii) $AD^2 - BD^2 = 2 \times BE \times ED$. [3]

Scanned with CamScanner
Downloaded from Stuguides.com

Point G is on minor arc EB such that BG bisects angle EBA.
Point H is on the circle such that $EGHD$ is a cyclic quadrilateral.

(c) Prove that angle $GHD = 90° -$ angle GBE. [3]

-End of Paper-

Scanned with CamScanner
Downloaded from Stuguides.com