Name:	Index No.:	Class:

PRESBYTERIAN HIGH SCHOOL

ADDITIONAL MATHEMATICS 4049/01
Paper 1

19 August 2024 Monday 2 hours 15 min

PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL

2024 SECONDARY FOUR EXPRESS / FIVE NORMAL (ACADEMIC) PRELIMINARY EXAMINATIONS

DO NOT OPEN THIS QUESTION PAPER UNTIL YOU ARE TOLD TO DO SO.

INSTRUCTIONS TO CANDIDATES
Write your name, index number and class in the spaces provided above.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.
Write your answers in the spaces provided below the questions.
Give non-exact numerical answers correct to 3 significant figures or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 90.

For Examiner's Use													
Qn	1	2	3	4	5	6	7	8	9	10	11	12	*Marks Deducted*
Marks													

Category	Accuracy	Units	Notations	Others
Question No.				

TOTAL MARKS
90

This question paper consists of **19** printed pages and **1** blank page.

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 Express $\dfrac{3x^2+14x+6}{x^2(x+3)}$ as the sum of 3 partial fractions. [5]

Downloaded from SingaKids.com

2 **(a)** State, in terms of π,

(i) the principal value of $\tan^{-1}\left(-\sqrt{3}\right)$, [1]

(ii) the values between which the principal value of $\sin^{-1}x$ must lie. [1]

(b) Given that A is a reflex angle and $\cos A = -\frac{4}{5}$, find the exact value of $\cos(A + 30°)$ without the use of a calculator. [3]

Downloaded from SingaKids.com

3 **(a)** Find the set of values of the constant k for which the curve $y = -kx^2 - 2x + 2k - 3$ does not intersect the x-axis. [3]

(b) Using the answer in **part (a)**, explain whether it is possible for $-kx^2 - 2x + 2k - 3$ to be positive for all x. [2]

Downloaded from SingaKids.com

4 A roller coaster is being designed such that the height, h m, of a rider above the ground in a section of the roller coaster ride is given by $h = 10x - 2x^2 - 4$, where x is the horizontal distance of the rider from the starting point and $1 \leq x \leq 4$.


(a) Express h in the form $a + b(x + c)^2$ where a, b and c are constants to be determined. [3]

(b) Hence, explain why the rider cannot reach a height of 10 m. [1]

(c) After testing the prototype, the roller coaster designer wants to make the ride more exciting by moving the highest point of this section of the roller coaster ride up by 0.2 m and left by 0.1 m. Write down a possible new expression for h. [1]

Downloaded from SingaKids.com

5 Water is poured into an empty inverted conical container with radius 6 cm and slant height 10 cm. After t seconds, the radius of the top surface of the water is r cm.



(a) Show that the surface area, S, of water in contact with the container at any time is given by $S = \frac{5}{3}\pi r^2$.

[The curved surface area of a cone of base radius r and slant height l is πrl.] [2]

(b) Water is poured into the container such that r increases at a constant rate. Given that it takes 30 seconds to completely fill up the empty container, write down the rate at which r increases. [1]

Downloaded from SingaKids.com

(c) Hence calculate the rate at which S increases when the container is one-eighth filled. [4]

Downloaded from SingaKids.com

6 A curve has equation $y = \dfrac{30(2x-1)^3}{e^{3x}}$, where $x > 0$.


(a) Find the x-coordinates of the stationary points of the curve. [5]

(b) Determine the nature of each of the stationary points. [3]

(c) Using your answer in **part (b)**, infer and write down the set of values of x for which y is an increasing function. [2]

Downloaded from SingaKids.com

7 The diagram shows part of the curve $y = \frac{40}{x+8} - 20$, where $x > -8$. The curve intersects the x-axis and y-axis at P and Q respectively. The tangent to the curve at R is parallel to line PQ.



(a) Show that the coordinates of point R are $(-4, -10)$. [5]


Downloaded from SingaKids.com

(b) Given that the point S has coordinates $(-2, -5)$, find the area of the quadrilateral $PRQS$. [2]

(c) Determine, with reason, whether $PRQS$ is a parallelogram. [3]

Downloaded from SingaKids.com

8 The diagram shows part of the curve $y = \dfrac{30}{2x-1}$ and the line $y = -\dfrac{4}{3}x + \dfrac{38}{3}$. The curve intersects the line at the points A and B.



(a) Find the coordinates of the points A and B. [3]

Downloaded from SingaKids.com

(b) Find the area of the shaded region. [4]

(c) Show that the area bounded by the curve, the y-axis and the lines $y = 2$ and $y = 6$ can be expressed in the form $15\ln p + q$, where p and q are constants to be determined. [3]

Downloaded from SingaKids.com

9 The function f is given by $f(x) = x^3 - x^2 + ax - 1$.

It is given that $\frac{f(x)}{x-1} = Q(x) + \frac{3}{x-1}$, where $Q(x)$ is a quadratic function.

(a) Explain how you can use Remainder Theorem to show that $a = 4$. [3]

(b) By using long division, divide $f(x)$ by $x^2 + 1$.

Hence, express $\frac{f(x)}{x^2+1}$ in the form $px + q + \frac{rx}{x^2+1}$, where p, q and r are constants to be determined. [2]

Downloaded from SingaKids.com

(c) Find $\frac{d}{dx}\ln(x^2+1)$. [1]

(d) Hence, using your results from **part (b) and (c)**, find $\int \frac{f(x)}{x^2+1}dx$. [2]

Downloaded from SingaKids.com

10 The equation of a circle, C_1, with centre A, is $x^2 + y^2 - 6x + 8y - 75 = 0$.

(a) Find the radius of the circle and the coordinates of its centre, A. [4]


(b) A second circle , C_2, with centre B, has radius 12 units.
The equation of the perpendicular bisector of AB is $y = -\frac{1}{2}x + 5$.
Find the equation of the second circle, C_2. [5]

Downloaded from SingaKids.com

(c) P is a point on the perpendicular bisector of AB, where P is **not** the midpoint of A and B.
Explain whether triangle PAB is an isosceles triangle. [1]

Downloaded from SingaKids.com

11 (a) Sketch the graph of $y = e^{-x}$ on the given axes. Label any axial intercepts. [1]




(b) The curve $y = f(x)$ is such that $f'(x) = e^{2x+1} + e^{-x}$.

(i) Explain why the curve has no stationary points. [2]

(ii) The curve passes through the point (ln 4, 8e). Without the use of a calculator, find an expression for $f(x)$. Show all working clearly. [6]

Downloaded from SingaKids.com

12 The diagram shows the graph of $y = \mathrm{f}'(x)$.



(a) Write down the range of values of x for which the **function** f decreases as x increases. [1]

(b) The above function f is given by $\mathrm{f}(x) = 12k^2x - 2x^3 - 3kx^2$, where k is a positive constant. By solving a suitable inequality, use the answer in **part (a)** to find the value of the constant k. [4]

(c) Explain how you can use the above graph of $y = \mathrm{f}'(x)$ to determine the value of x for which f(x) is a maximum value. [1]

END OF PAPER

Downloaded from SingaKids.com

Downloaded from SingaKids.com

Name:	Index No.:	Class:

PRESBYTERIAN HIGH SCHOOL

ADDITIONAL MATHEMATICS **4049/01**
Paper 1

19 August 2024 Monday 2 hours 15 min

PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL
PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL PRESBYTERIAN HIGH SCHOOL

2024 SECONDARY FOUR EXPRESS / FIVE NORMAL (ACADEMIC)
PRELIMINARY EXAMINATIONS

MARKING SCHEME

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\ldots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 Express $\dfrac{3x^2+14x+6}{x^2(x+3)}$ as the sum of 3 partial fractions. [5]

$\dfrac{3x^2+14x+6}{x^2(x+3)} = \dfrac{A}{x} + \dfrac{B}{x^2} + \dfrac{C}{x+3}$ **[M1, accept $\dfrac{Ax+B}{x^2} + \dfrac{C}{x+3}$]**

$3x^2+14x+6 = Ax(x+3) + B(x+3) + Cx^2$ **[M1 for comparing numerator]**

Subst. $x=0$: $6 = 3B$ **[M1 correct method of subst. or compare coeff.]**

$B = 2$

Subst. $x=-3$: $-9 = 9C$

$C = -1$

Subst. $x=1$: $23 = 4A+7$

$A = 4$

Therefore, $\dfrac{3x^2+14x+6}{x^3+3x^2} = \dfrac{4}{x} + \dfrac{2}{x^2} - \dfrac{1}{x+3}$ **[A1 for 2 correct terms; A2 for all correct]**

Downloaded from SingaKids.com

2 **(a)** State, in terms of π,

(i) the principal value of $\tan^{-1}\left(-\sqrt{3}\right)$, [1]

(ii) the values between which the principal value of $\sin^{-1}x$ must lie. [1]

(b) Given that A is a reflex angle and $\cos A = -\frac{4}{5}$, find the exact value of $\cos(A+30°)$ without the use of a calculator. [3]

(a) **(i)** $-\frac{\pi}{3}$ **[B1]**

(ii) $-\frac{\pi}{2} \leqslant \sin^{-1}x \leqslant \frac{\pi}{2}$ **[B1]**

(b) A lies in the 3rd quadrant.

$$\sin A = -\frac{3}{5} \quad \textbf{[B1]}$$

$$\cos(A+30°) = \cos A\cos 30° - \sin A \sin 30°$$

$$= \left(-\frac{4}{5}\right)\left(\frac{\sqrt{3}}{2}\right) - \left(-\frac{3}{5}\right)\left(\frac{1}{2}\right) \quad \left[\textbf{M1 for } \cos 30° = \frac{\sqrt{3}}{2} \textbf{ or } \sin 30° = \frac{1}{2}\right]$$

$$= \frac{3}{10} - \frac{2}{5}\sqrt{3} \quad \textbf{[A1]}$$

Downloaded from SingaKids.com

3 **(a)** Find the set of values of the constant k for which the curve $y = -kx^2 - 2x + 2k - 3$ does not intersect the x-axis. [3]

$b^2 - 4ac < 0$

$(-2)^2 - 4(-k)(2k-3) < 0$ [**M1 for $b^2 - 4ac < 0$ and at least 2 correct substitution of a, b, c**]

$4 + 8k^2 - 12k < 0$

$2k^2 - 3k + 1 < 0$

$(2k-1)(k-1) < 0$ [**M1 for factorisation/finding x – intercepts**]

$\frac{1}{2} < k < 1$ [**A1**]

(b) Using the answer in **part (a)**, explain whether it is possible for $-kx^2 - 2x + 2k - 3$ to be positive for all x. [2]

Since $\frac{1}{2} < k < 1$, **coefficient of $x^2 = -k < 0$.** **[M1]**

The curve lies entirely below the axis, therefore it is **not possible** for $-kx^2 - 2x + 2p - 3$ to be positive for all x. **[A1]**

Downloaded from SingaKids.com

4 A roller coaster is being designed such that the height, h m, of a rider above the ground in a section of the roller coaster ride is given by $h = 10x - 2x^2 - 4$, where x is the horizontal distance of the rider from the starting point and $1 \leq x \leq 4$.

(a) Express h in the form $a + b(x + c)^2$ where a, b and c are constants to be determined. [3]

$$h = 10x - 2x^2 - 4$$
$$= -2(x^2 - 5x + 2) \qquad \textbf{[M1 factorise]}$$
$$= -2\left[\left(x - \frac{5}{2}\right)^2 - \left(\frac{5}{2}\right)^2 + 2\right]$$
$$= 8.5 - 2\left(x - \frac{5}{2}\right)^2 \qquad \textbf{[B1, B1 for each term]}$$

(b) Hence, explain why the rider cannot reach a height of 10 m. [1]


Since the **maximum value of the rider is 8.5 m** and **8.5 < 10**, the rider cannot reach a height of 10 m. **[B1]**

(c) After testing the prototype, the roller coaster designer wants to make the ride more exciting by moving the highest point of this section of the roller coaster ride up by 0.2 m and left by 0.1 m. Write down a possible new expression for h. [1]

$h = 9.7 - 2(x - 2.4)^2$ **[B1]**

Downloaded from SingaKids.com

5 Water is poured into an empty inverted conical container with radius 6 cm and slant height 10 cm. After t seconds, the radius of the top surface of the water is r cm.



(a) Show that the surface area, S, of water in contact with the container at any time is given by $S = \frac{5}{3}\pi r^2$. [2]

By similar triangles, $\frac{l}{10} = \frac{r}{6}$

$$l = \frac{5}{3}r \qquad \textbf{[M1]}$$

$$S = \pi(r)\left(\frac{5}{3}r\right) = \frac{5}{3}\pi r^2 \qquad \textbf{[A1]}$$

(b) Water is poured into the container such that r increases at a constant rate. Given that it takes 30 seconds to completely fill up the empty container, write down the rate at which r increases. [1]

$$\frac{dr}{dt} = \frac{6}{30} = \frac{1}{5} \text{ cm/s} \qquad \textbf{[B1]}$$

(c) Hence calculate the rate at which S increases when the container is one-eighth filled. [4]

$$\frac{dS}{dt} = \frac{dS}{dr} \times \frac{dr}{dt} \qquad \textbf{[M1 for chain rule]}$$

$$= \frac{10}{3}\pi r \times \frac{1}{5} \qquad \left[\textbf{B1 for } \frac{10}{3}\pi r\right]$$

When the container is one-eighth filled,

$$\left(\frac{r}{6}\right)^3 = \frac{1}{8}$$

$$r = \frac{1}{2} \times 6 = 3 \qquad \textbf{[B1 for } r = 3\textbf{]}$$

When $r = 3$, $\frac{dS}{dt} = \frac{2}{3}\pi(3) = 2\pi$ (or 6.28 cm/s to 3 s.f.) **[A1]**

Downloaded from SingaKids.com

6 A curve has equation $y = \dfrac{30(2x-1)^3}{e^{3x}}$, where $x > 0$.

(a) Find the x-coordinates of the stationary points of the curve. [5]

$$\frac{dy}{dx} = \frac{e^{3x} \cdot 30(3)(2x-1)^2(2) - 30(2x-1)^3 \cdot 3e^{3x}}{e^{6x}}$$

M1 for chain rule
i.e. $3(2x-1)^2(2)$ or $3e^{3x}$ seen
M1 for correct use of quotient rule even w/o chain rule

$$= \frac{90e^{3x}(2x-1)^2(2-2x+1)}{e^{6x}}$$

$$= \frac{90(2x-1)^2(3-2x)}{e^{3x}}$$ **[M1 factorising]**

At stationary point, $\dfrac{dy}{dx} = 0$

$(2x-1)^2 = 0$ or $3-2x=0$ **[M1]**

$x = \dfrac{1}{2}$ or $x = \dfrac{3}{2}$ **[A1]**

(b) Determine the nature of each of the stationary points. [3]

x	$\left(\frac{1}{2}\right)^-$	$\frac{1}{2}$	$\left(\frac{1}{2}\right)^+$
$\frac{dy}{dx}$	+	0	+
	↗	—	↗

x	$\left(\frac{3}{2}\right)^-$	$\frac{3}{2}$	$\left(\frac{3}{2}\right)^+$
$\frac{dy}{dx}$	+	0	−
	↗	—	↘

[M1 for first derivative test]


Therefore, the stationary point at $x = \dfrac{1}{2}$ is a **point of inflexion** and the stationary point at $x = \dfrac{3}{2}$ is a **maximum point**. **[A1, A1]**

(c) Using your answer in **part (b)**, infer and write down the set of values of x for which y is an increasing function. [2]

$0 < x < \dfrac{3}{2}, x \neq \dfrac{1}{2}$ **[B1,B1]** **OR** $0 < x < \dfrac{1}{2}, \dfrac{1}{2} < x < \dfrac{3}{2}$ **[B1,B1]**

Downloaded from SingaKids.com

7 The diagram shows part of the curve $y = \frac{40}{x+8} - 20$, where $x > -8$. The curve intersects the x-axis and y-axis at P and Q respectively. The tangent to the curve at R is parallel to line PQ.



(a) Show that the coordinates of point R are $(-4, -10)$. [5]

When $x = 0$, $y = -15$.

When $y = 0$, $x = -6$ **[B1 for either coordinates of *P* and *Q*]**

$m_{PQ} = \frac{0-(-15)}{-6-0} = -\frac{5}{2}$ **[M1]**

$\frac{dy}{dx} = -\frac{40}{(x+8)^2}$ **[B1]**

Let $-\frac{40}{(x+8)^2} = -\frac{5}{2}$

$(x+8)^2 = 16$

$x+8 = \pm 4$

$x = -12$ (reject) or -4 **[A1 for −12 and not choosing it]**

When $x = -4$, $y = -10$. Therefore $R(-4, -10)$. **[A1]**

Downloaded from SingaKids.com

(b) Given that the point S has coordinates (–2, –5), find the area of the quadrilateral $PRQS$. [2]

$$\text{Area} = \frac{1}{2}\begin{vmatrix} -6 & -4 & 0 & -2 & -6 \\ 0 & -10 & -15 & -5 & 0 \end{vmatrix} \qquad \textbf{[M1]}$$

$$= \frac{1}{2}(60 + 60 - 30 - 30)$$

$$= 30 \text{ units}^2 \qquad \textbf{[A1]}$$

(c) Determine, with reason, whether $PRQS$ is a parallelogram. [3]

$$m_{PR} = \frac{0-(-10)}{-6-(-4)} = -5$$

$$m_{QS} = \frac{-15-(-5)}{0-(-2)} = -5$$

$$m_{RQ} = \frac{-10-(-15)}{-4-0} = -\frac{5}{4}$$

$$m_{PS} = \frac{0-(-5)}{-6-(-2)} = -\frac{5}{4}$$

[M1 for 1 pair of gradient
M2 for 2 pairs of gradient]

Since $m_{PR} = m_{QS}$ and $m_{RQ} = m_{PS}$, $PRQS$ is a parallelogram **[A1]**

OR


$$\text{Midpoint of } PQ = \left(\frac{-6+0}{2}, \frac{0-15}{2}\right) = (-3, -7.5) \qquad \textbf{[M1]}$$

$$\text{Midpoint of } RS = \left(\frac{-4-2}{2}, \frac{-10-5}{2}\right) = (-3, -7.5) \qquad \textbf{[M1]}$$

Since the diagonals bisect each other, $PRQS$ is a parallelogram. **[A1]**

Downloaded from SingaKids.com

8 The diagram shows part of the curve $y = \frac{30}{2x-1}$ and the line $y = -\frac{4}{3}x + \frac{38}{3}$. The curve intersects the line at the points A and B.



(a) Find the coordinates of the points A and B. [3]

$$\frac{30}{2x-1} = -\frac{4}{3}x + \frac{38}{3}$$

$90 = (-4x + 38)(2x - 1)$ **[M1 simplify to quad func]**

$90 = -8x^2 + 4x + 76x - 38$

$0 = 8x^2 - 80x + 128$

$0 = x^2 - 10x + 16$

$0 = (x - 8)(x - 2)$

$x = 8$ or $x = 2$ **[A1]**

Therefore $A(2, 10)$ and $B(8, 2)$. **[A1]**

Downloaded from SingaKids.com

(b) Find the area of the shaded region. [4]

$$\text{Area} = \int_2^8 -\frac{4}{3}x + \frac{38}{3} - \frac{30}{2x-1}\,dx \qquad \textbf{[M1 for correct order of subtraction]}$$

$$= \left[-\frac{2}{3}x^2 + \frac{38}{3}x - 15\ln(2x-1)\right]_2^8 \qquad \begin{bmatrix}\textbf{B1 for 2 terms;}\\ \textbf{B1 for 3rd term}\end{bmatrix}$$

$$= 58\frac{2}{3} - 15\ln 15 - \left(22\frac{2}{3} - 15\ln 3\right)$$

$$= 36 + 15\ln 3 - 15\ln 15$$

$$= 36 - 15\ln 5$$

$$= 11.9 \text{ units}^2 \text{ (to 3 } s.f.\text{)} \qquad \textbf{[A1]}$$

OR

$$\text{Area} = \frac{1}{2}(10+2)(6) - \int_2^8 \frac{30}{2x-1}dx \qquad \begin{bmatrix}\textbf{M1 for correct integral}\\ \textbf{and subtraction}\end{bmatrix}$$

$$= 36 - [15\ln(2x-1)]_2^8 \qquad \begin{bmatrix}\textbf{B1 for 36;}\\ \textbf{B1 for integrated term}\end{bmatrix}$$

$$= 36 - (15\ln 15 - 15\ln 3)$$

$$= 11.9 \text{ units}^2 \text{ (to 3 } s.f.\text{)} \qquad \textbf{[A1]}$$

(c) Show that the area bounded by the curve, the y-axis and the lines $y = 2$ and $y = 6$ can be expressed in the form $15\ln p + q$, where p and q are constants to be determined. [3]

$$y = \frac{30}{2x-1}$$

$$2x - 1 = \frac{30}{y}$$

$$x = \frac{1}{2}\left(\frac{30}{y} + 1\right)$$

$$= \frac{15}{y} + \frac{1}{2}$$

$$\text{Area} = \int_2^6 \frac{15}{y} + \frac{1}{2}\,dy \qquad \textbf{[M1 for making } x \textbf{ the subject to form integral]}$$

$$= \left[15\ln y + \frac{y}{2}\right]_2^6 \qquad \textbf{[M1 for correct integration]}$$

$$= 15\ln 3 + 2 \qquad \textbf{[A1]}$$

Downloaded from SingaKids.com

9 The function f is given by $f(x) = x^3 - x^2 + ax - 1$.

It is given that $\frac{f(x)}{x-1} = Q(x) + \frac{3}{x-1}$, where $Q(x)$ is a quadratic function.

(a) Explain how you can use Remainder Theorem to show that $a = 4$. [3]

Since the **remainder is 3** when f(x) is divided by $x - 1$, **[B1]**

we let $f(1) = 3$ **[M1 for f(1)]**

$1 - 1 + a - 1 = 3$

$a = 4$ (shown) **[A1]**

(b) By long division, divide $f(x)$ by $x^2 + 1$.

Hence, express $\frac{f(x)}{x^2+1}$ in the form $px + q + \frac{rx}{x^2+1}$, where p, q and r are constants to be determined. [2]

$$x^2+1 \overline{)\, x^3 - x^2 + 4x - 1} \quad x - 1$$

$$\begin{array}{r} x^3 \quad + x \\ \hline -x^2 + 3x - 1 \\ -x^2 \quad - 1 \\ \hline 3x \end{array}$$

[M1]

$$\frac{f(x)}{x^2+1} = x - 1 + \frac{3x}{x^2+1}$$ **[A1]**

(c) Find $\frac{d}{dx}\ln(x^2+1)$. [1]

$$\frac{d}{dx}\ln(x^2+1) = \frac{2x}{x^2+1}$$ **[B1]**

(d) Hence, using your results from **part (b) and (c)**, find $\int \frac{f(x)}{x^2+1}\,dx$. [2]

$$\int \frac{f(x)}{x^2+1}\,dx = \int x - 1 + \frac{3x}{x^2+1}\,dx$$

$$= \frac{x^2}{2} - x + \frac{3}{2}\ln(x^2+1) + c$$ **[B1 for ln term, B1 for rest of 3 terms Allow ECF]**

Downloaded from SingaKids.com

10 The equation of a circle, C_1, with centre A, is $x^2 + y^2 - 6x + 8y - 75 = 0$.

(a) Find the radius of the circle and the coordinates of its centre, A. [4]

$(x-3)^2 + (y+4)^2 = 10^2$ **[M1 for $(x-3)^2$ or $(y+4)^2$ seen, M1 for 10^2]**

$A\,(3, -4)$ and radius = 10 units **[A1, A1]**

(b) A second circle, C_2, with centre B, has radius 12 units.
The equation of the perpendicular bisector of AB is $y = -\frac{1}{2}x + 5$.
Find the equation of the second circle. [5]

$m_{AB} = 2$ **[B1]**

Equation of AB is

$y - (-4) = 2(x-3)$

$y = 2x - 10$ **[M1]**

$2x - 10 = -\frac{1}{2}x + 5$

$\frac{5}{2}x = 15$

$x = 6$ **[M1]**

$y = 2$

Let centre of second circle be $(a,\ b)$

$\left(\frac{3+a}{2}, \frac{-4+b}{2}\right) = (6, 2)$ **[M1]**

$a = 9,\ b = 8$


Therefore, equation of circle is $(x-9)^2 + (y-8)^2 = 144$ **[A1]**

(c) P is a point on the perpendicular bisector of AB, where P is **not** the midpoint of A and B.
Justify whether triangle PAB is an isosceles triangle. [1]

Since ***P* is equidistant from *A* and *B***, i.e. $PA = PB$, the triangle is isosceles. **[B1]**

Downloaded from SingaKids.com

11 **(a)** Sketch the graph of $y = e^{-x}$ on the given axes. Label any axial intercepts. [1]



[B1]

(b) The curve $y = f(x)$ is such that $f'(x) = e^{2x+1} + e^{-x}$.

(i) Explain why the curve has no stationary points. [2]

Since $e^{2x+1} > 0$ and $e^{-x} > 0$ for all real values of x, **[B1]**

$f'(x) \neq 0$, therefore the curve has no stationary points. **[B1]**

(ii) The curve passes through the point (ln 4, 8e). Without the use of a calculator, find an expression for $f(x)$. Show all working clearly. [6]

$f(x) = \frac{e^{2x+1}}{2} - e^{-x} + c$ **[B1 for 2 terms, B1 for 3rd term]**

$8e = \frac{e^{2\ln 4+1}}{2} - e^{-\ln 4} + c$

$8e = \frac{e^{\ln 16+1}}{2} - e^{\ln(4)^{-1}} + c$ **[M1 for power law i.e. either ln16 or $\ln(4)^{-1}$ seen]**


$8e = \frac{16e}{2} - \frac{1}{4} + c$ **[M1 for $\times$e, M1 for $-\frac{1}{4}$ seen]**

$c = \frac{1}{4}$

Therefore, $f(x) = \frac{e^{2x+1}}{2} - e^{-x} + \frac{1}{4}$ **[A1]**

Downloaded from SingaKids.com

12 The diagram shows the graph of $y = \mathrm{f}'(x)$.



(a) Write down the range of values of x for which the **function** f decreases as x increases. [1]

$x < -10$ or $x > 5$ **[B1]**

(b) The above function f is given by $\mathrm{f}(x) = 12k^2x - 2x^3 - 3kx^2$, where k is a positive constant. By solving a suitable inequality2, use the answer in part **(a)** to find the value of the constant k. [4]

$\mathrm{f}'(x) = 12k^2 - 6x^2 - 6kx$ **[B1]**

For f to be decreasing,

$12k^2 - 6x^2 - 6kx < 0$

$x^2 + kx - 2k^2 > 0$

$(x + 2k)(x - k) > 0$ **[M1 for correct factorisation or for $\mathrm{f}'(x) < 0$]**

$x < -2k$ or $x > k$ **[A1]**

Therefore $k = 5$ **[A1]**

(c) Explain how you can use the above graph of $y = \mathrm{f}'(x)$ to determine the value of x for which f(x) is a maximum value. [1]

From the graph, $\mathrm{f}'(x)$ changes from positive to negative at x =5, therefore f(x) is a maximum value when x = 5. **[B1]**

END OF PAPER

Downloaded from SingaKids.com