SWISS COTTAGE SECONDARY SCHOOL
SECONDARY FOUR AND FIVE
PRELIMINARY EXAMINATION

O

Name: _____ ( ) Class: _____

ADDITIONAL MATHEMATICS

4049/01

Paper 1

Monday 9 September 2024
2 hours 15 minutes

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 90.

Questions	1	2	3	4
Marks				

This document consists of **18** printed pages and **2** blank pages**.**

Setter: Mr Heng Teng Boon
Vetter: Mdm Zoe Pow

[Turn over

Downloaded from SingaKids.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + ... + \binom{n}{r}a^{n-r}b^r + ... + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1).....(n-r+1)}{r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$
$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$
$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$
$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$
$$\sin 2A = 2 \sin A \cos A$$
$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$
$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc\cos A$$
$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

Answer **all** the questions.

Section A (17 marks)

1 The equation of a curve is $y = 3x^3 + ax^2 + b$, where a and b are constants. If $a > 0$, find, in terms of a and/or b, the range of values of x for which y is increasing. [3]

2 The area of a rectangle is $\left(7 + b\sqrt{2}\right)$ cm^2. Given that the length of the rectangle is $\left(a + 4\sqrt{2}\right)$ cm and the breadth of the rectangle is $\left(5 - \sqrt{2}\right)$ cm, find the value of a and of b. [4]

Downloaded from SingaKids.com


3 The equation of a curve is $y = 2x^2 + 12x + 11$.

(a) Express $2x^2 + 12x + 11$ in the form $a(x+b)^2 + c$ where a, b and c are constants. [2]

(b) Find the range of values of p for which the line $y = px + 11$ intersects the curve at two distinct points. [3]

Downloaded from SingaKids.com

4



The diagram shows a triangle *BCE* whose vertices lie on the circumference of a circle. *AD* is a tangent to the circle at point *C* and *AB* is a tangent to the circle at point *B*. *BED* is a straight line.

(a) Prove that angle ABC + angle $CED = 180^\circ$. [3]

(b) Show that there **does not** exist a circle that passes through points *A*, *B*, *E* and *C*. [2]

Downloaded from SingaKids.com

BLANK PAGE

Downloaded from SingaKids.com

Name: _____ ( )

Class: _____

Questions	5	6	7	8	9	10	11	12	13
Marks									

Section B (73 marks)

5 **(a)** Solve the equation $\log_5 x + 2 = 3\log_x 5$. [5]

(b) Sketch the graph $y = \log_{0.5} x$. [2]

Downloaded from SingaKids.com

6 It is given that $\mathrm{f}(x) = 3\sin\left(\frac{x}{2}\right) + 4$.

(a) State the least and greatest value of $\mathrm{f}(x)$. [2]

(b) State the period of $\mathrm{f}(x)$. [1]

(c) Sketch the graph of $y = \mathrm{f}(x)$ for $0 \le x \le 4\pi$. [2]

(d) By drawing a suitable straight line on the same set of axes as the graph of $y = \mathrm{f}(x)$, state the number of solutions of the equation $\sin\left(\frac{x}{2}\right) = -\frac{x}{4\pi}$ for $0 \le x \le 4\pi$. [2]

Downloaded from SingaKids.com

7 A curve is such that $\frac{\mathrm{d}^2y}{\mathrm{d}x^2} = 3e^{-2x} + \cos 2x$. The curve passes through the point A (0, 3) and has a gradient of 5 at A. Find the equation of the curve. [7]

Downloaded from SingaKids.com

8 **(a)** The expansion of $\left(3x - \frac{2}{x^2}\right)^n$ has a term independent of x. By considering the general term in the expansion, explain why n is a multiple of 3. [3]

(b) It is given that $n = 9$.

Find the value of $\frac{\text{coefficient of term independent of } x}{\text{coefficient of } \frac{1}{x^6}}$. [4]

Downloaded from SingaKids.com

9 **(a)** Express $\dfrac{9x^2-4x+8}{(x-2)(x+1)^2}$ in partial fractions. [5]

(b) Hence, find $\int \dfrac{9x^2-4x+8}{(x-2)(x+1)^2}\,\mathrm{d}x$. [3]

Downloaded from SingaKids.com

10 A particle moves in a straight line such that its displacement, s cm, from a fixed point O is modelled by $s = -3t + e^{\frac{t}{2}}$, where t is the time in seconds since the start of motion.

(a) Show that the particle reaches instantaneous rest at $t = 2\ln 6$. [3]

(b) Explain why the particle passes through O during the first second. [2]

Downloaded from SingaKids.com


(c) Find the total distance travelled by the particle in the interval $t = 0$ to $t = 4$. [3]

Downloaded from SingaKids.com

11 The table shows, to 3 significant figures, the value, \$$C$, in thousands, of a car t years from 1st January 2024.

t	1	2	3	4
C	68.2	58.6	50.7	44.3

(a) On the grid below plot $\ln(C-15)$ against t and draw a straight line graph. The vertical $\ln(C-15)$-axis should start at 3.0 and have a scale of 2 cm to 0.2 units. The horizontal t-axis should have a scale of 2 cm to 1 unit. [3]




Downloaded from SingaKids.com

(b) Use your graph to find the gradient of your straight line and hence express C in the form $C = Ae^{-kt} + 15$, where A and k are constants. [4]

(c) Assuming that the model is still appropriate, find the year for which the value of the car is first below \$35 000. [3]

Downloaded from SingaKids.com

12



The diagram shows a parallelogram with vertices $A(-2,0)$, B, $C(7,7)$ and D. The side AB has equation $y = \frac{1}{2}x + 1$ and the length of $AB = 5\sqrt{5}$ units.

(a) Find the coordinates of B. [4]

Downloaded from SingaKids.com

(b) Prove that *ABCD* is a rectangle. [3]

(c) Calculate the area of *ABCD*. [2]

Downloaded from SingaKids.com

13 The equation of a curve is $y = \frac{6}{(2x-5)^3}$.

(a) Show that for $x > 2.5$, the curve has no stationary points. [3]

Downloaded from SingaKids.com

(b) The normal to the curve at $x = 1$ intersects another curve $36y = x^2 + 90x - 78$ at points A and B. Express the difference of the x-coordinates of A and B in the form $\sqrt{k}$, where k is an integer to be found. [7]

END OF PAPER

Downloaded from SingaKids.com

BLANK PAGE

Downloaded from SingaKids.com