1 It is given that P, Q and R are the angles of a triangle.

(a) Show that $\cos P = -\cos (Q + R)$. [2]

$\cos P = \cos [180° - (Q + R)]$ [M1 – for replacing]

$= \cos 180° \cos (Q + R) + \sin 180° \sin (Q + R)$

$= -\cos (Q + R) + 0$ [A1 – apply addition formula &

$= -\cos (Q + R)$ [evaluate to arrive at result]

Accept supplementary angles:

$\cos P = -\cos (180° - P)$ **M1**

$= -\cos (Q + R)$ **A1**

(b) Given that $Q = 45°$ and $R = 60°$, find $\cos P$ in the form $\frac{1}{4}(\sqrt{a} - \sqrt{b})$, where a and b are integers. [3]

$\cos P = -\cos (45° + 60°)$

$= -\cos 45° \cos 60° + \sin 45° \sin 60°$ [M1 – correct use of formula expansion]

$= -\frac{\sqrt{2}}{2}(\frac{1}{2}) + \frac{\sqrt{2}}{2}(\frac{\sqrt{3}}{2})$ [M1 – correct special angles trigo ratios]

$= \frac{1}{4}(\sqrt{6} - \sqrt{2})$ [A1]

Downloaded from SingaKids.com

2 Baking powder is poured onto a flat surface at a constant rate of 2π cm^3s^{-1} and formed a right circular cone. The radius of the cone is always $\frac{1}{18}$ of its height. Find the rate of change of the radius of the cone after 3 seconds of pouring. [5]

Vol. of cone, $V = \frac{1}{3}\pi r^2(18r)$

$= 6\pi r^3$

$\frac{dV}{dr} = 18\pi r^2$ [B1]

After 3 seconds, $V = 6\pi$

$6\pi r^3 = 6\pi$ [M1 - finding corresponding r]

$r = 1$ [A1]

$\frac{dV}{dt} = \left.\frac{dV}{dr}\right|_{r=1} \times \left.\frac{dr}{dt}\right|_{r=1}$ [M1 - connected rate of change]

$2\pi = 18\pi(1)^2 \times \left.\frac{dr}{dt}\right|_{r=1}$

$\left.\frac{dr}{dt}\right|_{r=1} = \frac{1}{9}$

The rate of change required is $= \frac{1}{9}$ cm/s. [A1 o.e.]

Downloaded from SingaKids.com

3 **(a)** Determine the set of values of m for which the equation $2x^2+4x+2m=6mx-2$ has real roots. [4]

$2x^2+(4-6m)x+2m+2=0$

$b^2-4ac\geq 0$ [M1 - correct Discriminant]

$(4-6m)^2-4(2)(2m+2)\geq 0$

$16-48m+36m^2-8(2m+2)\geq 0$

$36m^2-64m\geq 0$ [M1 – simplification in factors]

$m(9m-16)\geq 0$

$m\leq 0 \quad or \quad m\geq \frac{16}{9}$ [A2, minus 1 mark if inequality sign is wrong due to earlier wrong D sign]

(b) Hence state what can be deduced about the curve $y=2(x+1)^2$ and the line $y=6x-2$. Justify your statement. [2]

$2x^2+4x+2=6x-2$

By comparing with (a), $m=1$ [B1 – correct m value]

When $m=1$, it is not within the set of values of m for which there will be real roots, hence the curve will not meet the line/ the curve will not cut the line. [B1]

Downloaded from SingaKids.com

4 **(a)** Show that $\frac{d}{dx}\left(\ln\left(\cos x\right)\right) = -\tan x$. [2]

$\frac{d}{dx}\left(\ln\left(\cos x\right)\right) = \frac{1}{\cos x} \times \frac{d}{dx}\left(\cos x\right)$ [M1 – show working]

$= \frac{-\sin x}{\cos x} = -\tan x$ [A1 – show fraction]

(b) Differentiate $x \tan x$ with respect to x. [2]

$\frac{d}{dx} x \tan x = x \sec^2 x + \tan x$ [M1 – show product rule]

[A1 – correct ans for both]

(c) Using the results from part **(a)** and **(b)**, find $\int x \sec^2 x \, dx$ and hence show that $\int_0^{\frac{\pi}{4}} x \sec^2 x \, dx = \frac{\pi}{4} - \frac{1}{2}\ln 2$. [4]

From (b), $\int \left(x \sec^2 x + \tan x\right) dx = x \tan x + C$ [M1 – use part (b), 'C' must be seen]

$\int x \sec^2 x \, dx + \int \tan x \, dx = x \tan x + C$

$\int x \sec^2 x \, dx = x \tan x - \int \tan x \, dx + C$ [M1 – proper integration and final correct ans, 'C' must be seen]

$= x \tan x + \ln\left(\cos x\right) + C$

$\int_0^{\frac{\pi}{4}} x \sec^2 x \, dx = \left[x \tan x + \ln\left(\cos x\right)\right]_0^{\frac{\pi}{4}}$ [minus 1 mark if "C" is not seen]

$= \frac{\pi}{4}\tan\frac{\pi}{4} + \ln\left(\cos\frac{\pi}{4}\right) - 0 - \ln 1$

$= \frac{\pi}{4} + \ln\left(\frac{1}{\sqrt{2}}\right)$ [M1 – proper evaluation]

$= \frac{\pi}{4} + \ln 2^{-\frac{1}{2}}$

$= \frac{\pi}{4} - \frac{1}{2}\ln 2$ [A1]

Downloaded from SingaKids.com

5 **(a)** In the expansion of $(2+x)^n$, where n is a positive integer, the coefficient of x^2 is twice the coefficient of x. Find the value of n. [3]

$$(2+x)^n = 2^n + \binom{n}{1}2^{n-1}x + \binom{n}{2}2^{n-2}x^2 + \ldots$$

$$\binom{n}{2}2^{n-2} = 2\binom{n}{1}2^{n-1}$$ [M1 – correct coeff]

$$\frac{n(n-1)}{2}\left(2^{n-2}\right) = 2n\left(2^{n-1}\right)$$

$$\frac{n(n-1)}{2}\left(2^n \times 2^{-2}\right) = 2n\left(2^n \times 2^{-1}\right) \quad (n \neq 0)$$ [M1 – simplify]

$$\frac{n-1}{2(4)} = 1$$

$$n-1 = 8$$

$$n = 9$$ [A1 – with rejection]

(b) Find the value of the term that is independent of x in the expansion of $\left(2x - \frac{1}{4x^4}\right)^{15}$. [4]

general term

$$= \binom{15}{r}(2x)^{15-r}\left(-\frac{1}{4x^4}\right)^r$$

$$= \binom{15}{r}2^{15-r}x^{15-r}\left(-\frac{1}{4}\right)^r x^{-4r}$$ [M1]

$$= \binom{15}{r}2^{15-r}\left(-\frac{1}{4}\right)^r x^{15-5r}$$ [M1 - gather x and let power = 0]

$$15 - 5r = 0$$


$$r = 3$$ [M1 for r value]

$$\text{value} = \binom{15}{3}2^{12}\left(-\frac{1}{4}\right)^3 = -29120$$ [A1]

Max 2 marks if able to gather 15 – 5r for exponent of 'x' and equate to zero with correct r value.

Downloaded from SingaKids.com

6



The diagram shows a circle passing through the points P, Q and R. The point Q lies on the line RB. AB is a tangent to the circle at P. The points S and T lie on PR and PQ respectively. Given that AB is parallel to ST, prove that

(a) triangle PST is similar to triangle PQR, [3]

$\angle SPT = \angle QPR$ (common angle) [M1]

$\angle PST = \angle SPA$ (alt. $\angle$s, AB // ST) **OR** $\angle PTS = \angle TPB$ (alt. $\angle$s, AB // ST)

$= \angle PQR$ (Alt. Segment Thm) $= \angle PRQ$ (Alt. Segment Thm) [M1]

$\therefore$ triangle PST is similar to triangle PQR. [A1]

(2 pairs of corresponding angles are equal)

(b) $PQ \times PT = PR \times PS$, [2]

From above result, $\frac{PQ}{PS} = \frac{PR}{PT}$ [M1]

$\Rightarrow$ $PQ \times PT = PR \times PS$ [A1]

Downloaded from SingaKids.com

(c) Determine if $STQR$ is a cyclic quadrilateral. [4]

$QTS = 180° - \angle PTS$ (adj. $\angle$s on a st. line)

$= 180° - \angle QRS$ (from (a) result) [M1]

So $\angle QTS + \angle QRS = 180°$ [M1]

$\angle RST + \angle RQT = 360° - (\angle QTS + \angle QRS)$ ($\angle$ sum of a quadrilateral)

$= 360° - 180°$


$= 180°$ [M1]

By converse of angles in opposite segment, $STQR$ is a cyclic quadrilateral and all four vertices lie on the circumference of a circle.

[A1 – with correct reason, accept even if no mention of four vertices]

Downloaded from SingaKids.com

7



The diagram shows an isosceles triangle ABC in which $A(-3,0)$, $B\left(\frac{1}{2},3\right)$ and $C(5,2)$. M is the foot of perpendicular from B to AC.

(a) Find the coordinates of M. [1]

$$M = \left(\frac{-3+5}{2}, \frac{0+2}{2}\right) = (1, 1) \qquad \text{[B1]}$$

(b) Find the equation of the perpendicular bisector of AC. [2]

Gradient of $AC = \frac{2-0}{5-(-3)} = \frac{1}{4}$

Gradient of perpendicular BM $= -4$ [M1 – use $m \times m_{\perp} = -1$]

or $m_{BM} = \frac{3-1}{\frac{1}{2}-1} = -4$


Equation of perpendicular bisector:

$y - 1 = -4(x-1)$ [A1]

$y = -4x + 5$

Downloaded from SingaKids.com

(c) Given that $ABCD$ is a kite with $BM = \frac{2}{7}BD$, find the coordinates of D. [3]



Correct ratio: 2 : 5

$\frac{1}{2} : 1\frac{1}{4}$ M1

Coordinates of D

$= \left(1+\frac{1}{4}, 1-5\right)$ M1

$= \left(2\frac{1}{4}, -4\right)$ A1

Alternative: using vectors

$\overrightarrow{BM} = \frac{2}{7}\overrightarrow{BD}$

$\begin{pmatrix}0.5\\-2\end{pmatrix} = \frac{2}{7}\left(\overrightarrow{OD} - \overrightarrow{OB}\right)$ [M1 - position vectors]

$\overrightarrow{OD} = 3.5\begin{pmatrix}0.5\\-2\end{pmatrix} + \begin{pmatrix}0.5\\3\end{pmatrix} = \begin{pmatrix}2.25\\-4\end{pmatrix}$ [M1]

$D = \left(2\frac{1}{4}, -4\right)$ [A1]

(d) Find the area of the kite $ABCD$. [2]

Area of $ABCD = \frac{1}{2}\begin{vmatrix}-3 & 2\frac{1}{4} & 5 & \frac{1}{2} & -3\\ 0 & -4 & 2 & 3 & 0\end{vmatrix}$

M1 (ecf 1 – correct method)


$= \frac{1}{2}(31.5 + 28)$ (anti-clockwise)

$= 29.75$ units^2 $\left(\text{or } \frac{119}{4}\right)$ A1

(accept 29.7 with 3 s.f. if using other method such as Pythagoras'

Downloaded from SingaKids.com

8



A prototype consists of a cylindrical container of height h cm and radius r cm inscribed in a hollow sphere with centre O.

The sphere has a surface area of 6400π cm^2 and both the sphere and container have negligible thickness.

(a) Show that the volume of the cylinder container, V cm^3, is given by [3]

$$V = 2\pi r^2\sqrt{1600 - r^2}\ .$$

$4\pi R^2 = 6400\pi$	[M1 – find radius R]
$R = 40$ (radius R of sphere)	
By Pythagoras' Theorem,	
$r^2 + \left(\frac{h}{2}\right)^2 = 40^2$	[M1]
$h^2 = 4\left(1600 - r^2\right)$	
$h = \sqrt{4\left(1600 - r^2\right)}$	
$h = 2\sqrt{1600 - r^2}$	
$V = \pi r^2 h$	[A1]
$= 2\pi r^2\sqrt{1600 - r^2}$	

Downloaded from SingaKids.com

(b) Given that r can vary, find the value of r for which the volume V is stationary. [5]

$\frac{dV}{dr} = (2\pi r^2) \times \frac{1}{2}(1600 - r^2)^{-\frac{1}{2}} \times (-2r) + \sqrt{1600 - r^2} \times 2 \times 2\pi r$ [M2 – correct differentiation using product rule, deduct M1 if either term is incorrect]

$\frac{dV}{dr} = 4\pi r\sqrt{1600 - r^2} - \frac{2\pi r^3}{\sqrt{1600 - r^2}}$

For stationary values, $\frac{dV}{dr} = 0$

$4\pi r\sqrt{1600 - r^2} - \frac{2\pi r^3}{\sqrt{1600 - r^2}} = 0$ [M1 – equate to zero]

$\frac{2\pi r}{\sqrt{1600 - r^2}}\left[2(1600 - r^2) - r^2\right] = 0$ [M1 – simplify and show this or the next line working]

$\frac{2\pi r}{\sqrt{1600 - r^2}}\left[3200 - 3r^2\right] = 0$

$r = 0$ or $3200 - r^2 = 0$

(reject) $r = \frac{40\sqrt{6}}{3} = 32.7$ cm [A1]

(c) A scientist plans to launch this prototype into outer space carrying as much fuel as possible. Explain whether the prototype can satisfy the scientist's requirement. [2]

x	$\left(\frac{40\sqrt{6}}{3}\right)^-$	$\frac{40\sqrt{6}}{3}$	$\left(\frac{40\sqrt{6}}{3}\right)^+$
$\frac{dV}{dx}$	+	0	–
sketch of tangent	/	—	\

[M1 – first or 2nd derivative, ecf 1 for method]

Since this value of r give maximum V value, the prototype satisfies the scientist's requirement as he could maximise the fuel to be stored in the prototype. [B1]

Downloaded from SingaKids.com

9 It is given that $f(x) = 4 + \cos\left(\frac{x}{2}\right)$ and $g(x) = -2\sin x$.

(a) State the period and amplitude of $f(x)$. [2]

Period = 720° or 4π [B1]


Amplitude = 1 [B1]

(b) State the period and amplitude of $g(x)$. [1]

Period = 360° or 2π

Amplitude = 2 [B1 for both]

(c) Sketch, on the same axes, the graphs of $y = f(x)$ and $y = g(x)$ for $0° \le x \le 360°$. [3]



	f(x)	*g(x)*
Shape	cosine	negative sine
Amplitude	1	2
No. of cycle	Half	1
Shift	4	-

[B1 – for starts and ends at 'zero' for $g(x)$]

[B1 – for starts '5' and ends at '3' for $f(x)$]

[B1 – fully correct graphs]

Downloaded from SingaKids.com

10 **(a)** Express $\dfrac{1-3x-3x^2}{x(x+1)^2}$ in partial fractions. [5]

$$\frac{1-3x-3x^2}{x(x+1)^2}=\frac{A}{x}+\frac{B}{x+1}+\frac{C}{(x+1)^2}$$

$1-3x-3x^2=A(x+1)^2+Bx(x+1)+Cx$ [M1]

When $x=-1$, $C=-1$ [M1]

When $x=0$, $A=1$ [M1]

When $x=1$, $B=-4$ [M1]

$$\frac{1-3x-3x^2}{x(x+1)^2}=\frac{1}{x}-\frac{4}{x+1}-\frac{1}{(x+1)^2}$$ [A1]

(b) Hence find $\int\dfrac{1-3x-3x^2}{2x(x+1)^2}\,dx$. [4]

$$\int\frac{1-3x-3x^2}{2x(x+1)^2}\,dx=\frac{1}{2}\int\frac{1-3x-3x^2}{x(x+1)^2}\,dx$$

$$=\frac{1}{2}\int\frac{1}{x}-\frac{4}{x+1}-\frac{1}{(x+1)^2}\,dx$$ [M1 – take out 0.5]


$$=\frac{1}{2}\ln x-2\ln(x+1)+\frac{1}{2(x+1)}+C$$ [A1 for each term, must have constant C]

Accept $=\ln\sqrt{x}-\ln(x+1)^2+\dfrac{1}{2(x+1)}+C$

or $\ln\dfrac{\sqrt{x}}{(x+1)^2}+\dfrac{1}{2(x+1)}+C$

Downloaded from SingaKids.com

11



The horizontal distance of a child on a carousel, h m, from the starting point is modelled by the equation, $h = 2(1 - \cos kt)$, where k is a constant and t is the time in seconds after the child leaves the starting point. The time to complete one revolution is 20 seconds.

(a) Explain why this model suggests that the diameter of the carousel is 4 m. [1]

$h = 2(1 - \cos kt)$.

Since the diameter of the carousel = max value of h, $h = 2\,(1 - (-1)) = 4$ m

[B1 – must relate diameter to h, *do not accept amplitude method as question ask on the model equation*]

(b) Show that the value of k is $\frac{\pi}{10}$ radians per second. [2]

Period = 20 s

$\frac{2\pi}{k} = 20$ [M1]

$k = \frac{\pi}{10} rad/s$ (*shown*) [A1]

accept
2π rad in 20 s
$\frac{2\pi}{20}$ rad in 1 s
$\frac{2\pi}{20}t$ rad in t s
$k = \frac{\pi}{10}$

accept
$h = 4$ when $t = 10$
to find k value

Downloaded from SingaKids.com

(c) As the carousel turns, it is possible for the child on the carousel to view a landmark, provided that the horizontal distance of the child is within 1 m from the starting point.

Find the duration of time for which the child will not be able to view the landmark during one revolution. [5]

$1 = 2(1-\cos\frac{\pi}{10}t)$ accept $2(1-\cos\frac{\pi}{10}t) > 1$ [M1]

$\frac{1}{2} = \cos\frac{\pi}{10}t$


$basic\angle = \frac{\pi}{3}$ (solution in 1st and 4th quad)

$\frac{\pi}{10}t = \frac{\pi}{3}, \frac{5\pi}{3}$ [M1]

accept $t > \frac{10}{3}$ $t < \frac{50}{3}$


$t = \frac{10}{3}, \frac{50}{3}$ [M1]

Not able to view : $\frac{50}{3} - \frac{10}{3} = \frac{40}{3}s$ *or* $13.3\,s\ (3s.f.)$ [M1, A1]



Downloaded from SingaKids.com

12



The diagram shows part of the curve $y=\frac{10}{4x+1}$ intersecting the y-axis at $A(0, 10)$. The tangent to the curve at the point $P(1, 2)$ intersects the y-axis at B.

(a) Show that the coordinates of B is (0, 3.6). [4]

$y=\frac{10}{4x+1}=10(4x+1)^{-1}$

$\frac{dy}{dx}=-10(4x+1)^{-2}(4)$ [M1]

$\frac{dy}{dx}=-40(4x+1)^{-2}$

When $x=1$ $\frac{dy}{dx}=-40(4(1)+1)^{-2}$

$\frac{dy}{dx}=-1.6$ [M1]

$\frac{y-2}{0-1}=-1.6$ [M1]

$y-2=1.6$

$y=3.6$

Coordinate of B is (0, 3.6) [A1]

Downloaded from SingaKids.com

(b) Find the **exact** area of the shaded region. [5]

$$Area = \int_0^1 \frac{10}{4x+1}\,dx - \frac{1}{2}(3.6+2)(1) \qquad \text{[M1], [M1]}$$

$$Area = \left[\frac{10\ln(4x+1)}{4}\right]_0^1 - 2.8 \qquad \text{[M1]}$$

$$Area = \left[\frac{10\ln(4+1)}{4} - \frac{10\ln(1)}{4}\right] - 2.8 \qquad \text{[M1]}$$

$$Area = \frac{5}{2}\ln 5 - 2.8 \text{ unit}^2 \qquad \text{[A1]}$$

END OF PAPER

Downloaded from SingaKids.com

Downloaded from SingaKids.com

1 A curve has the equation $y = \dfrac{\sin 2x}{2 - \cos 2x}$.

(a) Show that the gradient function can be expressed in the form $\dfrac{k\cos 2x - 2}{(2 - \cos 2x)^2}$, where k is a constant. [3]

$$y = \frac{\sin 2x}{2 - \cos 2x}$$

$$\frac{dy}{dx} = \frac{(2 - \cos 2x)2\cos 2x - \sin 2x(2\sin 2x)}{(2 - \cos 2x)^2}$$

M1: correct quotient/pdt rule

M1: differentiate sin 2x and cos 2x correctly

$$= \frac{4\cos 2x - 2\cos^2 2x - 2\sin^2 2x}{(2 - \cos 2x)^2}$$

$$= \frac{4\cos 2x - 2(\cos^2 2x + \sin^2 2x)}{(2 - \cos 2x)^2}$$

A1: use of identity to reach answer

$$= \frac{4\cos 2x - 2}{(2 - \cos 2x)^2}$$

(b) Find the acute angle between the tangent to the curve at $x = \dfrac{\pi}{12}$ and the line $y = 0$. [3]

$$\text{gradient of tangent} = \frac{4\cos\frac{\pi}{6} - 2}{\left(2 - \cos\frac{\pi}{6}\right)^2} = 1.1386$$

M1: correct gradient value

$$\text{Angle required} = \tan^{-1}(1.1386) = 48.7° \text{ (1 dp)}$$

M1, A1 (accepts 0.850 rad)

Downloaded from SingaKids.com

2 **(a)** Factorise x^3+27k^3 as a product of a linear and a quadratic factor. [2]

$x^3+27k^3=(x+3k)(x^2-3kx+9k^2)$

M1: $(x+3k)$

A1

(b) Hence solve $x^3+27=(x+3)(x+10)$, expressing non-integer roots in surd form. [3]

$k=1: x^3+27=(x+3)(x^2-3x+9)$

$(x+3)(x^2-3x+9)=(x+3)(x+10)$

$x+3=0$ or $x^2-3x+9=x+10$

$x=-3$ or $x^2-4x-1=0$

$$x=\frac{4\pm\sqrt{16-4(-1)}}{2}$$

$$=2\pm\sqrt{5}$$

B1: identify x=-3 as a root from Hence

M1: apply quad formula correctly

A1

(c) Find the value of k given that x^3+27k^3 leaves a remainder of 351 when divided by $x-2$. [2]

Let $f(x)=x^3+27k^3$

$f(2)=2^3+27k^3=351$


$$k=\sqrt[3]{\frac{351-8}{27}}=\frac{7}{3}$$

M1: applying remainder thm correctly


A1

Downloaded from SingaKids.com

3 The diagram shows a L-shaped rod ABC where AB and BC have length 18 m and 8 m respectively and angle ABC is 90°. The rod is hinged to a wall at A so as to rotate in a vertical plane. The rod AB makes an acute angle θ with the vertical wall surface OA.



(a) Given that G is a point directly below C, show that $OG = p\cos\theta + q\sin\theta$, where p and q are constants to be found. [2]



Using triangle BCD: Angle $DBC = \theta$

$$\cos\theta = \frac{BD}{8}$$

$$BD = 8\cos\theta$$

B1: either correctly establishing BD or BE

Using triangle ABE: Angle $BAE = \theta$

$$\sin\theta = \frac{BE}{18}$$

$$BE = 18\sin\theta$$

$OG = BD + BE$

$= 8\cos\theta + 18\sin\theta$, $p = 8$, $q = 18$

B1: showing the other component and clear indication that OG is a sum of the 2 values

(b) Express OG in the form $R\cos(\theta - \alpha)$ where $R > 0$ and $0° \le \alpha \le 90°$. [3]

Downloaded from SingaKids.com

$OG = 8\cos\theta + 18\sin\theta$

$= \sqrt{8^2+18^2}\cos\left(\theta - \tan^{-1}\frac{18}{8}\right)$

$= \sqrt{388}\cos\left(\theta - \tan^{-1}\frac{9}{4}\right)$

$= 19.7\cos(\theta - 66.0°)$

M1: $\sqrt{8^2+18^2}$

M1: $\tan^{-1}\frac{18}{8}$

A1: correct evaluation and form, accept $\sqrt{388}$

(c) Find the length of OG and the corresponding value of θ if G is at maximum displacement from O. [3]

max $OG = \sqrt{388}$

$= 19.7$ m

when $\cos\left(\theta - \tan^{-1}\frac{9}{4}\right) = 1$

$\theta - \tan^{-1}\frac{9}{4} = 0$

$\theta = \tan^{-1}\frac{9}{4} = 66.0°$

A1: $\sqrt{388}$ or $19.7m$

M1: $\theta - \tan^{-1}\frac{9}{4} = 0$

A1: 66.0°

Downloaded from SingaKids.com

4 A circle C_1 has equation $x^2+y^2-6x+4y=12$.

(a) Find the radius and the coordinates of the centre of C_1. [3]

$x^2-6x+y^2+4y=12$

$(x-3)^2+(y+2)^2=12+3^3+2^2$ — M1: completing the square or using formula

$(x-3)^2+(y+2)^2=5^2$

Radius = 5 units — A1

Centre = (3, -2) — B1

(b) Find the equation of the tangent to the circle at the point $P(7,-5)$. [3]

Gradient of normal $=\dfrac{-2-(-5)}{3-7}=-\dfrac{3}{4}$ — M1

Gradient of tangent $=\dfrac{4}{3}$


Eqn of tangent: $y+5=\dfrac{4}{3}(x-7)$ — M1 award for correct pts and their -1/m used

$y=\dfrac{4}{3}x-\dfrac{43}{3}$ — A1, or $3y=4x-43$

(c) Another circle C_2 has centre $(-8, 4)$ and radius 7 cm. Find the shortest distance between the 2 circles. [2]

Distance between centres of circle $=\sqrt{(3+8)^2+(-2-4)^2}=\sqrt{157}$ — M1

Shortest distance $=\sqrt{157}-7-5=0.530$ cm (3 s.f.) — A1



Downloaded from SingaKids.com

5 **(a)** Prove the identity $\dfrac{(\sin A-\cos A)(1+\sin A\cos A)}{\cos^3 A}=\tan^3 A-1$. [4]

$$LHS=\frac{(\sin A-\cos A)(1+\sin A\cos A)}{\cos^3 A}$$

$$=\frac{\sin A-\cos A+\sin^2 A\cos A-\sin A\cos^2 A}{\cos^3 A}$$ M1: correct expansion of terms

$$=\frac{\sin A-\cos A+(1-\cos^2 A)\cos A-\sin A(1-\sin^2 A)}{\cos^3 A}$$ M1: applying identity

$$=\frac{\sin A-\cos A+\cos A-\cos^3 A-\sin A+\sin^3 A}{\cos^3 A}$$

$$=\frac{\sin^3 A-\cos^3 A}{\cos^3 A}$$ M1: expand & simplify

$$=\frac{\sin^3 A}{\cos^3 A}-\frac{\cos^3 A}{\cos^3 A}$$ A1: manipulation to RHS

$$=\tan^3 A-1$$

OR

$$LHS=\frac{(\sin A-\cos A)(1+\sin A\cos A)}{\cos^3 A}$$

$$=\frac{(\sin A-\cos A)(\sin^2 A+\cos^2 A+\sin A\cos A)}{\cos^3 A}$$ M1: applying identity

$$=\frac{\sin^3 A+\sin A\cos^2 A+\sin^2 A\cos A-\sin^2 A\cos A-\cos^3 A-\sin A\cos^2 A}{\cos^3 A}$$ M1: correct expansion of terms

$$=\frac{\sin^3 A-\cos^3 A}{\cos^3 A}$$ M1: simplify

$$=\frac{\sin^3 A}{\cos^3 A}-\frac{\cos^3 A}{\cos^3 A}$$ A1: manipulation to RHS

$$=\tan^3 A-1$$

Downloaded from SingaKids.com

(b) Hence solve $(\sin A - \cos A)(1 + \sin A \cos A) + 2\cos^3 A = 0$ exactly, for $-\pi \le A \le \pi$ radians. [4]

$(\sin A - \cos A)(1 + \sin A \cos A) = -2\cos^3 A$

$\dfrac{(\sin A - \cos A)(1 + \sin A \cos A)}{\cos^3 A} = -2$

$\tan^3 A - 1 = -2$ M1: simplification to single trigo equation

$\tan^3 A = -1$

$\tan A = -1$


Basic angle $= \dfrac{\pi}{4}$ M1: correct basic angle – must be acute

Quad: Q2, Q4

$A = \dfrac{3\pi}{4}, -\dfrac{\pi}{4}$ A1, A1

Downloaded from SingaKids.com

6 The diagram shows the graph of $y = x^2 + ax + 4$ and $y = x^2 + bx + c$. The graph of $y = x^2 + ax + 4$ touches the x-axis at P. Points M and O are the x-intercepts of the graph of $y = x^2 + bx + c$. The origin O is the mid-point of MP.



(a) Find the values of a, b and c. [4]

Since origin (0, 0) is on $y = x^2 + bx + c$, $c = 0$ B1

Discriminant of $y = x^2 + ax + 4 = 0$ since curve intersects x-axis once only:

$a^2 - 4(1)(4) = 0$

$a = \pm 4$

Since P is on positive x-axis, $a < 0$: $a = -4$

M1: using discriminant or the equation must be a perfect square

A1

OR

$$y = x^2 + ax + 4 = \left(x + \frac{a}{2}\right)^2 - \frac{a^2}{4} + 4.$$

Since curve intersects x-axis once only:

$$4 - \frac{a^2}{4} = 0$$

$a = \pm 4$

Since P is on positive x-axis, $a < 0$: $a = -4$

M1: using discriminant or the equation must be a perfect square

A1

Coor of P $= \left(-\frac{a}{2}, 0\right) = (2, 0)$

Coor of M $= (-b, 0) = (-2, 0)$

$b = 2$

B1

Downloaded from SingaKids.com

(b) The graph of $y=x^2+ax+4$ and $y=x^2+bx+c$ intersects at N. Find the coordinates of N. [2]

$x^2-4x+4=x^2+2x$

$6x=4$

$x=\frac{2}{3}$

M1: equating and solving correct x (allows ecf)

$y=\left(\frac{2}{3}\right)^2+2\left(\frac{2}{3}\right)=\frac{16}{9}$

Coordinates of $N=\left(\frac{2}{3},\frac{16}{9}\right)$

A1

(c) The graph $y=px^2+qx+r$ has its turning point at N and passes through point P. Find the values of p, q and r, where $r>0$. [3]

Graph with turning point at N and passes P (downward opening):

$y=-p\left(x-\frac{2}{3}\right)^2+\frac{16}{9}$, $p<0$

M1: correct completed square form

At P $(2,0)$, $0=-p\left(2-\frac{2}{3}\right)^2+\frac{16}{9}$

$p=-1$

$y=-\left(x-\frac{2}{3}\right)^2+\frac{16}{9}$

$=-\left(x^2-\frac{4}{3}x+\frac{4}{9}\right)+\frac{16}{9}$

$=-x^2+\frac{4}{3}x+\frac{4}{3}$

A1: correct p

A1: correct q & r

$p=-1,\ q=r=\frac{4}{3}$

OR (longer method): form 3 equations with coor of N, P and either 2nd x-intercept or derivative and solve

$y=px^2+qx+r$

$\frac{dy}{dx}=2px+q$

Turning point at $\left(\frac{2}{3},\frac{16}{9}\right)$: $2p\left(\frac{2}{3}\right)+q=0$

$4p=-3q$ ---- (1)

M1: establishing 3 equations correctly

Graph passes through P: $0=p(2)^2+q(2)+r$

$4p+2q+r=0$

Downloaded from SingaKids.com

Subst (1): $q = r$ ----- (2)

Graph passes through N: $\frac{16}{9} = p\left(\frac{2}{3}\right)^2 + q\left(\frac{2}{3}\right) + r$

$4p + 6q + 9r = 16$

Subst (1) and (2):

$-3q + 6q + 9q = 16$

$12q = 16$

$q = \frac{4}{3}$

A1: correct p

A1: correct q & r

7 A particle P, travels in a straight line, so that its displacement, s m, from O at time t seconds, is modelled by $s = \frac{1}{3}t^3 - 5t^2 - 3$.

(a) Find the value of t when particle P return to its initial position. [2]

Initial position when $t = 0$, $s = -3$

$\frac{1}{3}t^3 - 5t^2 - 3 = -3$ M1: setting s = -3

$\frac{1}{3}t^3 - 5t^2 = 0$

$t^2\left(\frac{1}{3}t - 5\right) = 0$

$t = 0$ or $t = 15s$

Particle returns to initial position at t = 15s. A1

(b) Find the minimum velocity of particle P. [3]

$v = t^2 - 10t$

$= (t-5)^2 - 25$

Minimum velocity = -25 m/s^2

M1: $v = t^2 - 10t$

M1: completing the square

A1

OR

$v = t^2 - 10t$

At stationary v,

$\frac{dv}{dt} = 2t - 10 = 0$

$t = 5$

$v = (5)^2 - 10(5) = -25$ m/s

M1: $v = t^2 - 10t$

M1: solving for t = 5

$\frac{d^2v}{dt^2} = 2 > 0$, minimum v at -25m/s

A1: includes check for minimum

Downloaded from SingaKids.com

(c) Another particle, Q, travels in a straight line from O such that its velocity, v m/s, at time t seconds after passing O is given by $v = 24\left(e^{-\frac{t}{6}} - e^{-1}\right)$.

Find the value of t at which the particle Q is instantaneously at rest. [2]

$v = 24\left(e^{-\frac{t}{6}} - e^{-1}\right) = 0$

$e^{-\frac{t}{6}} = e^{-1}$

$t = 6s$

M1: Setting $v = 0$

A1

(d) Find the total distance travelled by particle Q for the first 9 seconds. [4]

$s = 24\left(\dfrac{e^{-\frac{t}{6}}}{-\frac{1}{6}} - te^{-1}\right) = -24\left(6e^{-\frac{t}{6}} + te^{-1}\right) + C$

M1: correct integration (award even if without +C)

When t=0, s=0: $s = -24(6) + C = 0$

$C = 144$

$s = 144 - 24\left(6e^{-\frac{t}{6}} + te^{-1}\right)$

M1: correct S

When $t = 6$, $s = 144 - 24\left(6e^{-1} + 6e^{-1}\right) = 38.051m$ (5 sf)

When $t = 9$, $s = 144 - 24\left(6e^{-\frac{9}{6}} + 9e^{-1}\right) = 32.407m$ (5 sf)

M1: sub t= 6 and t=9

Total distance travelled = $s = 38.051 + (38.051 - 32.407) = 43.7m$ (3 sf)

A1

OR

Displacement covered for 1st 6 sec =

$\int_0^6 24\left(e^{-\frac{t}{6}} - e^{-1}\right) dt = 24\left[\dfrac{e^{-\frac{t}{6}}}{-\frac{1}{6}} - te^{-1}\right]_0^6$

M1: correct integration

$= -24\left[6e^{-\frac{t}{6}} + te^{-1}\right]_0^6$

$= -24\left[6e^{-\frac{6}{6}} + 6e^{-1} - 6e^{-\frac{0}{6}} - 0\right]$

$= 38.051m$ (3sf)

M1

Displacement covered for next 3 sec =

Downloaded from SingaKids.com

$$\int_6^9 24\left(e^{-\frac{t}{6}}-e^{-1}\right)\mathrm{d}t=-24\left[6e^{-\frac{t}{6}}+te^{-1}\right]_6^9$$

$$=-24\left[6e^{-\frac{9}{6}}+9e^{-1}-6e^{-\frac{6}{6}}-6e^{-\frac{6}{6}}\right]$$

$$=-5.6434m \text{ (5sf)}$$

M1

Total distance $= s=38.051+(38.051-32.407)=43.7m$

A1

8 The height, h cm, of a plant is modelled by $h=\dfrac{80}{1+10\mathrm{e}^{-0.4t}}$, where t is the number of months after the first observation.

(a) Show that h is an increasing function. [3]

$$\frac{\mathrm{d}h}{\mathrm{d}t}=80\frac{\mathrm{d}}{\mathrm{d}t}\left(1+10\mathrm{e}^{-0.4t}\right)^{-1}$$

$$=80(-1)\left(1+10\mathrm{e}^{-0.4t}\right)^{-2}\left(10(-0.4)\mathrm{e}^{-0.4t}\right)$$

M1: differentiate $1+10\mathrm{e}^{-0.4t}$ correctly

$$=\frac{320\mathrm{e}^{-0.4t}}{\left(1+10\mathrm{e}^{-0.4t}\right)^2}$$

M1: simplification

Since $\mathrm{e}^{-0.4t}>0$ for $t>0$, $320\mathrm{e}^{-0.4t}>0$ and $\left(1+10\mathrm{e}^{-0.4t}\right)^2>0$

$$\frac{\mathrm{d}h}{\mathrm{d}t}=\frac{320\mathrm{e}^{-0.4t}}{\left(1+10\mathrm{e}^{-0.4t}\right)^2}>0$$

Therefore, h is an increasing function.

A1: Clear explanation on why derivative is positive and therefore h is increasing

(b) Find the value of t when the height of the plant first exceeds four times its initial observation. [3]

Since h is increasing,

$$\frac{80}{1+10\mathrm{e}^{-0.4t}}>\frac{4\times80}{1+10\mathrm{e}^{0}}$$

M1: correct set up of equation (accepts equal if students round up at the end, preferably with explanation that h is an increasing function)

$$1+10\mathrm{e}^{-0.4t}<\frac{11}{4}$$

$$\mathrm{e}^{-0.4t}<\frac{\frac{11}{4}-1}{10}=0.175$$

M1: simplification to exponential eqn

$$-0.4t<\ln 0.175$$

$$t>4.3574 \text{ (5 s.f.)}$$

$$t=4.36 \text{ (3 s.f.)}$$

A1 (accepts 5, though context need not be integer)

Downloaded from SingaKids.com

(c) The height, y cm, of another species of plant, t months after the first observation is given by $y=\frac{1}{a+be^{-t}}$, where a and b are constants. Explain clearly how a straight line graph can be drawn to represent this relationship. You should state which variables should be plotted on each axis and explain how the values of a and b can be calculated. [4]

$y=\frac{1}{a+be^{-t}}$

$\frac{1}{y}=a+be^{-t}$ M1: simplification to Y=mX + c form

Plot $\frac{1}{y}$ on the vertical axis against e^{-t} on the horizontal axis to obtain a straight line graph. [A1] The gradient of the graph would give the value of b [A1] and the y-intercept will give the value of a [A1].

Note: no award of full A1s if M1 not awarded.

9 **(a)** Without using a calculator, solve the equation $x\sqrt{15}+\sqrt{5}=x\sqrt{2}+\sqrt{6}$. Leave your answer in the form $p\sqrt{10}+q\sqrt{3}$, where p and q are fractions. [4]

$x\sqrt{15}+\sqrt{5}=x\sqrt{2}+\sqrt{6}$

$x\left(\sqrt{15}-\sqrt{2}\right)=\sqrt{6}-\sqrt{5}$

$x=\frac{\sqrt{6}-\sqrt{5}}{\sqrt{15}-\sqrt{2}}$ M1: $x=\frac{\sqrt{6}-\sqrt{5}}{\sqrt{15}-\sqrt{2}}$

$=\frac{\sqrt{6}-\sqrt{5}}{\sqrt{15}-\sqrt{2}}\times\frac{\sqrt{15}+\sqrt{2}}{\sqrt{15}+\sqrt{2}}$ M1: correct rationalisation fraction

$=\frac{\sqrt{90}-\sqrt{10}-\sqrt{75}+\sqrt{12}}{15-2}$

$=\frac{3\sqrt{10}-\sqrt{10}-5\sqrt{3}+2\sqrt{3}}{13}$ M1: expansion of numerator and denominator

$=\frac{2\sqrt{10}-3\sqrt{3}}{13}$

$=\frac{2}{13}\sqrt{10}-\frac{3}{13}\sqrt{3}$ A1: in $p\sqrt{10}+q\sqrt{3}$, where p and q are fractions

Downloaded from SingaKids.com

(b) Without using a calculator, solve the equation $\log_2 x - \log_x 16 = 0$. [4]

$\log_2 x - \log_x 16 = 0$

$\log_2 x - \dfrac{\log_2 16}{\log_2 x} = 0$ — M1: correct change of base

$(\log_2 x)^2 = 4$

$\log_2 x = \pm 2$ — M1: simplification of $\log_2 16$ and reduced to a solvable quadratic equation

$x = 2^2$ or $x = 2^{-2}$

$x = 4$ or $x = \dfrac{1}{4}$ — A1, A1

(c) Solve the equation $3^{x+2} - 2\left(3^{-x}\right) = 17$. [4]

$3^{x+2} - 2\left(3^{-x}\right) = 17$

$9\left(3^x\right) - \dfrac{2}{3^x} = 17$

$9\left(3^x\right)^2 - 17\left(3^x\right) - 2 = 0$ — M1: correct application of indices rules and rewriting into quadratic eqn in 3^x

Let $u = 3^x$

$9u^2 - 17u - 2 = 0$

$(u-2)(9u+1) = 0$ — M1: factoring or formula to solve quadratic eqn


$u = 2$ or $u = -\dfrac{1}{9}$ (rej since $3^x > 0$) — A1: Solving of 3^x with rejection

$3^x = 2$

$x = \dfrac{\ln 2}{\ln 3} = 0.631$ (3 s.f.) — A1: do not accept $x = \log_3 2$ as no way of computing

Downloaded from SingaKids.com

10 **(a)** The diagram shows part of a straight line graph which passes through $(2,1)$ and $(4,3)$.



Find the equation of the straight line in the form $y=\dfrac{x}{a+b\sqrt{x}}$, where a and b are constants. [3]

gradient of straight line: $\dfrac{3-1}{4-2}=1$ — M1: gradient of line

equation:

$$Y-1=1(X-2)$$

$$Y=X-1$$ — M1: eqn of line

$$\frac{x}{y}=\sqrt{x}-1$$

$$y=\frac{x}{\sqrt{x}-1}$$ — A1

(b) The table below shows the experimental values of two variables x and y.

x	1	2	3	4	5	6
y	63	127	258	510	1000	2100

It is known that x and y are related by an equation of the form $y=\dfrac{b^x}{10^a}$, where a and b are constants.

(i) On the grid next page, plot $\lg y$ against x and draw a straight line graph. [3]

(ii) Use your graph to estimate the value of a and of b. [3]

$y=\dfrac{b^x}{10^a} \Rightarrow$

Downloaded from SingaKids.com

$\lg y = \lg\left(\frac{b^x}{10^a}\right)$

$\lg y = \lg b^x - \lg 10^a$

$\lg y = x\lg b - a$ M1

$-a = 1.5$

$a = -1.5$ A1 (accept -1.54 to -1.46)

$\lg b = 0.3$

$b = 2.00$ A1 (accept 1.82 to 2.19)

(iii) Explain how you would use the graph to find the value of x for which $(10b)^x = 10^{a+1}$. [2]

$(10b)^x = 10^{a+1}$

$10^x b^x = 10^a \times 10$

$\frac{b^x}{10^a} = \frac{10}{10^x}$ M1

$y = 10^{1-x}$

$\lg y = \lg\left(10^{1-x}\right)$

$\lg y = 1 - x$

Draw the line $\lg y = 1 - x$ and find the x-coordinate of the point of intersection. A1

OR

$(10b)^x = 10^{a+1}$


$x\lg(10b) = (a+1)\lg(10)$

$x\lg(10) + x\lg(b) = a+1$

$x\lg(b) - a = x+1$

$\lg y = x+1$

Downloaded from SingaKids.com



Downloaded from SingaKids.com

Downloaded from SingaKids.com