DUNMAN SECONDARY SCHOOL

CANDIDATE NAME			
CLASS		**INDEX NUMBER**	

PRELIMINARY EXAMINATION 2025
SECONDARY 4 EXPRESS

ADDITIONAL MATHEMATICS 4049/02

Paper 2

25 August 2025

2 hours 15 minutes

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Calculator Model

Write your name, class and index number on all the work you hand in.
Write in dark blue or black pen.
You may use a pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total marks for this paper is 90.

Total Marks

Downloaded from SingaKids.com

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + ... + \binom{n}{r}a^{n-r}b^r + ... + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)...(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2\sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2\tan A}{1 - \tan^2 A}$$

Formula for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

Downloaded from SingaKids.com

1 **(a)** A ball is thrown vertically upwards. Its height h m, above the ground at time t seconds after being thrown is given by the formula $h = 7 + 20t - 5t^2$.
Find the maximum height attained by the ball and the time at which this occurs. [4]

(b) Find the set of values of the constant k for which the curve $y = x^2 + 6x + k$ lies entirely above the line $y = k(x - 1)$. [4]

Downloaded from SingaKids.com

2 A and B are acute angles, such that $\cos A = \frac{3}{5}$ and $\tan B = \frac{12}{5}$.

Find the exact values of


(a) $\tan 2A$, [2]

(b) $\sin\left(B + \frac{3\pi}{2}\right)$, [3]

(c) $\cos\frac{B}{2}$. [3]

Downloaded from SingaKids.com

3 In the diagram, PQS lies on a circle and $PCBQ$ lies on another circle.
PQ is a common chord of the two circles. BPS is a straight line.
QC is a tangent to the circle PQS at Q. QC and BP intersect at M.



(a) Prove that BC is parallel to QS. [3]

(b) Prove that triangle PQM is similar to triangle QSM. [2]

(c) Hence show that $QM^2 = PM \times SM$. [2]

Downloaded from SingaKids.com

4 **(a)** Show that $\frac{\mathrm{d}}{\mathrm{d}x}\left(\cot x\right) = -\operatorname{cosec}^2 x$. [4]

(b) Hence find the value of $\int_{\frac{\pi}{4}}^{\frac{\pi}{3}} -\cot^2 x \, \mathrm{d}x$. [4]


Downloaded from SingaKids.com

5 **(a)** Given that the constant term in the binomial expansion of $\left(x-\frac{k}{x^3}\right)^8$ is 7, find the value of the positive constant k. [4]

(b) Using the value of k found in part **(a)**, show that there is no constant term in the expansion of $\left(1+x^4\right)\left(x-\frac{k}{x^3}\right)^8$. [4]

Downloaded from SingaKids.com

6 The diagram below shows an isosceles triangle ABC, where $AB = AC = 2$ cm, and a square $BCDE$. M is the midpoint of BC.
Angle $BAM = \theta$ and can vary.



(a) Show that the total area of $ABEDC$ is $16 \sin^2 \theta + 4 \sin \theta \cos \theta$ cm^2. [3]

(b) Show that $16 \sin^2 \theta + 4 \sin \theta \cos \theta = 2 \sin 2\theta - 8 \cos 2\theta + 8$. [2]

Downloaded from SingaKids.com

(c) Express $2 \sin 2\theta - 8 \cos 2\theta$ in the form $R \sin (2\theta - \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. [3]

Downloaded from SingaKids.com

7 **(a)** Prove the identity $\dfrac{\cos^2 x - \sin^2 x}{1 + 2\sin x \cos x} = \dfrac{1 - \tan x}{1 + \tan x}$. [4]

(b) Solve the equation $\dfrac{\cos^2 x - \sin^2 x}{1 + 2\sin x \cos x} = \dfrac{2}{3}\tan x$ for $0 \le x \le 2\pi$. [4]

Downloaded from SingaKids.com


8 **(a)** Using long division, divide $2x^3 - 7x^2 + 2x + 3$ by $2x - 4$. [2]

(b) Solve $2x^3 - 7x^2 + 2x + 3 = 0$. [4]

(c) Kun Ye claims that the solutions to the equation $2x^3 - 7x^2 + 2x + 3 = 0$ can be used to solve $3y^6 + 2y^4 - 7y^2 + 2 = 0$. Explain how he is correct. [2]

Downloaded from SingaKids.com

9 The diagram shows part of the curve $y = 3x^2$.
The points P and Q lie on the curve, and have the same y-coordinates.



(a) Show that the area of the triangle OPQ, A cm^2, is given by $A = 3x^3$. [2]

Downloaded from SingaKids.com

The points P and Q move along the curve such that their y-coordinates are changing at a rate of 3 units per second.

Find, at the point where $x = 4$,

(b) the rate of change of PQ, [4]

(c) the rate of change of A. [3]

Downloaded from SingaKids.com

10 The lines $y = 9$, $y = -1$ and $3y = 4x + 9$ are tangents to a circle.

The x-coordinate of the centre of the circle is positive.

(a) Explain why the y-coordinate of the centre of the circle is 4. [1]


(b) By considering the discriminant, or otherwise, show that the x-coordinate of the centre of the circle is 7. [7]

Downloaded from SingaKids.com

(c) Determine if the point $(3,0)$ lies within the circle. [2]

Downloaded from SingaKids.com

11 The diagram shows a lake (shaded) bordered by a straight level road AXB.
Point C is on the surface of the lake and B is the point closest to C.
The distance AB is 30 m and BC is 20 m.



Natalie is at point A on the road and wants to get to point C as quickly as possible. She can run at a speed of 4 m/s and swim at a speed of 2 m/s.
She realizes that in order to minimize the total time to get to C, it is necessary to leave the road at some point X between A and B, and then swim directly to C.

Let AX be x m and T be the total time, in seconds, taken by Natalie to get from A to X and then from X to C.

(a) Show that $T = \frac{x}{4} + \frac{1}{2}\sqrt{x^2 - 60x + 1300}$. [2]

Downloaded from SingaKids.com

(b) Find an expression for $\frac{\mathrm{d}T}{\mathrm{d}x}$. [2]

(c) Find the value of x for which the total time taken is minimum. You are **not** required to justify that this value of x leads to the minimum total time. [4]

END OF PAPER

Downloaded from SingaKids.com

BLANK PAGE

Downloaded from SingaKids.com